

Visit us at https://onlinejima.com





Chief Patron Past President, IMA, WMA, MCI Dr Ketan Desai

Hony. Joint Secretary

Hony. Joint Secretary

from Tamilnadu Dr M Thiraviam Mohan







National Vice President Dr. Suresh Gutta





Imm.Past National President Dr Sharad Kumar Agarwal



National Vice President Dr Shiv Kumar Utture







Hony. Joint Secretary nominated by NP Dr A.V. Jayakrishnan



National Vice President

Dr R Gunasekaran

Hony. Asstt. Secretary from NCR Dr Thakur Padmanabhan



Hony. Asst. Secretary from rest of the country Dr Paramjit Singh Maan

from rest of the country Dr Satish joshi



Hony.Jt Finance Secretary from rest of the country Dr Mahendra Nath Thareja



Hony. Joint Finance Secretary stationed at Calcutta Dr Sarbari Dutta

IMA COLLEGE OF GENERAL PRACTITIONERS



Hony. Joint Secretary from NCR

Dr Satyajit Borah



Hony.Jt.Secretary from Tamilnadu Dr D Senthil Kumar



Dr Poonam Singh



Hony, Joint, Secretary from rest of the country Dr Sunil Bhikhabhai Chenwala



Hony. Secretary Dr. R Anburajan



Hony. Joint. Secretary from rest of the country Dr Yeshwant Vasantrao Gade



Hony.Joint.Secretary. from rest of the country Dr Pavankumar N Patil



04

Hony. Secretary General Dr Anilkumar J Nayak

Journal of the Indian Medical Association

TEAM IMA 2024





Hony. Joint Secretary stationed at Calcutta

















TEAM IMA 2024



Hony. Editor Dr Sanjoy Banerjee



Dr Ranjan Bhattacharyya





Hony. Associate Editor Dr Prasanta Kr. Bhattacharyya



Hony. Secretary Dr Sibabrata Banerjee



05

Hony. Asstt. Secretary Dr Minakshi Ganguly



Chairman

Dr Nomeeta Shiv Gupta Dr Nibedita Pani







Hony. Joint Secretary

Dr Hiren S. Kothari

OF MEDICAL SPECIALITIES



Hony Editor Annals

Dr Shilpa Basu Roy

Hony. Exe Editor Annals

Dr Rajiv Ranjan Prasad

IMA AKN SINHA INSTITUTE OF CONTINUING MEDICAL AND HEALTH EDUCATION & RESEARCH

Hony. Joint Secretary

Dr D. Shekhar Reddy



Vice Chairman

Director Dr Ramneek Singh Bedi

Hony. Editor

Dr Kakoli Sen Mandal



Hony. Secretary

Dr Srirang Abkari

Hony. Executive Secretary Dr Sanjiv Ranjan Kr. Singh

YOUR HEALTH OF IMA



Hony. Associate Editor Dr Sankar Sengupta



Hony. Associate Editor Dr Bibartan Saha



Dr Parul Vadgama



Hony. Secretary Dr Samarendra Kumar Basu

IMA HOSPITAL BOARD OF INDIA









Chairman

Dr A K Ravikumar

Hony. Secretary Dr Dinesh B. Thakare



Hony. Editor Dr Sudhir Singh

Hony. Associate Editor Dr Arun Kumar Tripathi

Hony. Associate Editor Dr Shailendra Kumar Singh

APKA SWASTHYA OF IMA

Hony. Secretary Dr Ritu Garg



Dr Rajeev B. Agarwal

Hony. Joint Secretary Dr Deepak Kr. Singh











Treasurer

JIMA COMMITTEE - 2024



Dr. R V Asokan National President, IMA



Dr Sanjoy Banerjee Hony. Editor, JIMA



Prof (Dr) Tamonas Chaudhuri Member, JIMA Committee



Dr. Anilkumar J Nayak Hony Secretary General, IMA

Dr Ranjan Bhattacharyya

Hony. Associate Editor,

.IIMA

Dr Samrendra Kumar Basu

Member, JIMA Committee



Dr Prasanta Kumar Bhattacharyya Hony. Associate Editor, JIMA



Dr Sanjay Banerjee Member, JIMA Committee





Dr. Sibabrata Banerjee Hony. Secretary, JIMA



Dr. Sarbari Datta

Hony. Jt. Finance Secretary, Hqs



Dr. Minakshi Gangopadhyay Hony. Assistant Secretary, JIMA



Dr Udas Chandra Ghosh Member, JIMA Committee

ANDMARK ACHIEVEMENT OF JIMA



The National Medical Commission (NMC) guidelines for medical faculty in India requires publications of meta-analysis, systematic reviews, original articles, or case series in journals indexed in Medline, PubMed Central, Citation index, Sciences Citation Index Expanded (SCIE), Embase, Scopus, or Directory of Open Access Journals (DOAJ).

Journal of the Indian Medical Association (JIMA) is already indexed in SCOPUS since 2020.

Once again, we are successful to index JIMA in yet another indexing in EMBASE in 2024 with retrospective effect (Serial No. 5532 in the Embase Jan 2024 Journal List).

Work is going on for PUBMED CENTRAL indexing in full swing.

We are really grateful to **Dr. R. V. Asokan**, our beloved National President and **Dr. Anilkumar J. Nayek**, our Hony. Secretary General for round the year support to JIMA Committee.

I express my heartfelt gratitude to all the JIMA Committee members, the Reviewers and Staffs of JIMA for this historical achievement of JIMA.

Dr. Sanjoy Banerjee Hony. Editor, JIMA

Listed in Embase at serial No.5532

	A	В	С	D
1	Embase journal titles (Jan 2024)	Abbreviated title	ISSN .	EISSN
5526	Journal of the History of Ideas	J. Hist. Ideas	00225037	10863222
5527	Journal of the History of Medicine and Allied Sciences	J. Hist. Med. Allied Sci.	00225045	14684373
5528	Journal of the History of the Behavioral Sciences	J. Hist. Behav. Sci.	00225061	15206696
5529	Journal of the History of the Neurosciences	J. Hist. Neurosci.	0964704X	17445213
5530	Journal of the Hong Kong College of Cardiology	J. Hong Kong Coll. Cardiol.	10277811	
5531	Journal of the Indian Chemical Society	J. Indian Chem. Soc.	00194522	
5532	Journal of the Indian Medical Association	J. Indian Med. Assoc.	00195847	
5533	Journal of the Intensive Care Society	J. Intensive Care Soc.	17511437	8
5534	Journal of the International AIDS Society	J. Int. AIDS Soc.		17582652
5535	Journal of the International Association of Providers of AIDS	J. Int. Assoc. Providers AIDS Care	23259574	23259582

JIMA Editorial Advisory Board Members (National and International)



Dr. Vedprakash Mish Physiology Maharashtra







Dr D P Singh Respiratory Medicine Bhagalpur, Bihar



Dr V G Pradeep Kumar Neurologist Kozhikode, Kerala



Dr C Daniala Radiologist Shillong, Meghalaya



Dr Gautamananda Roy Acute & Stroke Medicine UK



Dr Fazila TN Malik Cardiologist Dhaka Bangladesh



Dr Vinay Aggarwal Physician New Delhi



Dr. T. Nirmal Fredrick Ophthalmologist Tamilnadu



Dr Surya Kant Respiratory Medicine Lucknow



Dr V AmuthanEmeritus Cardiologist Tamil Nadu



Dr Anju Grewal Anaesthesiologist Punjab



Dr Colin Robertson A&E Medicine UK



Dr. Ricardo Escalante Colorectal Surgeon Venezuelan



General Surgeon Maharashtra



Dr Mansukh R Kanani Paediatrician Gujarat



Dr G Narsimulu Rheumatologist Hyderabad



Dr V Mohanan Nair Public Health Ananthapuri



Dr Vikram Kate Gastro Surgeon Puducherry



Dr Shohael M Arafat Medicine Bangladesh



Dr SM Mostafa Zaman Cardiologist Dhaka, Bangladesh



Radiologist Bengal



Dr Ravindra Kute Oncosurgeon Maharashtra



Dr S.M. Kadri Public Health Kashmir



Dr. C Palanivelu Robotic Gastro Surgeon Coimbatore



Dr Om Tantia Bariatric Surgeon Kolkata



Dr Narimantas E Samalavicius Robotic Surgeon Lithuania







Dr J.A. Jayalal Surgeon Tamilnadu



Dr. Subramanian Nallasivan Rheumatlogist Tamiinadu



Dr Apurba Ghosh Paediatric Medicine kolkata



Dr Alok Pandit Neurologist Kolkata



Dr Bibhuti Saha Tropical Medicine Kolkata



Prof Roman Jaeschke Medicine Canada



DrWJW Nunco - Mensah, Colorectal Surgeon London



Dr Bipin M Patel Anaesthesiologist Gujarat



Dr Shashank Joshi Endocrinologist Mumbai



Dr. Tanu Raj Sirohi Internal Medicine Uttar Pradesh



Dr Deepraj Bhandarkar Minimal Access Surgeon Mumbai







Dr Partha Sarathi Roy Neurologist UK



Dr Aminur Rahman Neurologist Dhaka, Bangladesh

www.dishaeye.org



Launching Disha Multi-speciality Hospital

The future of affordable healthcare is now in the heart of New Town!



Disha, the largest eyecare provider in Eastern India, has been dedicated to serving the patients of Eastern India for the past 25 years. We are embarking on a new venture with the upcoming Disha Multi-speciality Hospital this Q4 2024, focusing on healthcare that's accessible for all.

Built on 3-acre land | A huge 5 lakh sq. ft. structure | B+G+5 floors | 600-beds

We invite collaboration with experienced doctors who can bring their expertise to individual subspecialties and contribute to the hospital's development with enthusiasm and accountability.

Come, let's together redefine our healthcare standards in this greenfield initiative.





Volume 122 (JIMA) Number 12

December 2024 KOLKATA ISSN 0019-5847

JOURNAL Of the INDIAN MEDICAL ASSOCIATION

11 Editorial

23

29

35

40

44

47

2024 : A Year of Transformative Strides in Medicine, Surgery and Public Health — Sanjoy Banerjee, Shambo S Samajdar

19 Original Articles

Diagnostic Yield of Bronchoscopy in Benign Lung Diseases at Tertiary Health Care Hospital in South Gujarat — Jignesh Karamshibhai Gengadiya, Grinish Pravinbhai Tamakuwala, Parul Kishorbhai Vadgama, Jeel Vijaybhai Mangrola, Gopika Premchandra Khatri

A Study of the Clinico-demographic Profile and Clinical Outcomes of COVID-19 Patients in a Tertiary Care Hospital of India — *Rupankar Nath, Deepannita Sutradhar, Riturag Thakuria*

Levels and Risk Factors of Diabetes Mellitus among Tuberculosis Patients Attending District Tuberculosis Centre, Vijayapur — *Mallikarjun C Yadavannavar, Tanuja P Pattankar, Daneshwari P Kouttalmatt*

Extent of Disability and Behavioral Disturbances of the Children with Intellectual Disability and the Coping Mechanisms of their Parents : An Observational Study from a tertiary Care Hospital of Eastern India — *Paramita Patra, Amitava Dan, Sharmila Sarkar, Asish Mukhopadhyay, Debasish Sanyal*

Neutrophil to Lymphocyte Ratio (NLR) as an Inflammatory Marker in Predicting Severity in Acute Ischemic Stroke Patients — Deepti Sharma, Gordhan Lal Nagar, Narendra Fageria, Anup Kumar Mangal, Pawan Kumar

Microvascular Density of Invasive Breast Carcinoma as an Independent Prognostic Factor in Comparison with Already Established Ones : A Study from Northeast India — *Junu Devi, Asreen Suhana*

Comparative Analysis of Sensitivity and Specificity of Creatinine Kinase MB and Cardiac Troponin I for the Diagnosis of Acute Myocardial Infarction — *Nidhi Rohan Purandare, Astha Goyal, Sarita Shinde, Shilpa Jain*





Of the INDIAN MEDICAL ASSOCIATION

Volume 122 (JIMA) Number 12 December 2024 KOLKATA ISSN 0019-5847

Prevalence of Depression and Anxiety among Elderly Patients Visiting Tertiary Care Hospital — Samreen Kour, Manmeet Singh, Parul Gupta, Raghav Arora, Urvi Gupta

Evaluating Farmers' Knowledge, Attitude and Practice Regarding Pesticide Use and Its Impact on Human Health in Northern Karnataka — A Cross Sectional study — *Nagendra Angirekula, Shailaja S Patil*

61 **Review Article**

JOURNAL

52

57

69

Cardiac Steatosis — An Emerging Entity — Aradhya Sekhar Bagchi, Ananda Bagchi

66 Case Reports

Scrub Typhus with Unilateral Parotitis and Encephalopathy : A Rare Case Report and Literature Review — Umakanta Mahapatra, Soham Das Bakshi, Soumyadip Mukhopadhyay, Biplab Kumar Gayen

Neuropsychiatric Lupus with Apla Syndrome and Auto-immune Haemolytic Anaemia in a Patient of Hansen's Disease — A Diagnostic and Therapeutic Challenge — *Sudipto Chakraborty, Parthajit Das, Kishalaya Karan, Jayanta Sharma*

72 Letter to the Editor

Does only Mupirocin enough for Nasal Decolonization in Methicillinresistant *Staphylococcus Aureus* Carriers or do we need more ? — *Sahjid Mukhida, Amitesh Datta, Preethy Edavaloth, Vishal Pawar, Sundip Mukherjee*

73 Index to Volume 122

2024 : A Year of Transformative Strides in Medicine, Surgery and Public Health

As 2024 draws to a close, the Journal of the Indian Medical Association (JIMA) reflects with pride on the remarkable advancements and challenges that have defined the year in medicine and public health. These 12 months were a testament to the resilience, innovation, and dedication of the global medical community in addressing some of the most pressing healthcare issues of our time.

This year was not just about scientific discoveries but also about fostering a deeper understanding of the human condition and its interplay with health systems, policies, and technology. From groundbreaking therapies that redefined chronic disease management to policy interventions aimed at ensuring equitable access to healthcare, 2024 encapsulated the diversity and depth of the global healthcare landscape.

Through its editorials, JIMA captured the pulse of these developments, offering a platform for thought leadership and dialogue on the multifaceted challenges in healthcare. Themes ranging from the ethical integration of artificial intelligence and advancements in surgical technologies to the evolving landscape of mental health and the persistent fight against inequities in healthcare access have illuminated the path forward.

This retrospective highlights the pivotal moments and overarching themes of 2024, offering insights into how these advancements are shaping our collective pursuit of health and well-being. It also serves as a reminder of the enduring challenges that require our unwavering commitment as we move into 2025. By reflecting on the strides made and the lessons learned, JIMA reaffirms its dedication to fostering knowledge, innovation, and collaboration in the ongoing journey toward better health for all.

ADVANCING THERAPEUTICS AND TECHNOLOGIES

Breakthroughs in Diabetes Care :

The year 2024 marked pivotal advancements in the treatment of Type 2 Diabetes Mellitus (T2DM), with oral semaglutide emerging as a landmark therapeutic option for more than last 2 years. This novel oral GLP-1 receptor agonist, marketed under the brand name Rybelsus, revolutionized diabetes care by addressing the long-standing challenges associated with injectable GLP-1 agonists. Leveraging innovative SNAC (Sodium N-[8-(2-hydroxybenzoyl) Amino] Caprylate) technology, oral semaglutide ensures stable absorption of the peptide in the gastrointestinal tract, overcoming degradation barriers and offering efficacy comparable to its injectable counterparts. This innovation not only simplifies the management of

T2DM but also significantly enhances patient adherence and satisfaction by eliminating needlerelated barriers such as fear and discomfort.

Additionally, the development of once-weekly insulins like Insulin Icodec and Efsitora Alfa heralded a transformative step in basal insulin therapy. These ultra-long-acting insulins, with pharmacokinetic profiles offering stable glycemic control over a week, reduce the need for daily injections, mitigating the logistical and psychological burdens on patients. Phase III trials revealed their efficacy in achieving glycemic targets comparable to daily basal insulins, with no additional risks of hypoglycemia. By minimizing the frequency of injections, these insulins are poised to redefine diabetes management, making therapy more accessible and sustainable for a larger population.

Progress in Robotic-Assisted Surgery :

The field of surgical technology witnessed groundbreaking enhancements in Robot-assisted Surgery (RAS). The da Vinci Xi system, a masterslave robotic platform, continued to set the standard for precision, offering unparalleled three-dimensional magnified views and ergonomic control for surgeons. These advancements translated into reduced intraoperative complications, such as blood loss, and improved surgical outcomes across specialties, including urology, gynecology, and oncology.

To address one of the key limitations of robotic platforms – the lack of tactile sensation – innovations like VerroTouch, a haptic feedback system under development, have garnered significant attention. This system integrates vibrotactile feedback and stereo sound to replicate the sensation of touch, enabling surgeons to navigate complex anatomical structures with greater confidence. Complementing these advancements, augmented vision systems and realtime tracking technologies allowed precise overlay of imaging data, such as MRI or CT scans, onto the surgical field. This integration of imaging and robotics transformed procedures like radical prostatectomies, enhancing both oncological precision and functional outcomes .

Artificial Intelligence : Revolutionizing Medicine

Artificial Intelligence (AI) maintained its trajectory as a transformative force across healthcare in 2024. AI technologies expanded their footprint, from predictive analytics in cardiology to diagnostic imaging in oncology. For instance, AI algorithms trained on large datasets demonstrated superior accuracy in predicting cardiovascular risks compared to traditional methods. In oncology, Al-powered tools optimized histopathological image analysis, facilitating earlier cancer diagnoses and personalized treatment strategies.

Despite these breakthroughs, challenges persist. Ensuring the clinical validation of AI tools remains critical to gaining widespread trust and adoption. Many models show promise in controlled environments but lack robustness in real-world scenarios, raising concerns about generalizability. Furthermore, ethical issues, such as data privacy, algorithmic biases, and the potential over-reliance on AI systems, highlighted the need for a balanced approach that integrates technological innovation with human oversight.

The path forward lies in embracing AI as a collaborative partner, augmenting rather than replacing human expertise. This partnership must prioritize ethical frameworks, transparency, and rigorous validation to harness AI's full potential while preserving the human touch central to patient care.

CHRONIC DISEASES AND PREVENTIVE CARE

Tackling End-Stage Kidney Disease : The Promise of Bioartificial Kidneys

Chronic diseases like End Stage Kidney Disease (ESKD) remain a growing global health challenge, with increasing prevalence fueled by lifestyle diseases such as diabetes and hypertension. In India, the burden of ESKD is particularly daunting, with over 90% of patients unable to access renal replacement therapies due to financial constraints. Against this backdrop, 2024 saw significant editorial focus on bioartificial kidneys, a revolutionary development in nephrology.

Bioartificial kidneys represent a quantum leap in renal replacement therapy. These devices are designed to mimic the filtration, absorptive, and secretory functions of human kidneys, offering an alternative to conventional dialysis or transplantation. Leveraging breakthroughs in nanotechnology, they integrate silicon nanopore membranes and cultured proximal tubular epithelial cells to deliver continuous renal support without the need for immunosuppressive therapy. Their portability, efficiency, and independence from donor organs position them as a transformative solution for ESKD patients, especially in resource-limited settings.

Moreover, wearable and implantable versions of these devices have advanced to preclinical trials, demonstrating promising outcomes. These technologies could alleviate the dependency on dialysis centers, reduce treatment costs, and improve patients' quality of life, representing a beacon of hope for millions struggling with kidney failure globally.

Headache Disorders : Evolving Strategies for a Changing Landscape

Primary headache disorders, including migraine and tension-type headaches, constitute a significant but often underdiagnosed public health concern in India. These disorders affect a broad demographic, with recent studies highlighting their disproportionate impact on women and younger populations. The prevalence of migraine, for example, reaches as high as 23% in certain regions, emphasizing the need for targeted interventions tailored to local epidemiological trends.

Editorials in 2024 highlighted the shifting clinical landscape of headache disorders, focusing on improvements in diagnosis, management, and public health strategies. Advances in neuroimaging and diagnostic tools have facilitated better differentiation of primary headache types, enabling more precise treatment plans. The integration of pharmacological therapies, including triptans for acute attacks and beta-blockers or CGRP (Calcitonin Gene-Related Peptide) inhibitors for prophylaxis, has transformed patient outcomes. Additionally, non-pharmacological approaches, such as cognitive behavioral therapy and mindfulness-based interventions, have gained traction as complementary options.

Public health initiatives, including awareness campaigns and training programs for primary care physicians, have also been pivotal in bridging gaps in care. These efforts are particularly crucial in rural and underserved areas, where access to neurologists and specialized headache clinics remains limited. The emphasis on preventive care, lifestyle modification, and early intervention has positioned India to better address the substantial personal and societal burden imposed by headache disorders.

WOMEN'S HEALTH AND ENDOMETRIOSIS

Understanding the Complexity of Endometriosis :

Endometriosis, a chronic and often debilitating gynecological condition, was a focal point in 2024 editorials due to its multifaceted nature and significant impact on women's health. Affecting an estimated 10-15% of women of reproductive age globally, this enigmatic disease is characterized by the presence of endometrial-like tissue outside the uterus, leading to symptoms such as chronic pelvic pain, dysmenorrhea, dyspareunia, and subfertility. The varied clinical manifestations and the elusive pathophysiology of endometriosis continue to pose challenges for timely diagnosis and effective management.

Diagnostic Challenges :

A recurring theme in the discussions was the complexity of diagnosing endometriosis. While advances in imaging techniques like pelvic ultrasonography and MRI have enhanced diagnostic accuracy, limitations persist. Notably, the reliance on clinical presentation and the use of biomarkers such as CA-125, which have limited diagnostic value, underline the need for improved non-invasive diagnostic tools. Additionally, the editorial highlighted a paradigm shift from laparoscopic confirmation as the gold standard to more patient-friendly diagnostic strategies that prioritize symptom assessment and imaging.

Tailored Medical Interventions :

Hormonal therapies remain the cornerstone of endometriosis management. Editors emphasized the efficacy of Dienogest, a selective progestogen known for its ability to reduce pain and suppress disease progression. Long-term therapies such as Levonorgestrel-releasing intrauterine systems (LNG-IUS) and GnRH agonists were also noted for their role in mitigating recurrent symptoms and enhancing patient satisfaction. Continuous Combined Oral Contraceptives (COCs) were highlighted as an effective, non-invasive option for managing dysmenorrhea and other cyclical symptoms, providing sustained relief and improving quality of life.

Despite these advances, hormonal therapies are not without limitations, including side effects like irregular bleeding, weight gain, and potential impacts on bone mineral density. The editorials underscored the importance of tailoring treatments to individual patient profiles, considering factors such as symptom severity, fertility goals, and tolerance to medications.

Innovations in Surgical and Assisted Reproductive Technologies :

Surgical intervention remains indispensable for certain patients, particularly those with severe pain, Deep Infiltrating Endometriosis (DIE), or ovarian endometriomas. The editorials advocated for evidence-based surgical approaches, emphasizing the benefits of excision over ablation for reducing pain and recurrence. However, the potential impact of surgery on ovarian reserve necessitates a balanced discussion with patients, particularly those seeking fertility preservation.

In cases of endometriosis-associated infertility, the role of Assisted Reproductive Technologies (ART)

was underscored as a pivotal option. Techniques such as In Vitro Fertilization (IVF) provide a pathway for achieving pregnancy in patients where other interventions have failed. Recent innovations in ART, including improved protocols for ovarian stimulation and embryo transfer, were highlighted as critical tools for addressing the unique challenges posed by endometriosis.

Bridging Gaps in Care :

The editorials drew attention to persistent gaps in care, including delays in diagnosis and limited access to specialized treatment in resource-constrained settings. The psychosocial impact of endometriosis, encompassing its effects on mental health, relationships, and professional life, was a recurring theme, advocating for holistic management approaches. Strengthening public awareness, training healthcare providers, and investing in research to unravel the pathophysiology of the disease were identified as critical priorities for reducing the burden of endometriosis.

The Road Ahead :

While significant progress has been made in the diagnosis and management of endometriosis, the editorials stressed the need for continued innovation and a patient-centered approach. By integrating advancements in medical and surgical therapies with broader systemic changes in healthcare delivery, there is potential to transform outcomes for millions of women affected by this condition.

GLOBAL HEALTH AND POLICY: HEALTH AS A HUMAN RIGHT

Advocating for Universal Health Coverage :

The editorial "My Health, My Right", published in alignment with World Health Day 2024, served as a powerful call to action, emphasizing the inextricable link between health and human rights. Framed around the theme of equitable access to healthcare, the editorial highlighted the urgency of achieving Universal Health Coverage (UHC) – a vision where all individuals, irrespective of socioeconomic status, geography, or gender, can access quality health services without financial hardship. With over 4.5 billion people globally lacking full access to basic healthcare services, the editorial underscored the critical need for systemic reforms to bridge this gap.

Progress and Challenges in South-East Asia :

The editorial presented a mixed picture of progress within the South-East Asia Region (SEAR). Notable achievements included an increase in the Universal Health Coverage service coverage index from 47 in 2010 to 62 in 2021, alongside a 68.5% reduction in maternal mortality and dramatic declines in neonatal and under-five mortality rates. Despite these advances, disparities remain stark: nearly 40% of the region's population still lacks access to basic healthcare, with preventable deaths from non-communicable diseases (NCDs) such as cancer, diabetes, and cardiovascular conditions persisting at alarmingly high rates.

Equity in Access to Care :

The editorial addressed the pervasive inequities in healthcare access, particularly for marginalized populations. Women, rural communities, and economically disadvantaged groups were identified as bearing the brunt of healthcare disparities. Genderbased inequities in diagnosing and treating conditions like diabetes and hypertension were highlighted, with women often receiving delayed or insufficient care compared to men. The editorial also pointed to the persistent public health crisis of violence against women, noting that over one-third of women in SEAR have experienced intimate partner violence, further undermining their health and well-being.

Policy Directions and Systemic Reforms :

To realize the vision of health as a fundamental right, the editorial advocated for multifaceted systemic reforms:

- Investing in Healthcare Infrastructure: Prioritizing underserved regions and building robust systems to ensure the availability of quality health services in both rural and urban areas.
- Human Resource Development: Expanding the healthcare workforce, with a focus on training primary care providers, midwives, and nurses to address shortages, particularly in rural areas.
- Health Financing: Implementing financial models that protect households from catastrophic health expenditures, such as public insurance schemes and government subsidies for essential services.
- Gender Equity in Healthcare: Developing targeted interventions to close gender gaps in diagnosis, treatment, and prevention, with an emphasis on diseases that disproportionately affect women.
- Community Engagement: Empowering communities through health education, awareness campaigns, and the promotion of preventive care practices.
- 6) Integrated Health Systems: Strengthening primary healthcare systems to serve as the foundation for comprehensive, accessible, and equitable health services.

Health Beyond Healthcare :

The editorial highlighted the importance of addressing broader determinants of health, such as clean water, adequate nutrition, safe housing, and education. It argued that achieving health equity requires tackling social and environmental inequities that perpetuate cycles of poverty and ill-health. By framing health as a cornerstone of human dignity and development, the editorial underscored its role in fostering inclusive and prosperous societies.

A Moral and Practical Imperative :

At its core, "My Health, My Right" reminded readers that equitable healthcare is not merely a moral obligation but also a practical necessity for sustainable development. As nations navigate complex public health crises and demographic shifts, ensuring access to healthcare for all emerges as a crucial strategy for building resilient communities and economies.

A Call to Action :

The editorial concluded with a resounding call for collective action, emphasizing the urgent need for governments, policymakers, and global stakeholders to come together in the pursuit of health equity. It highlighted the importance of aligning national and international efforts with the principles of Sustainable Development Goal 3 - to ensure healthy lives and promote well-being for all at all ages. By addressing systemic inequities, investing in inclusive healthcare systems, and fostering global cooperation, the vision of health as a universal right can evolve from a distant aspiration to a tangible reality. The editorial underscored that achieving this goal is not only a moral imperative but also a critical step towards creating resilient communities and sustainable futures, where no one is left behind in the journey toward health and well-being.

MENTAL HEALTH IN TRANSITION

Progress Shaped by the Mental Healthcare Act of 2017 :

The Mental Healthcare Act of 2017 represented a significant legislative milestone in India, laying the foundation for a rights-based approach to mental health. By defining access to mental healthcare as a fundamental right, the Act mandated the provision of affordable, accessible, and high-quality mental health services for all citizens. It also emphasized decriminalizing suicide, safeguarding patient rights, and ensuring that mental health care is provided with dignity and respect. The editorials in 2024 reflected

on the progress spurred by this Act, particularly in raising awareness and encouraging policy shifts that prioritize mental health as a critical component of public health .

Integration of Mental Health into Primary Care :

The District Mental Health Program (DMHP) emerged as a cornerstone initiative for integrating mental health services into primary healthcare. This program, operational under the broader National Mental Health Program (NMHP), was designed to decentralize mental health services, making them accessible to underserved populations at the district level. The DMHP focuses on early detection, treatment, rehabilitation, and community outreach, addressing conditions ranging from Common Mental Disorders (CMDs) like depression and anxiety to Severe Mental Disorders (SMDs) such as schizophrenia and bipolar disorder.

Key Components of the DMHP include :

- Training Primary Care Workers: Empowering general practitioners and healthcare workers with the skills to identify and manage mental health conditions.
- Community Engagement: Conducting mental health awareness campaigns to reduce stigma and promote early intervention.
- Strengthening Infrastructure: Establishing district mental health clinics and integrating psychiatric wings in general hospitals.
- Service Delivery Models: Promoting telepsychiatry and mobile outreach to bridge the gaps in rural and remote areas.

Persisting Challenges: Stigma and Resource Gaps :

Despite these advancements, significant barriers persist. Stigma surrounding mental illness remains one of the most formidable obstacles to care. Many individuals delay or avoid seeking treatment due to fear of social ostracism, compounded by limited awareness of mental health conditions. The editorials highlighted the need for sustained public education campaigns to challenge misconceptions and normalize discussions about mental health.

Additionally, resource constraints hinder the effective implementation of mental health programs. A glaring shortage of mental health professionals – including psychiatrists, psychologists, and social workers – limits the program's reach. India's mental health workforce density remains far below global recommendations, particularly in rural areas where access to specialized care is minimal. Financial

constraints further exacerbate the situation, with mental health services often receiving insufficient funding relative to other health sectors.

The Role of Technology in Mental Health :

Amid these challenges, technology is emerging as a transformative tool in mental health care. Telepsychiatry platforms, digital mental health apps, and virtual counseling sessions are increasingly being integrated into service delivery models. These innovations have expanded access to care, particularly for populations in remote and underserved areas and have shown promise in mitigating the stigma associated with in-person consultations.

Charting a Path Forward :

To build on the momentum created by the Mental Healthcare Act and the DMHP, a multi-pronged approach is essential:

- 1. Scaling Mental Health Services: Expanding the DMHP to cover all districts and ensuring consistent funding for program sustainability.
- Workforce Development: Investing in the training and recruitment of mental health professionals and incentivizing their deployment in rural areas.
- Community-Based Care Models: Strengthening partnerships with local organizations and self-help groups to create a supportive ecosystem for mental health care.
- 4. Policy Advocacy: Prioritizing mental health in national health budgets and fostering cross-sectoral collaborations to address social determinants of mental health.

Towards a Mentally Healthy India :

The editorials concluded that while the progress in mental health care has been encouraging, the road ahead demands unwavering commitment from stakeholders. Bridging the gaps in service delivery, addressing stigma, and integrating mental health into broader healthcare systems are pivotal to ensuring equitable access to care. By leveraging the potential of technology, fostering community engagement, and advocating for systemic reforms, India can chart a path toward a future where mental health is not only a policy priority but also a societal norm.

LEGAL AND ETHICAL DIMENSIONS IN MEDICINE

Navigating Medical Negligence : Professional Accountability

The issue of medical negligence was a prominent theme in the 2024 editorials, highlighting its critical impact on trust within doctor-patient relationships. Medical negligence, as defined by the Medical Council Act, involves a failure on the part of a registered practitioner to provide the standard of care expected, leading to harm or injury to the patient. The editorials emphasized that professional accountability is central to mitigating negligence. This includes maintaining up-to-date knowledge, practicing within the scope of expertise, and adhering to established protocols and guidelines. The importance of clear communication with patients and meticulous documentation of clinical decisions was also stressed as foundational to demonstrating accountability.

Patient Consent : A Cornerstone of Ethical Practice

Editorials underscored the pivotal role of informed consent in ethical medical practice. Consent must be obtained through a transparent process where patients are provided with clear, comprehensive information about their diagnosis, treatment options, risks, benefits, and potential alternatives. The editorials delineated between types of consent:

- Implied Consent, limited to routine examinations and non-invasive procedures.
- Written Consent, mandatory for invasive interventions, high-risk procedures and treatments involving potential complications.

The practice of obtaining "blanket consents," which lack specificity, was criticized as legally invalid. Instead, tailored, procedure-specific consents were advocated, witnessed by neutral parties to ensure ethical compliance. The inclusion of therapeutic privilege – where certain risks might be withheld if disclosure is deemed detrimental to the patient – was discussed as a nuanced aspect requiring cautious application.

Ethical Standards : A Guide to Medical Integrity

The editorials stressed adherence to ethical principles as the backbone of medical integrity. Key recommendations included:

- Respect for Patient Autonomy : Honoring patients' rights to make informed decisions about their care.
- Non-Maleficence and Beneficence : Balancing the duty to avoid harm with the imperative to promote well-being.
- **3)** Justice : Ensuring equitable access to care, regardless of socioeconomic or demographic factors.

Ethical concerns surrounding confidentiality, especially in sensitive cases, were highlighted. Practitioners were advised to uphold privacy while navigating scenarios where third-party disclosures might be necessary for public safety.

Preventive Strategies to Mitigate Litigation :

To reduce the incidence of medical litigation, the editorials outlined practical measures:

- 1) Continuing Medical Education (CME) : Regularly updating knowledge and skills to meet evolving standards of care.
- Professional Etiquette : Maintaining respectful communication with patients and colleagues to foster trust.
- Detailed Medical Records : Keeping thorough, chronological, and tamper-proof documentation as evidence of due diligence.
- Seeking Second Opinions : In complex or ambiguous cases, consulting peers to ensure comprehensive evaluation and shared responsibility.

Special attention was given to the importance of a professional demeanor, advising practitioners to communicate empathy, avoid overpromising outcomes, and ensure transparency during adverse events.

Fostering Trust in the Doctor-Patient Relationship

The editorials acknowledged that trust is the cornerstone of effective healthcare delivery. Instances of negligence, whether perceived or real, erode this trust and often lead to disputes. To rebuild and sustain this trust, practitioners were encouraged to adopt proactive communication strategies, foster openness in discussing complications and prioritize patient safety above all.

The Ethical and Legal Horizon :

Looking ahead, the editorials advocated for systemic reforms to support practitioners and patients alike. These include:

- Legal Safeguards for Practitioners : Implementing frameworks that protect doctors from frivolous lawsuits while holding them accountable for genuine lapses.
- Patient Advocacy Systems : Establishing mechanisms to mediate disputes and provide patients with a fair hearing before litigation.
- Ethics Committees : Strengthening institutional review boards to guide practitioners through ethically complex decisions.

The discussions on medical negligence and ethics in 2024 emphasized that a culture of transparency, accountability and respect is essential for advancing healthcare. By adhering to high ethical standards and implementing robust preventive strategies, the medical community can not only reduce litigation risks but also foster a stronger, trust-based doctor-patient relationship that ultimately enhances the quality of care.

The Road Ahead :

As we stand at the threshold of a new year, the editorial reflections from 2024 offer a moment of pause – both to celebrate the strides we have made and to acknowledge the work that remains. The year has been marked by transformative progress across healthcare, from the introduction of groundbreaking therapies to advancements in technology and policy frameworks aimed at fostering equitable health systems. These developments serve as a testament to the resilience and ingenuity of the global medical community, yet they also underscore the complexity of the challenges that lie ahead.

Expanding Access to Groundbreaking Therapies

The advent of innovative therapies, such as oral semaglutide for diabetes and bioartificial kidneys for chronic kidney disease, has shown the potential of modern medicine to not only enhance patient outcomes but also improve accessibility and adherence. The road ahead demands a concerted effort to ensure these advancements are not limited to affluent populations but reach underserved communities globally. Scaling distribution networks, reducing costs, and fostering partnerships between governments and private sectors will be critical to achieving universal access to these life-saving interventions.

Building Equitable Health Systems :

The call for universal health coverage and the recognition of health as a fundamental right have gained momentum, but significant inequities persist. The path forward must prioritize closing gaps in healthcare access, particularly for marginalized populations, women, and rural communities. Strengthening primary healthcare, addressing social determinants of health and investing in infrastructure and human resources will be foundational in building resilient and inclusive health systems. This endeavor will also require innovative financing mechanisms and policies that ensure no one is left behind in the pursuit of well-being.

Navigating the Ethical Integration of AI :

Artificial Intelligence continues to redefine the boundaries of medicine, offering unprecedented opportunities to improve diagnostics, personalize treatments and optimize healthcare delivery. However, its integration is fraught with ethical challenges, including issues of privacy, bias and the potential for over-reliance on machine learning systems. The road ahead must include robust regulatory frameworks, transparent validation processes, and an unwavering commitment to balancing Al's capabilities with the irreplaceable human touch. Embracing AI as a collaborative tool rather than a replacement for clinical expertise will be key to its responsible and effective implementation.

Advancing Research and Innovation :

Innovation remains the lifeblood of medical progress, and the need for multidisciplinary research has never been greater. From precision medicine to regenerative therapies and sustainable public health strategies, the scope of research must expand to address the evolving challenges of our times. Collaboration between academia, industry, and policymakers will play a crucial role in translating discoveries into tangible healthcare improvements.

Fostering a Global Dialogue :

The challenges of global health – from pandemics to chronic disease burdens – demand collective action and shared learning. JIMA reaffirms its commitment to being at the forefront of this evolving narrative, serving as a platform for disseminating knowledge, fostering dialogue among stakeholders, and inspiring innovation. By championing evidence-based practices, ethical integrity and inclusivity, the journal aims to contribute meaningfully to shaping the future of healthcare.

A Call to Action :

The journey ahead is as demanding as it is promising. As we navigate the complexities of a rapidly evolving healthcare landscape, the vision of a healthier, more equitable world must guide our efforts. This requires not only technical excellence but also a steadfast dedication to compassion, collaboration, and the well-being of all. The reflections from 2024 remind us that progress is possible when we unite in purpose and action, and they inspire us to meet the opportunities and challenges of 2025 with renewed resolve and optimism.

JIMA looks forward to continuing its legacy of thought leadership and advocacy, shaping a future where medical advancements, ethical practices, and equitable health systems converge to uplift humanity. Together, we can transform the aspirations of today into the achievements of tomorrow.

Hony Editor, JIMA	Sanjoy Banerjee ¹
MD, DM (Clinical Pharmacology)	Shambo S Samajdar ²
PG Dip Endo & Diabetes (RCP),	_
Fellowship in Respiratory and Critic	al Care (WBUHS),
Fellow Diabetes India, Fellow India	n Pharmacological Society;
ndependent Clinical Pharmacologis	st
Consultant Physician	

Original Article

Diagnostic Yield of Bronchoscopy in Benign Lung Diseases at Tertiary Health Care Hospital in South Gujarat

Jignesh Karamshibhai Gengadiya¹, Grinish Pravinbhai Tamakuwala², Parul Kishorbhai Vadgama³, Jeel Vijaybhai Mangrola⁴, Gopika Premchandra Khatri⁵

Methodology : Sixty patients with benign lung disorders like Suspected Sputum Negative Pulmonary Tuberculosis, Suspected Fungal Pneumonia, Suspected Bacterial Pneumonia and Suspected Interstitial Lung Disease were enrolled from April, 2020 to April, 2021 in this study. Detailed clinical history, physical examinations and necessary investigations were carried out. Eligible patients were subjected for fiberoptic video bronchoscopy. During Bronchoscopy Gross fiberoptic bronchoscopy findings noted and Bronchoalveolar lavage or tissue were sent for necessary investigations. Data entry and analysis were done using MS Excel sheet 2013.

Result : Overall diagnostic yield of bronchoscopy in 60 patients was 63.3%. Out of 37 Sputum negative suspected cases of Pulmonary Tuberculosis posted for bronchoscopy, 22(59.4%) cases were confirmed to have tuberculosis, only 1(2.7%) case was having Bacterial pneumonia and 2(3.4%) cases were diagnosed with Tuberculosis had super added bacterial infection. Other 13 patients suspected of having bacterial pneumonia, 7(53.3%) patients were confirmed to have bacterial pneumonia while 2(15.3%) cases were found positive for tuberculosis. Out of 5 suspected cases of fungal pneumonia and 5 suspected cases of Interstitial Lung Disease, each 2(40%) patients had positive findings respectively. Overall complications developed in 7(11.7%) patients.

Conclusion : Our study suggests, fiberoptic bronchoscopy can provide good sample material for diagnosis of suspected cases. Bronchoalveolar Lavage and Bronchial Brushing sent for AFB, CBNAAT, TB Culture, Cytology, KOH etc were cumulatively used to increase diagnostic yield of bronchoscopy.

[J Indian Med Assoc 2024; 122(12): 19-22]

Key words : Bronchoalveolar Lavage, Bronchial Brushing, CBNAAT.

iberoptic Bronchoscopy (FOB) is a very useful and safe procedure for diagnosis of various respiratory diseases. FOB can be performed under local anaesthesia in various clinical/hospital settings providing maximal visualization of tracheobronchial tree¹, if performed carefully, can be a thoroughly safe procedure². Many patients having clinical and radiological features of pulmonary tuberculosis have negative sputum smear examinations, even if repeated on several occasions and their sputum culture for Acid Fast Bacilli (AFB) may also turn to be negative. Incorrect diagnosis, inadequate antibiotic therapy, impaired host defence, atypical organisms, resistant

Received on : 25/07/2023 Accepted on : 09/09/2023

Editor's Comment :

Fiberoptic bronchoscopy is useful tool in diagnosing of suspected smear negative pulmonary tuberculosis, suspected bacterial, fungal and ILD with respect to their association with clinical and radiological profile at early stage despite not meeting the routine bacteriological criteria for diagnosis and helps in initiating early treatment which in turn improve patient outcome.

pathogens, non-infectious causes, tuberculosis, Endobronchial lesions etc are common causes of nonresolving pneumonia or slowly resolving pneumonia. Fiberoptic bronchoscopy frequently helps to identify the exact cause of difficult to treat pneumonia cases. FOB is immensely useful for making a conclusive diagnosis of lung disease, especially when there is an Endobronchial lesion, providing adequate tissue sample by Endobronchial biopsy, Bronchoalveolar Lavage (BAL) or brush cytology³.

AIMS AND OBJECTIVES

To assess the role of Bronchoscopy in diagnosis of Benign lung disorders like Smear Negative Pulmonary Tuberculosis, Non Resolving Pneumonia, Fungal Pneumonia, Bronchiectasis, Interstitial Lung Disease (ILD) etc.

¹MBBS, MD, Senior Resident, Department of Pulmonary Medicine, New Civil Hospital, Surat, Gujarat 395001

²MD (Respiratory Medicine), Assistant Professor, Department of Respiratory Medicine, Government Medical College, Surat, Gujarat 395001 and Corresponding Author

³MD (TB and Chest), Professor and Head, Department of Respiratory Medicine, Government Medical College, Surat, Gujarat 395001

⁴MBBS, Intern, Department of Respiratory Medicine, SMIMER Hospital & Medical College, Surat, Gujarat 395010

⁵MBBS, Junior Resident, Department of Respiratory Medicine, SMIMER Hospital & Medical College, Surat, Gujarat 395010

MATERIAL AND METHOD

This is a Prospective study of 60 patients was conducted over a period of one year, from April, 2020 to April, 2021 at Tertiary Health Care Hospital In South Gujarat.

Patient with Age >18 years, Suspected Smear negative pulmonary tuberculosis, Non Resolving pneumonia, Suspected fungal pneumonia, Suspected bacterial pneumonia, Bronchiectasis and Suspected Interstitial Lung Disease (ILD) patients were enrolled in our study.

Patient with Age <18years,Smear positive pulmonary Tuberculosis, critical cardiac or respiratory illness, malignancy, Active bleeding disorder, HIV/ AIDS Positive were excluded from our study.

Informed written consent was obtained from all the patients enrolled in study. Detailed clinical history, physical examinations and investigations were carried out. Eligible patients with Chest X-ray and CECT Thorax and clinical findings consisting with Sputum Negative Suspected Pulmonary Tuberculosis, Suspected Bacterial Pneumonia, Suspected Fungal Pneumonia and Non-Resolving Pneumonia were subjected for fiberoptic video bronchoscopy. During bronchoscopy detailed examination of Bronchial tree was done and specimens including BAL, Bronchial brushing, TBLB & TBNA (done in indicated patients), Post Bronchoscopy Sputum (PBS) was collected and sent for necessary investigations. Data entry, prelimary Data analysis and preparation of charts were done in MS Excel sheet 2013.

OBSERVATIONS

A total of 60 patients were included in our study. Out of which 39(65%) patients were Male and 21(35%) patients were Female. The most common age group involved in the study was 31-40 years (28.3%). The youngest patient was aged 19 years and the oldest was 85 years and mean age was 44.5±16.7 years.

In the present study ,cough and fever were the most common symptoms (76.67%) followed by loss of appetite in 36.67% patients and breathlessness in 33.3% patients while loss of weight and night sweat were seen in 26.67% and 16.67% patients respectively. About 12% patients were having chest pain.

Chest X-ray finding of pulmonary infiltrative lesions was the most common finding in majority (55%) of patients followed by consolidation in 33.33% patients. Cavitary lesions and nodular lesion were seen in 13.33% patients each. Bronchiectatic changes and reticulations were noticed in 8.33% cases. Pleural effusion was seen in 6.67% cases whereas lymphadenopathy and collapse of lung was found in 5% of patients. Most common CT thorax finding was consolidation in two-third of the patients, followed by lymphadenopathy in half of the patients and tree in bud appearance 43.33%. Whereas fibrocavitary lesion, ground glass opacity and nodules were seen in about one third of patients. Septal thickening, reticulations and miliary mottling were seen in 23.33%, 13.33% and 11.67% patients respectively. Only 10% patients showed Pleural effusion and honey-comb appearance.

Out of 60 patients based on clinical and radiological findings, 37 patients were suspected to have Pulmonary Tuberculosis, 13 Patients were suspected to have bacterial pneumonia,5 Patients were suspected to have fungal pneumonia and 5 Patients were suspected to have ILD. All patients were then subjected to fiberoptic bronchoscopy

Out of 60 patients, 29 patients had no gross FOB findings. 12 (20%) patients had hyperaemia; 9 (15%) patients had congestion and 8 (13.33%) patients had fibrotic and stenotic changes. 5 (8.33%) patients had growth; 5 (8.33%) patients had ulcerative lesions and remaining 4 patients had shown active bleeding (Fig 1).

Table 2 suggest whenever Bronchoalveolar Lavage and Bronchial Brushing sent for AFB staining, CBNAAT, TB Culture, Gram stain and culture sensitivity, Cytology, KOH staining and NTM culture were cumulatively used for increasing diagnostic sensitivity.

Table 3 show, out of 37 suspected sputum negative pulmonary tuberculosis cases posted for bronchoscopy, 22(59.4%) cases were found to be positive for tuberculosis, Only 1(2.7%) case had findings consistent with Bacterial pneumonia and 2(3.4%) cases of tuberculosis had super added bacterial infection. In 13 patients suspected of bacterial pneumonia, more than half found positive for bacterial pneumonia while 2(15.3%) cases were found positive for pulmonary tuberculosis. Out of 5 suspected cases of fungal pneumonia and 5 suspected cases of Interstitial Lung Disease, each 2(40%) patients had positive findings respectively.. Thus, overall diagnostic yield of bronchoscopy in 60 patients was 63.3% (Fig 2).



Fig 1 — Distribution of patients according to Gross FOB findings (n=60)

Table 1 — Comprehensive table showing results of various tests ($n=60$)						
Sample type	Performed in	Positive in	Percentage			
	no or patients	no or patients				
BAL AFB*	60	17	28.3%			
BAL CBNAAT#	60	24	40.0%			
BAL KOH ^{\$}	60	2	3.33%			
BAL fungal culture	5	2	40.0%			
BAL Galactomanan	5	2	40.0%			
BAL Cytology	20	2	10.0%			
BAL TB culture	60	23	38.3%			
BAL NTM Culture ⁺	5	2	40.0%			
BB-AFB	60	22	36.7%			
BB-CBNAAT	60	23	38.3%			
BB-TB culture	43	21	48.8%			
TBNA:HP®	6	2	33.3%			
Endobronchial Biops	y 6	4	66.7%			
PBS AFB ^è	60	14	23.3%			

(*AFB-Acid Fast bacilli; * CBNAAT-Cartridge Based Nucleic Acid Amplification Test; *NTM- Non-tuberculosis Mycobacteria; ^{\$}KOH-Potassium Hydroxide; [@]TBNA:HP-Trans Bronchial Needle Aspiration: Histopathology Examination; ^{\$}PBS-Post Bronchoscopy Sputum)

Table 2 — Cumulative result of Bronchoscopy specimen					
Bronchoscopy specimen Cumulative res					
BAL TB CULTURE + BB TB CULTURE	24(40%)				
BAL AFB + BB AFB	22(36.7%)				
BAL CBNAAT + BB CBNAAT	24(40%)				
BAL (AFB+CBNAAT + TB CULTURE)	24(40%)				
BB (AFB+CBNAAT + TB CULTURE)	23(38.3%)				
BAL (AFB + CBNAAT + TB CULTURE					
+ GMCS + KOH + NTM + CYTOLOGY)	38(63.3%)				

Present study found 14/60 (23.3%) positive results for Post Bronchoscopy Sputum (PBS)-AFB staining.

Out of 60 patients, majority 53 (88.3%) patients did not develop any complications. 5 (8.3%) patients developed minor bleeding and 2 (3.3%) patients developed bronchospasm.

DISCUSSION

The delay in diagnosis and treatment in patients of Non resolving Pneumonia may lead to rise in mortality by 3-5%. For this reason Fiberoptic Bronchoscopy is one of the most widely used diagnostic approach among sputum negative and difficult to treat pneumonia patients³.

With the advent of Fiberoptic bronchoscopy, diagnosis of PTB in sputum smear negative patients has become more promising. The main advantage with this instrument is the ability to visualize the bronchial tree and collect samples directly from the bronchial pathology site⁴.

Fungal pneumonia is classically found in neutopenic, imuno-compromised, malignancy and post-transplant patients there is high mortality if treatment is delayed to confirm a definite diagnosis. Different diagnostic modalities used to detect invasive fungal pneumonia and the decision to treat⁵.

In our study out of 60 patients, 29 patients were normal with no gross FOB findings. 12 (20%) patients had hyperaemia; 9 (15%) patients had congestion and 8 (13.33%) patients had fibrotic and stenosis changes. About 8.33% patients had growth and erosions/ulcer while remaining 4 patients had shown bleeding. In study of Atul Luhadia, *et al*⁶, congestion and hyperaemia (36%) and mucopurulent/mucoid secretions was seen in (32%) cases.

Present study found 17 (28.3%) patients with positive BALAFB staining.

In BAL CBNAAT findings, 40% patients found positive out of which 33.3% Rifampicin sensitive and 6.7% Rifampicin Resistance. 36.7% patients came positive for BB-AFB.BAL liquid culture was seen positive in 23 (38.3%) patients.TBLB done in 6 patients out of which 4(66.6%) patients biopsy report show Caseating granuloma. Quaiser et al⁷, showed a positive yield by BAL fluid smear examination and culture as 17.5% (7/40) and 30% (12/40) respectively. Study by Yong Suk Jo, *et al*⁶, showed BAL positive for AFB smear in 23% and 50% positive for culture. Willcox, *et al*⁶, obtained BB from 83 of the 89 patients and 35 (42%) of them were positive on smear, making early diagnosis possible in 42%.

PBS-AFB staining came positive in 14/60(23.3%) patients. Quaiser, *et al*⁷, reported the yield of PBS smear as 17.5% (7/40) and PBS culture as 27.5% (11/40).

Complications were seen in only 7(11.7%) patients in our study. In study of Vipparthi Surya kumara, *et* al^{10} , reported complication in 23.3% cases.

Thus, overall diagnostic yield of bronchoscopy in our study was 63.3% in 60 patients subjected for

Table 3 — Diagnostic yield of Bronchoscopy in suspected different benign Lung Disease					
Suspected case of	Total suspected case (n)	Diagnosis done	Diagnosis not confirmed by Bronchoscopy		
Tuberculosis	37	22(59.4%) tuberculosis (16-Rifampicin sensitive, 4-Rifampicin resistance, 2-NTM) 1(2.7%) Bacterial pneumonia 2(3.4%) Bacterial Pneumonia with Tuberculosis	12(32.4%)		
Bacterial pneumonia	13	7(53.3%) Bacterial pneumonia 2(15.3%) Tuberculosis	4(30%)		
Fungal pneumonia	5	2(40%)	3(60%)		
ILD	5	2(40%)	3(60%)		
Total	60	38(63.3%)	22(36.7%)		

Advantage of bronchoscopy in suspected patients with sputum negative pulmonary tuberculosis, isolation of mycobacteria at an early stage when the destruction of lung tissue is minimal and the risk of spreading the disease to contacts can be decreased by early diagnosis and treatment⁷. Bronchoscopy reveals a higher bacteriological

confirmation of diagnosis in patients with strong clinical and radiological evidence suggestive of Pulmonary Tuberculosis, Non resolving pneumonia and Interstial lung disease. Though FOB procedures have some risk of complications, it is considered to be a relatively safe procedure.

Ethical Consideration :

The study was conducted after ethical permission from the Institutional Ethical Committee. The institutional ethics committee has gone through the study proposal and had unanimously apporved this study. The study participants had been explained the nature and purpose of the study and written consent was obtained.

Limitations of study :

Sample size of the study is small beacause of the covid 19 pandemic and this study may not represent the whole indian populations due to smaller sample size and exclusion of patients with HIV infection, Bleeding disorders and suspected Malignancy

CONCLUSION

Role of Fiberoptic Bronchoscopy in malignant lung disorder is well established however, our study suggests that fiberoptic bronchoscopy can provide excellent material for diagnosis of suspected benign lung disorders.

Broncho alveolar Lavage and Bronchial Brushing subjected for AFB stain, CBNAAT, TB Culture, Cytology, KOH stain, NTM Culture etc. shall be cumulatively use for increase diagnostic yield of various etiology.

REFERENCES

- Kovnat DM, Rath GS, Anderson WM, Snider GL Maximal extent of visualization of bronchial tree by flexible fiberoptic bronchoscopy. *Amerrevrespdis* 1974; **110(1):** 88-90.
- Suratt PM, Smiddy JF, Gruber B Deaths and Complications Associated with Fiberoptic Bronchoscopy. *Chest [Internet]* 1976 Jun 1 [cited 2021 Sep 4]; 69(6): 747-51.



Fig 2 — Diagnostic yield of Bronchoscopy in suspected different Benign Lung Disease

- 3 Stover DE, Zaman MB, Hajdu SI, Lange M, Gold J, Armstrong D — Bronchoalveolar lavage in the diagnosis of diffuse pulmonary infiltrates in the immunosuppressed host. *Ann Intern Med* 1984; **101(1):** 1-7.
- 4 Narain R, Nair SS, Naganna K, Chandrasekhar P, Rao GR, Lal P — Problems in defining a "case" of pulmonary tuberculosis in prevalence surveys. *Bull World Health Organ* 1968; 39: 701-29
- 5 Shamim S, Agarwal A, Ghosh BK, Mitra M Fungal pneumonia in intensive care unit: When to suspect and decision to treatment: A critical review. J Assoc Chest Physicians [serial online] 2015 [cited 2023 Apr 8]; 3: 41-7.
- 6 Luhadia A, Luhadia SK, Jain S, MH Hanfe, Oza D, Zota A, et al Role of fiberoptic bronchoscopy in sputum smear negative suspected cases of pulmonary tuberculosis: a study conducted in Southern part of Rajasthan. *IJRMS* 2020; 8(12): 4302-6.
- 7 Quaiser S, Agarwal A, Khan R, Haque SF Fiberoptic bronchoscopy, as a valuable diagnostic option in sputum negative pulmonary tuberculosis: A prospective study. *Int J Appl basic Med Res [Internet]* 2012 [cited 2021 Oct 25]; **2(2)**: 123.
- 8 Jo YS, Park JH, Lee JK, Heo EY, Chung HS, Kim DK Discordance between MTB/RIF and real-time tuberculosisspecific polymerase chain reaction assay in bronchial washing specimen and its clinical implications. *PLoS One* 2016; **11(10).**
- 9 Baughman RP, Dohn MN, Loudon RG, Frame PT Bronchoscopy with bronchoalveolar lavage in tuberculosis and fungal infections. *Chest [Internet]* 1991 [cited 2021 Oct 25]; 99(1): 92-7.
- 10 Surya kumari V, Pathrudu BMS, Sunil Kumar R, Rao GS, Jakka S, Alavala SC, et al — Clinical and Etiological Profile of Unresolving Pneumonia Cases Attending Government Chest Hospital, Visakhapatnam" IOSR. Journal of Dental and Medical Sciences (IOSR-JDMS) 2015; 14(7): Ver. VII (July.2015), 56-61
- 11 Shin JA, Chang YS, Kim TH, Kim HJ, Ahn CM, Byun MK Fiberoptic bronchoscopy for the rapid diagnosis of smearnegative pulmonary tuberculosis. *BMC Infect Dis* 2012; 121 [Internet]. 2012 Jun 22 [cited 2021 Oct 25]; **12(1):** 1-7.
- 12 Bachh AA, Gupta R, Haq I, Varudkar HG Diagnosing sputum/smear-negative pulmonary tuberculosis: Does fibre-optic bronchoscopy play a significant role? *Lung India [Internet]* 2010 Apr 1 [cited 2021 Oct 25]; 27(2): 58-62.
- 13 Sarkar SK, Sharma GS, Gupta PR, Sharma RK Fiberoptic bronchoscopy in the diagnosis of pulmonary tuberculosis. *Tubercle [Internet]*. 1980 [cited 2021 Oct 25]; 61(2): 97-9.
- 14 Purohit SD, Sisodia RS, Gupta PR Fiberoptic bronchoscopy in the diagnosis of smear negative pulmonary tuberculosis. *Lung India* 1983; 1: 143-6.

Original Article

A Study of the Clinico-demographic Profile and Clinical Outcomes of COVID-19 Patients in a Tertiary Care Hospital of India

Rupankar Nath¹, Deepannita Sutradhar², Riturag Thakuria³

Background : The SARS-CoV-2-induced COVID-19 pandemic has presented substantial challenges to healthcare systems across the globe. This study aimed to investigate the clinico-demographic profile and clinical outcomes of COVID-19 patients in a tertiary care hospital in Barak Valley, Assam, India.

Methodology : A retrospective, single-center, hospital-based cross-sectional study was conducted, focusing on patients with laboratory-confirmed COVID-19 admitted between July, 2020 and December, 2020.

Results : A total of 353 patients were analyzed with 69% being male and 31% female. Patients were classified into three severity groups: mild, moderate and severe. The mean ages in the mild, moderate and severe groups were 41.58 years, 57.70 years and 59.54 years, respectively. Cachar district accounted for the majority of patients (68.83%), followed by Karimganj district (18%) and Hailakandi district (10%). The common presenting complaints included fever, cough and dyspnoea. Co-morbidities were present in 61.76% of patients, with diabetes and hypertension being the most prevalent. Pulmonary comorbidities showed a strong association with increased mortality rates. Vital signs and laboratory parameters worsened with disease severity with neutrophils increasing and lymphocytes decreasing. Ferritin and LDH levels also increased with severity, reflecting disease progression. Treatment involved remdesivir and convalescent plasma therapy, with the combination showing better outcomes compared to individual therapies.

Conclusion : Overall, the study provides valuable insights into the unique challenges and characteristics of COVID-19 patients in the low HDI region of Barak Valley, Assam. The findings can aid in targeted public health interventions, resource allocation and equitable healthcare delivery, ultimately mitigating the impact of the disease and improving patient outcomes in the region. However, the study has certain limitations and further prospective multi-center studies are needed to validate and expand upon these findings.

[J Indian Med Assoc 2024; 122(12): 23-8]

Key words : COVID-19, Co-morbidity, Clinico-demographic, Mortality, Treatment.

he COVID-19 pandemic, as declared by WHO on March 11, 2020, has presented significant challenges to the health care system Worldwide. COVID-19 is a viral illness caused by SARS CoV-2, a virus of the Coronaviridae family. It was first identified in Wuhan city in Hubei province of China in December, 2019 and was later named as 2019 novel Corona Virus (2019-nCoV)¹. Similar to the other two Corona Viruses of the family, SARS-CoV and MERS-CoV, which caused outbreaks in China (2003) and Saudi Arabia (2012), SARS-CoV-2 is also believed to have originated from bats that later spread to humans through an intermediate human host, although the exact origin is still being studied^{1,2}. While the overall fatality rate of COVID-19 is lower than that of SARS or MERS³, its high transmissibility has

²MD (Anaesthesiology), Associate Professor

³MD (Medicine), Associate Professor, Department of Medicine *Received on : 11/08/2023*

Accepted on : 28/11/2023

Editor's Comment :

This study highlights the importance of demographic factors, co-morbidities, and disease severity in determining COVID-19 outcomes in Barak Valley, Assam, emphasizing the need for targeted public health measures and equitable resource allocation. The findings indicate that older age, male gender and co-morbidities, particularly pulmonary conditions, significantly increase mortality risks. Enhanced clinical strategies, including the combination of Remdesivir and convalescent plasma therapy, could improve patient outcomes in severe cases.

resulted in a significantly larger number of infections and deaths Worldwide.

It has become evident that the clinical outcome and manifestation of COVID-19 vary widely among individuals, influenced by a wide range of clinicodemographic factors such as age, co-morbidities, socio-economic status, access to health care, population density and the effectiveness of public health measures implemented by different countries and regions. Performing a clinico-demographic study of COVID-19 in Assam, a low Human Development Index (HDI) region, is essential for understanding the unique challenges and characteristics of the local

Department of Anaesthesiology and Critical Care, Silchar Medical College and Hospital, Silchar, Assam 788014

 $^{^1\}mbox{MD}$ (Anaesthesiology), Associate Professor and Corresponding Author

population. This knowledge can inform targeted public health interventions, resource allocation and equitable healthcare delivery, ultimately helping to mitigate the impact of the disease and improve outcomes in the region. To the best of our knowledge, this is the first comprehensive study about the clinico-demographic profile of COVID-19 from Barak Valley, Assam. The primary objective of the study was to analyze the clinico-demographic profile of admitted COVID-19 patients and secondarily to correlate the association of age, sex and co-morbidities with mortality.

MATERIALS AND METHODS

Study Setting and Duration :

This retrospective, single-center, hospital-based cross-sectional study was conducted in a tertiary care hospital situated in Barak Valley, Assam. The study focused on the population of patients who were admitted to the hospital with laboratory-confirmed COVID-19 between July, 2020 and December, 2020.

Sampling Method :

The sampling approach employed was nonprobability convenience sampling, meaning all consecutive patients admitted during the specified study duration were included.

Inclusion and Exclusion Criteria :

All hospitalized patients with laboratory confirmed, COVID-19 (RAT or RTPCR) who consented to participate were included in the study. Duplicate samples, meaning repeat samples taken from patients who were already confirmed positive for COVID-19, were excluded from the study. Additionally, incompletely filled Surveillance Report Form (SRF) forms and/or incomplete inpatient hospital tickets were excluded from the analysis.

Data Analysis :

To gather the necessary data for analysis, clinicodemographic information about the patients was collected from the SRF forms and hospital bed tickets. This data was systematically entered into a predesigned form and further uploaded into a Microsoft Excel® sheet for analysis. Statistical correlation was done using GraphPad Prism® version 8.

RESULTS

This study involved the analysis of a total of 353 patients, comprising 246 Males (69%) and 107 Females (31%). The patients were categorized into three groups based on the severity of their condition: mild, moderate and severe, according to state protocol. 100 patients (28.3%) were classified as Mild,

169 (47.87%) as Moderate and 84 as severe (23.8%). The mean age of patients in the mild group was 41.58 years, while it was 57.7 years in the moderate group and 59.54 years in the severe group. Upon admission, 239 patients (67.7%) tested positive for Rapid Antigen Testing (RAT), while the remaining 114 patients (32.3%) tested positive for RT-PCR (Table 1).

The majority of patients seeking admission at Silchar Medical College were from Cachar district 68.83% (n=243), followed by 18% (n=65) from Karimganj and 10% (n=35) from Hailakandi district. The mortality of patients increased with increasing distance from the treating hospital (Table 2).

The most common presenting complaint among the patients was fever, reported by 61.76% (n=218) of patients, followed by cough in 53.82% (n=190) of patients. Other symptoms included dyspnoea in 42.78% (n=151) of patients, sore throat in 7.36% (n=26), diarrhoea in 4.25% (n=15), chest pain in 3.4% (n=12), headache in 3.68% (n=13), and loss of taste and smell in 0.28% (n=1) of patients (Fig. 1).

Table 1 — Demographic profile of COVID-19 patients						
Parameter Mild Moderate Severe						
Total Patients	100	169	84			
Male	65	122	59			
female	35	47	25			
Average Age (in years)	41.58	57.7	59.54			
Rat positive	239					
RT-PCR positive	114					

Table 2 — Demographic Distribution of cases and outcome								
	comparison							
Place Total Patients Mortality Mortality Percentage								
Cachar	243	37	15.22					
Hailakandi	35	7	20.00					
Karimganj	65	19	29.2					
Tripura	6	2	40.00					
Others	4	2	50.00					



Co-morbidities were present in 218 (61.76%) patients, out of which 48 (22%) expired. 19 (15.97%) patients expired who had no co-morbidities. Out of 14 patients with pulmonary comorbidities, 6 (42.86%) expired. The mortality in patients with diabetes, hypertension (HTN), or both as co-morbidities was 19.09% (n= 21). In patients with diabetes, HTN or both along with additional comorbidities was 19.72% (n=14). In patients without



Journal of the Indian Medical Association

Fig 2 — Association between co-morbidities and mortality

diabetes or HTN but with other co-morbidities, the mortality was 30.43% (n=7). (Fig 2).

Among patients with severe disease, 20.96% (n=74) patients were having age more than 40 and 2.83% (n=10) patients were less than 40. The mortality among patients with an age more than 40 was 17% (n=60) and mortality at age below 40 was 1.98% (n=7). The percentage of Males in severe group was 70.23% (n=59) and Females 29.77% (n=25) while 21.14% (n=52) were males as compared to 22.39% (n=15) females out of the total mortality (n=67) (Table 3).

The vital signs and laboratory parameters varied across the three severity categories. In the mild category, patients had an average respiratory rate of 19.56 breaths per minute, oxygen saturation (SpO2) of 96.8%, heart rate of 92.7 beats per minute, and a mean blood pressure of 75 mm Hg. In the moderate category, the average respiratory rate was 23.75 breaths per minute, SpO2 was 92%, heart rate was 101.3 beats per minute and mean blood pressure was 70 mm Hg. In the severe category, the average respiratory rate was 25.86 breaths per minute, SpO2 was 72.8%, heart rate was 125.8 beats per minute, and mean blood pressure was 50 mm Hg (Table 4).

The average Hb levels were 12.46 gm%, 11.8 gm% and 11.39 gm% in mild, moderate and severe group respectively. The total count increased with severity from 6281/microliters in mild group to 14329/microliters in severe group with neutrophilia progressively increasing from 62.82% per mm³ in mild group to 85% per mm³ in severe group. Lymphocytes progressively decreased with severity from 25.5% per mm³ in mild group to 8.12% per mm³ in severe group. Furthermore, the severity of the patients' conditions

Table 3 — Association of Age and sex with mortality						
	Severity	Percentage	Mortality	Percentage		
Age >40	74/353	20.96	60/353	17.00		
Age <40	10/353	2.83	7/353	1.98		
Male	59/84	70.23%	52/67	77.61		
Female	25/84	29.77%	15/67	22.39		

Table 4 — Vital parameters of admitted patients						
Mild Moderate Severe						
RR	19.56	23.75	25.86			
SPO2%	96.8	92	72.8			
HR (bpm)	92.7	101.3	125.8			
Mean BP (mmHg)	75	70	50			
RR - Respiratory Rate (breaths per minute); HR - Heart Rate; BP - Blood Pressure						

was reflected in their ferritin and LDH levels, as well as the findings from Chest X-rays. In the mild group, ferritin levels averaged at 129.7 ng/mL, while LDH levels were 391 IU/L. Among the patients in this group, 47 individuals had abnormal Chest X-ray findings. In the moderate group, ferritin levels increased to an average of 330 ng/mL and LDH levels rose to 436 IU/ L. The number of patients with abnormal chest X-ray findings also increased to 107. In the severe group, ferritin levels further increased to an average of 416 ng/mL and LDH levels were notably elevated at 594 IU/L. The number of patients with abnormal Chest Xray findings remained significant, with 33 individuals displaying such findings (Table 5).

Treatment for the patients followed the state protocol, which included symptomatic care. Two major modalities were employed: Pharmacotherapy with Remdesivir and experimental Convalescent Plasma Therapy (CPT). Different treatment combinations were administered based on the severity of the disease. Specifically, 11 only patients received convalescent plasma therapy and 6 of them (54.54%) did not survive. Remdesivir alone was administered to 126 patients, of whom 31 (24.60%) expired. Among those who received both CPT and Remdesivir (91 patients), the mortality rate was 21.90% with 20 patients expiring. 25 patients did not receive either CPT and Remdesivir of whom 10 patients (40%) expired (Table 6).

	Table 5 — Laboratory and Radiology Parameters in different groups							
Parameters Hb gm%		TLC p microlit	er Lympl ers per	hocytes mm ³	Neutrophils per mm ³	Ferritin ng/mL	LDH IU/L	Positive Chest x-ray
Mild	12.46	6281	1 25	5.55	62.82	129.7	391	47
Moderate	11.87	8918	3 16	3.26	75.88	330	436	107
Severe	11.39	1432	98	.12	85	416	594	33
Hb - Haei	Hb - Haemoglobin; TLC - Total Leucocyte Count							
Table 6 — Outcome comparison after experimental therapies CPT- Convalescent Plasma Therapy								
CF	CPT Remdesivir CPT+ Remdesivir No CPT No Remdesivir							
Total	Expired	Total	Expired	Total	Expired	Total	E	xpired
Patients	(%)	Patients	(%)	Patient	s (%)	Patient	s	(%)
11	6(54.54)	126	31(24.60)	91	20(21.98)	25		10(40)

Journal of the Indian Medical Association

In terms of outcome, within the mild group, 90 patients were successfully cured, 7 remained admitted at the time of data collection and 3 were discharged against medical advice. There were no deaths in the group. In the moderate group, 23 individuals expired, while 123 patients recovered. Additionally, 17 patients were still admitted and 4 patients left against medical advice. Within the severe group, 44 patients expired while 32 were successfully cured, additionally, 5 patients remained admitted and 1 patient left against medical advice (Fig 3).

DISCUSSION

Demographically, the study revealed that the majority of patients seeking admission to Silchar Medical College were Male (69%) compared to Female patients (31%). This observation aligns with previous studies that have also shown a higher susceptibility of males to COVID-19⁴⁻⁶. The underlying reasons for this gender disparity are not fully understood and may involve biological such as the higher expression of ACE2 receptor⁷, behavioural such as smoking, excessive alcohol consumption and less adherence to preventive measures like wearing



Fig 3 — Outcome comparison among different groups

mask and social distancing and societal factors such as occupation and workplace exposure and traditional social roles like sharing the majority of outdoor responsibility^{5,8-10}. The severity and mortality of the illness also had a male preponderance in our study. However, there is disparity in observations with few studies showing positive correlation between severity as well as mortality and male gender like^{6,11-17} and others showing negative or no correlation with male gender⁸. It is important for future research to investigate the mechanisms behind this disparity to form targeted interventions and public health strategies

Our study confirms previous findings that increased age is linked with higher disease severity and mortality¹⁹. This population requires particular focus on prevention, early detection and management strategies.

Furthermore, the study reported that the majority of patients seeking admission were from Cachar district (68.83%), followed by Karimganj (18%) and Hailakandi (10%) districts. It was also seen that both the disease severity and mortality were higher among cases from Karimganj and Hailakandi. These regional variations might be influenced partly by factors such as population density, accessibility to healthcare facilities and the level of awareness about COVID-19 in different regions and also because most of the cases from these two districts were referred. The study provides important regional data that can aid in resource allocation and public health planning at the local level though further studies with particular focus on these factors are awaited.

The common presenting complaints in our study were fever, cough and dyspnoea in the descending order of presentation followed by sore throat, diarrhoea, chest pain, headache and loss of taste and smell the last being the least frequent symptom. These findings are consistent with the typical symptoms associated with COVID-19, as reported by various studies^{6,20,21}.

Co-morbidities were found to be present in a significant proportion of the patient, among which diabetes and hypertension were the most prevalent. It was also observed that Pulmonary Co-morbidity was strongly associated with increased mortality rates, surpassing other Co-morbidities in terms of mortality risk. These findings are consistent with previous studies that have highlighted the association between these Co-morbidities and increased morbidity and mortality in COVID-19^{5,17,22,23}. It is thus crucial for healthcare providers to be aware of and address these underlying Co-morbidities in the management of COVID-19 patients.

Increasing disease severity correlates with escalating abnormalities in vital signs particularly increase in respiratory rate, SpO2 and blood pressure reflecting respiratory distress and multiorgan dysfunction in severe COVID-19. A progressive decrease in the lymphocyte count and an increase in the neutrophil count, LDH and ferritin levels with increasing disease severity was also observed as a finding that aligns with previous studies^{21,24}.

The outcome in terms of mortality was seen to be favourable in patients who received a combination of plasma therapy and Remdesivir as compared to those receiving either of the two therapies though outcome was slightly better in Remdesivir group as compared to Plasma therapy alone. This finding of ours was in accordance with another study by Diaz G²⁵. Further focussed Randomized controlled trials are required to establish the efficacy and safety of these treatment modalities.

It was observed that the severe group had the highest mortality rate which highlights the critical need for timely and appropriate management of severe COVID-19 cases to improve patient outcomes.

It is important to acknowledge the limitations of this study. First, the study design was retrospective, which introduces inherent biases and limitations in data collection. Prospective studies with larger sample sizes and control groups are needed to further validate and expand upon these findings. Second, the study was conducted at a single centre, which may limit the generalizability of the results to other settings. Multi centre studies involving diverse populations are necessary to obtain a more comprehensive understanding of COVID-19 characteristics and outcomes.

CONCLUSION

This study at Silchar Medical College and Hospital highlights the impact of factors like demographics, Co-morbidities, vital signs and treatment outcomes on COVID-19 severity and mortality in the population of Barak valley. The findings can inform Government policies for improved clinical management and public health interventions. Further research is needed to validate and explore additional factors influencing COVID-19 outcomes.

REFERENCES

- 1 The origin of SARS-CoV-2 Burki, Talha. *The Lancet Infectious Diseases*, **20(9):** 1018-9.
- 2 Rabaan AA, Al-Ahmed SH, Haque S, Sah R, Tiwari R, Malik YS, et al — SARS-CoV-2, SARS-CoV and MERS-COV: A comparative overview. *Infez Med* 2020.
- 3 Petrosillo N, Viceconte G, Ergonul O, Ippolito G, Petersen E — COVID-19, SARS and MERS: are they closely related? *Clinical Microbiology and Infection* 2020; **26(6):** 729-34.
- 4 Prasad AK, Khyriem AB, Lyngdoh WV, Lyngdoh CJ, Phukan AC, Bhattacharya PK, et al Clinico-demographic profile of COVID-19 positive patients first wave versus second wave an experience in north-east India. J Infect Dev Ctries 2023; 17(2): 166-77.
- 5 Jamil M, Bhattacharya PK, Barman B, Topno N, Barman H, Nongpiur VN, et al — Clinical and Demographic Profile of COVID-19 Patients: A Tertiary Level Hospital-Based Study From Northeast India. *Cureus [Internet]*. 2021 Oct 19 [cited 2023 Jun 21]; 13(10). Available from: https:// pubmed.ncbi.nlm.nih.gov/34820212/
- 6 Tiwari L, Gupta P, N Y, Banerjee A, Kumar Y, Singh PK, et al — Clinicodemographic profile and predictors of poor outcome in hospitalised COVID-19 patients: a single-centre, retrospective cohort study from India. BMJ Open 2022; 12(6): e056464.
- 7 Jin JM, Bai P, He W, Wu F, Liu XF, Han DM, *et al* Gender Differences in Patients With COVID-19: Focus on Severity and Mortality. *Front Public Health* 2020 Apr 29; 8.
- 8 Pradhan A, Olsson PE Sex differences in severity and mortality from COVID-19: are males more vulnerable? Vol. 11, Biology of Sex Differences. BioMed Central Ltd; 2020.
- 9 Mohamed MO, Gale CP, Kontopantelis E, Doran T, de Belder M, Asaria M, et al — Sex Differences in Mortality Rates and Underlying Conditions for COVID-19 Deaths in England and Wales. *Mayo Clin Proc* 2020; **95(10):** 2110-24.
- 10 Bwire GM Coronavirus: Why Men are More Vulnerable to COVID-19 Than Women? Available from: https://doi.org/ 10.1007/s42399-020-00341-w
- Shah S, Singhal T, Davar N, Thakkar P No correlation between Ct values and severity of disease or mortality in patients with COVID 19 disease. *Indian J Med Microbiol* 2021; **39(1):** 116-7.
- 12 Soni S, Kajal K, Yaddanapudi L, Malhotra P, Puri G, Bhalla A, et al — Demographic & amp; clinical profile of patients with COVID-19 at a tertiary care hospital in north India. Indian Journal of Medical Research. 2021;153(1):115.

- 13 Ortiz-Prado E, Simbaña-Rivera K, Barreno LG, Diaz AM, Barreto A, Moyano C, et al — Epidemiological, socio-demographic and clinical features of the early phase of the COVID-19 epidemic in Ecuador. *PLoS Negl Trop Dis* 2021; **15(1)**: e0008958.
- 14 Biccard BM, Gopalan PD, Miller M, Michell WL, Thomson D, Ademuyiwa A, et al — Patient care and clinical outcomes for patients with COVID-19 infection admitted to African high-care or intensive care units (ACCCOS): a multicentre, prospective, observational cohort study. *The Lancet* 2021; **397(10288)**: 1885-94.
- 15 Grasselli G, Zangrillo A, Zanella A, Antonelli M, Cabrini L, Castelli A, et al — Baseline Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to ICUs of the Lombardy Region, Italy. JAMA 2020; 323(16): 1574.
- 16 Cummings MJ, Baldwin MR, Abrams D, Jacobson SD, Meyer BJ, Balough EM, *et al* — Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *The Lancet* 2020; **395(10239)**: 1763-70.
- 17 Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, *et al* — Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* 2020; **323(20)**: 2052.
- 18 Hamza A, Shah N, Azad A, Ghanshyam O, Khan Z Impact of age, gender and comorbidities affecting the severity of COVID-19 infection in Kashmir. *J Family Med Prim Care* 2022; 11(4): 1519.

- 19 Wu Z, McGoogan JM Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China. JAMA 2020; 323(13): 1239.
- 20 Gallo Marin B, Aghagoli G, Lavine K, Yang L, Siff EJ, Chiang SS, et al — Predictors of <scp>COVID</scp> 19 severity: A literature review. *Rev Med Virol* 2021; **31(1):** 1-10.
- 21 Guan W jie, Ni Z yi, Hu Y, Liang W hua, Ou C quan, He J xing, et al — Clinical Characteristics of Coronavirus Disease 2019 in China. New England Journal of Medicine 2020; 382(18): 1708-20.
- 22 Parohan M, Yaghoubi S, Seraji A, Javanbakht MH, Sarraf P, Djalali M — Risk factors for mortality in patients with Coronavirus disease 2019 (COVID-19) infection: a systematic review and meta-analysis of observational studies. *The Aging Male* 2020; **23(5):** 1416-24.
- 23 Singh AK, Misra A Impact of COVID-19 and comorbidities on health and economics: Focus on developing countries and India. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* 2020; **14(6):** 1625-30.
- 24 Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* 2020; **395(10223):** 497-506.
- 25 Diaz GA, Christensen AB, Pusch T, Goulet D, Chang SC, Grunkemeier GL, et al — Remdesivir and Mortality in Patients With Coronavirus Disease 2019. *Clinical Infectious Dis*eases 2022; **74(10):** 1812-20.

If you want to send your queries and receive the response on any subject from JIMA, please use the E-mail or Mobile facility.

Know Your JIMA

Website :	https://onlinejima.com
	www.ejima.in
For Reception :	Mobile : +919477493033
For Editorial :	jima1930@rediffmail.com
	Mobile : +919477493027
For Circulation :	jimacir@gmail.com
	Mobile : +919477493037
For Marketing :	jimamkt@gmail.com
	Mobile : +919477493036
For Accounts :	journalaccts@gmail.com
	Mobile : +919432211112
For Guideline :	https://onlinejima.com

Original Article

Levels and Risk Factors of Diabetes Mellitus among Tuberculosis Patients Attending District Tuberculosis Centre, Vijayapur

Mallikarjun C Yadavannavar¹, Tanuja P Pattankar², Daneshwari P Kouttalmatt³

Background : The co-existence of diabetes and tuberculosis is becoming a public health problem in India, as they are the cause of morbidity, mortality and pose a burden on healthcare system.

Aims and Objectives : To assess the risk factors of diabetes mellitus and its status among Tuberculosis patients. To compare the response to treatment and drug resistance among diabetic and non-diabetic patients. And to study the prevalence of MDR-TB among Diabetic patients with Tuberculosis.

Methodology: It was a cross sectional study done in Vijayapura district during June, 2019 to July, 2019. The patients who had attended the District Tuberculosis Centre in previous 3 months of the beginning of the study were considered and data was collected from the records maintained in the center. The sample size calculated to be 300 cases. Data was collected from them.

Results : The majority of 21.7% of study participants belonged to the age group 26-35 years. The overall prevalence of diabetes was 21.6 % among TB patients. Among males the prevalence was 67.3% and among females was 32.7%. When prevalence of diabetes mellitus was compared with the age it was found that majority (30.7%) of the diabetes were in the age group of 56-65 years and it was statistically significant.

Conclusion : Diabetes is a prevalent comorbidity in both drug sensitive and drug resistant tuberculosis. Age group of more than 55 years is significantly associated with diabetes mellitus and other factors like alcohol consumption is also a associated factor.

[J Indian Med Assoc 2024; 122(12): 29-34]

Key words : TB, Diabetes Mellitus, MDR TB.

orldwide Tuberculosis being one among the top 10 leading cause of death¹. 24% of TB patients & 29% deaths are due to TB in India. The factors attributed for this are HIV, Diabetes Mellitus, Smoking & Malnutrition². Among 10.4 million new active TB case, one million of them have TB-DM comorbidities³. There is an evidence showing association between diabetes and tuberculosis referred as 'intersecting epidemic' by WHO⁴. According to International Federation of Diabetes, about 382 million adults had Diabetes in the year of 2013, among which 80% belong to low-middle income countries, with prediction of increase in global burden reaching 592 million by 2035⁵. India is stepping towards becoming the Diabetic capital, as every fifth diabetic in the World is Indian⁶. Diabetes increases the risk of active infection, progression of latent to active case and transmission of infection, besides these it has led to poor outcome

Received on : 27/08/2022

Accepted on : 08/06/2024

Editor's Comment :

 What's Already Known : Diabetes Mellitus (DM) is a significant comorbidity among tuberculosis (TB) patients, contributing to worse treatment outcomes and complications. The association between DM and TB is well-established globally, but regional variations in prevalence and risk factors remain underexplored.
 What This Study Adds : This study reports a 21.6% prevalence of diabetes among TB patients in Northern Karnataka, identifying key risk factors such as age, education level, alcohol consumption and family bioter.

history of diabetes. It underscores the dual burden of TB and DM in the region.
How It's Useful for Policy : The findings advocate for integrated screening and management of both TB and diabetes. Policy should promote bidirectional screening to improve treatment outcomes and

reduce complications, enhancing public health interventions in areas with high TB burden. of treatment with increased risk of failure of treatment⁴. In addition glycemic control is being worsened by TB infection. Added to this Drug interaction is deterioting the effectiveness of both TB and Diabetes⁷. DM also increases the risk of TB by 3 times and thus rendering for slow decline of global TB incidence which may hinder the goal of achieving

global milestone of 50% reduction in TB incidence and 75% reduction of deaths due to TB by 2025⁸. Hence, this bidirectional association between TB-DM

Department of Community Medicine, BLDE(DU) Shri B M Patil Hospital, Medical College and Research Center, Vijayapura, Karnataka 586103

 $^{^1\}mbox{MD}$ (Community Medicine), Professor and Head and Corresponding Author

²MD (Community Medicine), Assistant Professor ³MBBS, Student

is being major concern for physicians, as DM affects the disease prognosis and treatment outcome of TB vice versa^{9,10}. Bidirectional screening and integrated management will help in early diagnosis and health outcomes of both conditions, unfortunately there is inadequate evidence for this on feasibility and effectiveness of this approach⁴. The impact of these two converging epidemics has led WHO to declare Tuberculosis and Diabetes as global epidemics¹¹.

Apart from this the emergence of Multi Drug Resistant Tuberculosis (MDR-TB) across the world is posing a major threat, as treatment is difficult and expensive hence causing a large economic burden on the developing nations. In addition to this Diabetes is an added risk factor for the development of MDR-TB because many studies have shown increased risk of MDR-TB among Tuberculosis patients with comorbid Diabetes Mellitus, although Diabetes Mellitus is not an independent risk factor for MDR-TB¹²⁻¹⁷.

Tuberculosis and Diabetes Mellitus co-morbidity is one of the raising public health problems. Data available regarding tuberculosis co-morbid with diabetes mellitus is very sparse in this part of Northern Karnataka, hence the study was undertaken.

AIMS AND OBJECTIVES

(1) To study the risk factors of diabetes mellitus and its status among Tuberculosis patients.

(2) To compare the response to treatment and drug resistance among diabetic and non-diabetic patients.

(3) To study the prevalence of MDR-TB among Diabetic patients with Tuberculosis.

MATERIALS AND METHODS

It is a Cross-sectional study done in 2 months from June, 2019 to July, 2019. The patients who had attended the District Tuberculosis Centre in previous 3 months of the beginning of the study were considered and data was collected from the records maintained in the centre and also by contacting the patient (after obtaining their consent) with the help of contact details available in the records for further information. With anticipated prevalence rate of diabetes among TB cases is 24%¹⁸ at 95% confidence level and 5% absolute error, sample size calculated is 280.

Using statistical formula : $n=Z^2p^*q/d^2$, 280 sample size was obtained which was rounded to 300 cases.

Tuberculosis patients (18 years and above) who attended District Tuberculosis Centre for past 3 months are included in the study. Pregnant lactating mothers and patients with incomplete records were excluded from the study. **Data Analysis :** The data obtained will be entered in a Microsoft Excel sheet and statistical analysis will be performed using statistical package for the social sciences (Version 20). Results will be presented as Mean (Median) ±SD, counts and percentages and diagrams. Categorical variables will be compared using Chi square test. Association between psychosocial changes and resiliency factors will be assessed using regression analysis, p<0.05 will be considered statistically significant. All statistical tests will perform two tailed.

RESULTS

Table 1 shows that out of 300, majority (21.7%) of them belonged to the age group 26-35 years followed by 56-65 years (21%) and 36-45 years (17%).This shows that most of population belong to the reproductive age group. In our study males were predominant (64.3%) and female constituted about (35.7%).

Majority of study participants belong to urban (51.7%) when compared to rural background (49.3%). Most of the study participants studied upto Primary

Table 1 — Distribution of Socio-demographic Characteristics of Study Populations (N=300)							
Study variable Number(n) Percentage (%)							
Age (years) :							
15-25	43	14.3					
26-35	65	21.7					
36-45	51	17					
46-55	48	16					
56-65	63	21					
>65	29	9.7					
Sex :							
Male	193	64.3					
Female	107	35.7					
Place :							
Rural	145	48.3					
Urban	155	51.7					
Education Status :							
Illiterate	44	14.7					
Primary	162	54					
Secondary	73	24.3					
PUC	20	6.7					
Graduate	1	0.3					
Marital Status :							
Married	268	89.3					
Unmarried	19	6.3					
Widowed	13	4.3					
Type of Family :							
Nuclear	285	95					
Joint	15	5					
Socio-economic Status	:						
Upper class	7	2.3					
Upper middle class	10	3.3					
Middle class	25	8.3					
Lower middle class	93	31					
Lower class	165	55					

school (54%), followed by secondary school (24.3%), 14.7% were illiterates.

89.3% of them were married , 6.3% were unmarried and 4.3% widowed participated in study. 95% belonged to nuclear family and only 5% belonged to joint family.

In our study 55% belonged to lower class followed by lower middle class (31%) (Modified BG Prasad's classification).

In our study among 300 TB patients 65 were Diabetic (21.6 %)(Fig 1). Prevalence of diabetes in male and female is 67.3% and 32.7% respectively (Fig 2)(Table 2).

When prevalence of diabetes mellitus was compared with the age it was found that majority (30.7%) (Fig 3) of the diabetes were in the age group of 56-65 years and it was statistically significant with the p value of <0.001 (Table 3).

On comparing the association between prevalence of diabetes mellitus and educational status, it was found that among diabetes most of the study

DIABETIC





Fig 2 — Distribution of Male and Female Among Diabetic and Non Diabetic TB Patients

Table 2 — Association of Prevalence of DM and Age								
Age	Diabetic		Non Diabetic		Chi square	e P value		
(Years)	n	%	n	%	test			
15-25	0	0	43	18.3				
26-35	2	3.1	63	26.8	χ²=60.252	P<0.0001		
36-45	8	12.3	43	18.3		HS		
46-55	18	27.7	30	12.8				
56-65	20	30.7	43	18.3				
>65	17	26.1	12	5.1				
Total	65	100	234	100				
HS - Highly significant								



Fig 3 — Distribution of Male and Female among Diabetic and Non Diabetic TB Patients

Table 3 — Association of Prevalace of DM and Educational Status								
Educational	Dia	abetic	Non Diabetic		Chi square	P value		
Status	n	%	n	%	test			
Illiterate	14	21.5	30	12.75	χ ² =	P=		
Primary	43	66.15	119	50.6	16.521	0.0024		
Secondary	4	6.15	69	29.36		HS		
PUC	4	6.15	16	6.8				
Graduate	0	0	1	0.4				
Total	65	100	235	100				
HS - Highly significant								

participants (66.2%)(Fig 4) studied upto primary school and it was statistically found significant (Table 4] ie, illiteracy and primary education was significantly associated (Table 4).

The prevalence of was more in non-alcoholic 70.8% (Fig 5) and this was found statistically significant (Table 5). Hence, consumption of alcohol was not risk factor in our study.

We found that prevalence of diabetes was not significant when compared with BMI (Table 6).

89.2% of Diabetes Mellitus prevalence was found

Table 4 — Association of Prevalance of DM and Alcohol Consumption								
Habit of	Diab	etic	Non	Diabetic	Chi square	P value		
Alcohol	n	%	n	%	test			
Alcoholic	19	29.4	28	11.9	$\chi^2 =$	P=		
Non Alcoholic	46	70.6	207	88.1	11.55	0.0007		
Total	65	100	235	100		HS		
HS - Highly significant								



Alcohol intake

Fig 5 — Distribution of Alcohol Intake among Diabetic and Non Diabetic TB Patients

Table 5 — Association of Prevalnce of Diabetes and Body Mass Index								
BMI	Dia	betic	Non	Diabetic	Chi square	P value		
	n	%	n	%	test			
Underweight	40	61.5	177	75.3	$\chi^2 =$	P=		
Normal	24	36.9	55	23.4	4.897	0.0864		
Overweight/	1	1.59	3	1.27		NS		
Obese								
Total	65	100	235	100				
NS - Not significant								

among patients not having family history of DM [Figure no.07] and was statistically significant (Table 7).

We found that prevalence of DM was not significant when compared to HIV status (Fig 8).

DISCUSSION

The present study was conducted to determine the levels and risk factors of Diabetes mellitus among Tuberculosis patients registered under RNTCP in District Tuberculosis centre Vijaypura.

In our study it was found that large number of TB patients were in the age group 26-55 years (21.7%) most of them were in reproductive age group. Similar results were found in a study done by Damtew E, *et al*¹¹ in Addis Ababa Ethiopia where majority of patients were in age group 25-44 years of age.

In study done by Balakrishnan S, et al¹⁹ in Kerala

Table 6 — Association of Family History and Prevalence of Diabetes among TB Patients								
Family	Diabetic		Non Diabetic		Chi square	P value		
History	n	%	n	%	test			
No	58	90.2	232	98.7	χ ² =	P=		
Yes	7	9.8	3	1.3	14.239	0.0002		
Total	65	100	235	100		HS		
HS - Highly significant								
Table 7 — A	Associa	ation of	Preval	ence of	DM and HI\	/ Status		
Table 7 — A HIV	A <i>ssocia</i> Diab	a <i>tion of</i> etic	<i>Preval</i> Non D	<i>ence of</i> Diabetic	<i>DM and HI</i> Chi square	/ <i>Status</i> P value		
Table 7 — A HIV Status	A <i>ssocia</i> Diab n	ation of etic %	<i>Preval</i> Non D	ence of Diabetic %	DM and HIN Chi square test	/ <i>Status</i> P value		
Table 7 — A HIV Status Postive	Associa Diab n 4	etion of etic % 6.15	Preval Non D n 27	ence of Diabetic % 11.48	DM and HIN Chi square test χ²=	/ <i>Status</i> P value P=		
Table 7 — A HIV Status Postive Negative	Associa Diab n 4 61	ation of etic % 6.15 93.85	Preval Non E n 27 208	ence of Diabetic % 11.48 88.52	DM and HN Chi square test χ ² = 1.564	/ <i>Status</i> P value P= 0.2110		
Table 7 — A HIV Status Postive Negative Total	Associa Diab n 4 61 65	etic % 6.15 93.85 100	Preval Non D 1 27 208 235	ence of Diabetic % 11.48 88.52 100	DM and HN Chi square test χ ² = 1.564	/ <i>Status</i> P value P= 0.2110 NS		

most of the patients belond to the age group 45-54 years. Other studies done by Kishan, *et al* in Patiala, Punjab and Dutt N, *et al*²⁰ in Ahemdabad revealed 40-60 year's age group as most commonly involved. This difference in age group may be due to varied geographical condition and study design setting.

In our study 64.3% (193) subjects were male and 35.7% (107) were females similar to the study conducted by Tahir Z, $et a l^{21}$ in Pakistan where 69.1% were males and 30.9% were females. In study done



Fig 6 — Distribution of BMI among Diabetic and Non Diabetic TB Patients



Fig 7 — Distribution of Family History among Diabetic and Non Diabetic TB Patients



Fig 8 — Distribition of HIV Status among Diabetic and Non Diabetic TB Patients

by Alisjahbana B, et $a\beta^2$ in Indonesia in which 52.4% of patients were males. The reason is fear and stigma associated with tuberculosis which resulted in undernotification in case of females and also the tendency to seek health care.

In our study 51.7% (155) of the patients were from urban back ground and 48.3% (145) of were from rural back ground, in other study conducted by Damtew E, *et al*¹¹ in Addis Ababa, Ethiopia in which 70% of patients belong to Urban back ground. In the study done by Vishwanathan V, *et al*²³ in Tamil Nadu which had majority of Tuberculosis from urban back ground. This difference is due to different study setting.

In our study more than half 54% (162) tuberculosis patients studied up to primary schooling, 24.7% (73) of them completed high school education and 14.7% (44) were illiterates, where it was observed that most of tuberculosis patients had received less schooling which was consistent with other studies.

In contrast with the study conducted by Sarkar M, *et al*²⁴ in Bangladesh 40.7% of patients were illiterates, 25.1% received primary schooling and 19.8% of them received secondary schooling.

Another study done by Tahir Z, *et a* P1 in Pakistan has shown 51.6% as illiterates, 34.7% as primary schooling and 10.7% as secondary schooling.

In our study 89.3% (268) were married followed by $6.3\%^{19}$ unmarried and $4.3\%^{13}$ widowed, this finding was similar with study conducted by Damtew E, *et al*¹¹ in Ethiopia 59.2% were married, 36.7% were unmarried and 1.7% widowed.

In our study, 55%(165) of participants belonged to lower class, 31 %(93) belonged to lower middle class and 8.3%(25) belonged to middle class according to modified B G Prasad's Socio-economic classification. These findings were similar to the study conducted by Agarwal A K, *et al*⁷ in Madhya Pradesh in which 37.8% belonged to lower class and 15.7% belonged to lower middle class.

The prevalence of DM among TB patients in our study is 21.6%.

Other studies like institutional based cross sectional study done by Padmalatha P, $et al^{18}$ in Andra Pradesh showed prevalence of DM as 30.6%.

Another study conducted by Raghuraman S, *et al*²⁵ in Pondicherry (2017) reported 29% as DM prevalence.

In contrast to above findings, another study in Nigeria by Oliyanka AO, *et al*²⁶ found prevalence to be 5.7% which could be attributed to difference in demographic characteristics.

In our study 30.7% of Diabetes Mellitus prevalence was found among tuberculosis patients aged between 56-65 years which was similar to the study done by Kottarnath MD, *et al*²⁷ in Pariyaram Kerala where it was >61 years. In the study conducted by Ekeke N, *et al*²⁸ in Nigeria where 16.9% of DM prevalence was found among 56-65 years of age.

In our study majority (67.6%) of prevalence among males similar to the study by Agarwal AK, *et al*⁷ in Madhya Pradesh (77%) and in another study conducted by Vishwanathan V, *et al*²³. The higher prevalence of DM among males might be due to risk factors such as smoking tobacco and consumption of alcohol, which affects both TB and DM.

In our study prevalence of DM was high among who had less education (66.1% primary school and 21.5% illiterates) which is similar to the study done by Kornfeld H, *et a* P9 in South India where 51.4% of diabetic patients were illiterates. This is due to lack of awareness about diasease among less educated patients.

Our study has shown significant association between prevalence of DM and Alcohol consumption which is similar to the study conducted by Mkhontfo MM, *et a*^{β 0} in the state of Florida.

CONCLUSION

The present study showed that, the prevalence of DM among TB patients registered under RNTCP in District Tuberculosis Centre Vijayapura, was 21.6%. The DM TB were more between 56-65 years of age majority of them studied upto primary school and belonged to lower socio-economic status. Most of them were underweight due to DM TB co-morbidity. Various factors like age, educational status, addictive habits , family history of DM have influenced the prevalence.

Acknowledgement : I am thankful for ICMR STS to fund this project.

Conflict of Interest : None

REFERENCES

- 1 WHO Global tuberculosis report 2018.
- 2 PandeT, Huddart S, Xavier W, Kulavalli S, Chen T, Pai M, et al — Prevalence of diabetes mellitus amongst hospitalized tuberculosis patients at an Indian tertiary care center : a descriptive analysis. PLoS ONE 2018; 13(7) : e0200838
- 3 Shewade HD, Jeyashree K, Mahajan P, Shah AN, Kirubakaran R, Rao R, et al Effect of glycemic control and type of diabetes treatment on unsuccessful TB treatment outcomes among people with TB-Diabetes: A systematic review. PLoS ONE 2017; 12(10): e0186697.
- 4 Segafredo G, Kapur A, Robbiati C, Joseph N, de Sousa JR, Putoto G, et al — Integrating TB and non-communicable diseases services: Pilot experience of screening for diabetes and hypertension in patients with Tuberculosis in Luanda, Angola. PLoS ONE 2019; 14(7): e0218052.
- 5 IDF. IDF Diabetes ATLAS 2013 [cited 2018 Apr 27]. 6th:[Available from: www.idf.org/diabetesatlas.
- 6 Mahishale V, Avuthu S, Patil B, Lolly M, Eti A, Khan S Effect of Poor Glycemic Control in Newly Diagnosed Patients with Smear- Positive Pulmonary Tuberculosis and Type-2 Diabetes Mellitus. *Iran J Med Sci* 2017; **42(2)**: 144-51.
- 7 Agarwal AK, Gupta G, Marskole P, Agarwal A A Study of the Patients Suffering from Tuberculosis and Tuberculosis-diabetes Comorbidity in Revised National Tuberculosis Control Program Centers of Northern Madhya Pradesh, India. *Indian J Endocrinol Metab* 2017; **21(4)**: 570-6. doi:10.4103/ ijem.IJEM_89_17.
- 8 Mave V, Nimkar S, Prasad H Tuberculosis screening among persons with diabetes mellitus in Pune, India. *BMC Infect Dis* 2017; **17(1)**: 388. Published 2017 2. doi:10.1186/s12879-017-2483-9.
- 9 Siddiqui AN, Khayyam KU, Sharma M Effect of Diabetes Mellitus on Tuberculosis Treatment Outcome and Adverse Reactions in Patients Receiving Directly Observed Treatment Strategy in India: A Prospective Study. *Biomed Res Int* 2016; 2016;7273935. doi:10.1155/2016/7273935.
- 10 Nagar V, Gour D, Pal DK, Singh AR, Joshi A, Dave L A study on prevalence of diabetes and associated risk factors among diagnosed tuberculosis patients registered under Revised National Tuberculosis Control Programme in Bhopal District. J Family Med Prim Care 2018; 7(1): 130-6. doi:10.4103/jfmpc.jfmpc_289_17
- 11 Damtew E, Ali I, Meressa D Prevalence of Diabetes mellitus among Active Pulmonary Tuberculosis Patients at St. Peter Specialized Hospital, Addis Ababa, Ethiopia. World J Med Sci 2014; 11(3): 389-396.
- 12 Workneh MH, Bjune GA, Yimer SA Prevalence of associated factors of diabetes mellitus among Tuberculosis patients in South Eastern Amhara region, Ethiopia : A Cross Sectional Study. *PLoS ONE* 2016; **11(1):** 1-15.
- 13 Dobler CC, Flack JR , Marks GB Risk of Tuberculosis among people with Diabetes Mellitus : An Australian nationwide cohort study. *BMJ Open* 2012; **2**: 1-5.

- 14 Umaaknth M, Rishikesavan S Prevalence of diabetes mellitus among tuberculosis patient in Batticaloa district, Sri Lanka. Int J med Res 2017; 2(2): 21-3.,
- 15 Zheng C Hu M, Goa F Diabetes and pulmonary tuberculosis: a global overviewwith special focus on the situation in asian countries with high TBDM burden, *Global Health Action* 2017; **10(1):** 1-11.
- 16 Tahir NB, Uddin QT, Munir SS Waheed A, Waheed S, Daud HM — Prevalence of Diabetes Mellitus in tuberculosis patients at DOTS regional centre KMU-IMS teaching Hospital Kohat. *Eur Acad Res* 2014; **2(7)**: 9948-59.
- 17 Chaudhary LA, Essa EB, Solaiman Sa, Al-Sindi K Prevalence of diabetes type -2 and pulmonary tuberculosis among Filipino and treatment outcomes: A surveillance study in the eastern Saudi Arabia. *Int J Myobact* 2012; 106-9.
- 18 Padmalatha P, Hema K Study on prevalence of Diabetes Mellitus in Tuberculosis patients attending a tertiary care hospital in Guntur, Andhra Pradesh. *IJBAMR* 2014; **4(1):** 494-8.
- 19 Balakrishnan S, Vijayan S, Nair S, Subramoniapillai J, Mrithyunjaya S — High Diabetes Prevalence among Tuberculosis cases in Kerala, India. *PLoS ONE* 2012; **7(10)**: 1-7.
- 20 Kishan J, Garg K Tuberculosis And Diabetes Mellitus: A Case Series of 100 patients. *Saarc J Tuber Lung Dis* HIV/ AIDS 2010; 7(2): 34-8.
- 21 Tahir Z, Ahmed M, Akhtar AM, Yaqub T, Mushtaq MH, Javed H Diabetes mellitus among tuberculosis patients: a cross sectional study from Pakistan. *Afri Health Sci* 2016; **16(3)**: 671-6.
- 22 Alisjahbana B Diabetes mellitus is strongly associated with tuberculosis in Indonesia. Int J Tuberc Lung Dis 2006; 10(6): 696-700.
- 23 Vishwanathan V, Kumpatla S, Aravindalochanan V, Rajan R, Chinnaswamy C, Srinivasan R, et al — Prevalence of Diabetes and Pre-Diabetes and Associated Risk Factors among Tuberculosis Patients in India. PLoS ONE 2012; 7(7): 1-9.
- 24 Sarkar M, Barua M, Guerra F, Saha A, Aftab A, Latif AHMM, et al — Double Trouble: Prevalence and Factors Associated with Tuberculosis and Diabetes Comorbidity in Bangladesh. PLoS ONE 2016; 11(10): 1-15.
- 25 Raghuraman S, Vasudevan KP, Govindarajan S, Chinnakali P, Panigrahi KC — Prevalence of diabetes mellitus among tuberculosis patients in Urban Puducherry. *North Am J Med Sci* 2014; **6**: 30-4.
- 26 Olayinka AO, Anthonia O, Yetunde K Prevalence of diabetes in persons with tuberculosis in a tertiary health centre in Lagos, Nigeria. *Indian J Endocr Metab* 2013; **17**: 486-9.
- 27 Kottarnath MD, Mavila R, Achuthan V, Nair S Prevalence of diabetes mellitus in tuberculosis patients – A hospital based study. Int J Res Med Sci 2015; 3(10): 2810-4.
- 28 Ekeke N Screening for diabetes mellitus among tuberculosis patients in Southern Nigeria : a multi-centre implementation study under programme settings. Sci Rep 2017; 7: 1-8.
- 29 Kornfeld H High Prevalence and Heterogeneity of Diabetes in Patients with TB in South India. *Chest* 2016; **149(6)**: 1501-8.
- 30 Mkhontfo, Mandzisi Mbongeni An Examination of The Distribution of Diabetes Mellitus Among TB Patients with Pulmonary Tuberculosis and Drug Resistance Tuberculosis In The State of Florida, USA. (2016) [cited on Sep 20 2017]. Graduate Theses and Dissertations. Available from:http:// scholarcommons.usf.edu/edu/etd/6325

Original Article

Extent of Disability and Behavioral Disturbances of the Children with Intellectual Disability and the Coping Mechanisms of their Parents : An Observational Study from a Tertiary Care Hospital of Eastern India

Paramita Patra¹, Amitava Dan², Sharmila Sarkar³, Asish Mukhopadhyay⁴, Debasish Sanyal⁵

Background : Parents experience great challenge in bringing up their children with Intellectual Disability (ID). The present study was undertaken to examine the level of disability, magnitude and pattern of Behavioral Disturbances of the children with ID and the coping mechanism of their parents to combat against this great challenge.

Materials and Methods : The current observational study was conducted on consecutive 92 children (\leq 18 years) with mental retardation [Intellectual Disability (ID)] as per International Classification of Mental and Behavioral Disorders, Tenth Edition, Diagnostic Criteria for Research version (ICD-10, DCR) and their parents (<65 years) who was living with their children with ID. Socio-demographic data of both children and parents, Intellectual Quotient (IQ), level of disability of the children were assessed. Behavioral Disturbance of the children was assessed by Behavioral Assessment Scale for Indian Children with Mental Retardation (BASIC-MR) and coping mechanism of the parents was assessed by Brief Cope Questionnaire (BCQ).

Results : Majority of the children were male with mean age of the sample was 11.34 (SD 5.07) years with mean IQ was 79.87 (SD 11.28). Majority of the children (53%) had severe disability (\geq 90%). Children had high level of behavioral problems having higher scores in domain 1, 2 and 3 of BASIC-MR with mean scores were 105.79, 113.13 and 118.48 respectively. Majority of the parents were above 40 Years. A large portion of families came from upper middle class followed by lower middle-class background. Among the domains of BCQ, both parents had applied 'acceptance' (mean score of fathers 6.89, mothers 7.00), 'positive refraining' (mean score of fathers 5.93, mothers 6.00) and 'active coping' (mean score of fathers 6.13, mothers 6.00) mostly.

Conclusion : Parents, both the fathers and the mothers were using their coping strategies well to combat against their stress, on account of care-giving their disabled children.

[J Indian Med Assoc 2024; 122(12): 35-9]

Key words : Intellectual Disability (ID), Disability, Behavioral Disturbances, Coping Mechanisms, Parents.

According to World Health Organization, Mental Retardation (MR) or Intellectual Disability (ID) is a condition of arrested or incomplete development of the mind, characterized by impairment of skills and overall intelligence in areas such as cognition, language and motor and social abilities (WHO, 1993)¹. In developing country like India families especially parents have to take the prime role to raise their children with disabilities at home and in the society. Greater financial stress, frequent disruption

Received on : 26/11/2024

Accepted on : 27/11/2024

Editor's Comment :

- Children with intellectual disability attending Tertiary Care Hospital were at a significant level of disability and behavioral disturbances.
- The therapists should educate the parents as primary caregivers to enhance on several components of coping skills such as humor, planning, venting, self distraction etc to continue their battle of upbringing of their children as well as improve their own quality of life.

in family functioning, reduced social interaction outside the family etc. are the basic issues that the parents have to deal with. It often becomes very challenging for parents to provide long term care to the child, can become a strain and may result into impacts on psychological and physical health and overall Quality of Life of the parents²⁻⁸.

Parents of children with developmental disabilities experience higher level of stress to upbring their children in comparison to the parents with normally developed children⁹⁻¹². Parents can experience stress in several spheres of their life facing the challenges of caregiving of their children, regarding relational

¹MD, Assistant Professor, Department of Psychiatry, Tamralipto Government Medical College, Tamluk, West Bengal 721636

²MD, Associate Professor, Department of Psychiatry, Burdwan Medical College & Hospital, Burdwan, West Bengal 713104 and Corresponding Author

³MD, Former Professor of Psychiatry, Department of Psychiatry, Calcutta National Medical College & Hospital, Kolkata, West Bengal 700014

⁴MD, DCH, Professor, Department of Psychiatry, NRS Medical College & Hospital, Kolkata 700014

⁵MD, Professor, Department of Psychiatry, KPC Medical College & Hospital, Kolkata, West Bengal 700032

problem¹³, with financial constraints¹⁴, feeling of decreased parenting efficacy¹⁵ etc. The severity of stress perceived by the parents depends on the severity of the adversities in various sub-domains of their lives and their cognitive appraisal about the issues. Chronic unresolved stress can often compromise their psychological health¹⁶. By natural effort parents always try to cope with their stress by their own way actively or passively, that often depends on their psychological, social and cultural background and availability of resources. The term 'coping' refers to the behavioral strategies that an individual applies to reduce the effect(s) and or demand(s) of stress¹⁷. Folkman and Lazarus narrated coping efforts as the interaction of two separate cascades, ie, management of the person-environment relationship and regulation of associated stressful emotions. Lazarus (1980) defined coping as "the cognitive and behavioral efforts made to master, tolerate, or reduce external and internal demands appraised as taxing or exceeding the resources of the individual". These coping styles or mechanisms were described in the literature in different categories, subcategories and items¹⁸. In comparison to using maladaptive pattern of coping strategies such as negative emotion focused coping, using of healthy coping strategies such as problem focused coping, persons may manage their stress relatively in rational way and protect their psychological health better. There are studies available in the literature on the coping mechanism of parents with developmental disabilities but Indian studies are very scant¹⁹⁻²⁴. The pattern of using coping of Indian parents might be different from the western counterpart as there is significant difference in their socio-cultural-economic milieu. Even fathers and mothers may have contrast in pattern of using coping strategies as per their social role in that specific family and in that society or community at large.

This present study was undertaken in a Tertiary Care Hospital of Eastern India, to understand magnitude and pattern of behavioral disturbances and the extent of disability in the children with ID and the pattern of coping behaviors of their parents, observed differentially for fathers and mothers.

MATERIALS AND METHODS

In the current observational cross-sectional study, which was conducted at Psychiatry OPD of a tertiary care teaching hospital in Kolkata, West Bengal and consecutive 92 children (\leq 18 years) with ID as per ICD-10, DCR version¹ and their parents (<65 years, having children living with them) were included.

Parents with chronic medical illness or ID were excluded from the study. Research protocol was approved from the Institutional Ethics Committee. Intelligence Quotient (IQ) was tested using standard scales [eg, Vineland Social Maturity Scale (VSMS)²⁵, The Denver Developmental Screening Test (DST)²⁶ Seguin Form Board (SFB)²⁷ or Binet Kamat Test (BKT) for Intelligence²⁸ as applicable by an authorized clinical psychologist. Level of disability of the children was measured in percentage as per Gazette of India notification²⁹. Behavioral assessment of their child was done using Behavioral Assessment Scale for Indian Children with MR (BASIC-MR)³⁰. Coping mechanisms of the parents were assessed using Brief Cope Questionnaire (BCQ)³¹. BASIC-MR is a scale to assess the current behavior of the children with Intellectual Disability (ID). As Brief Cope Questionnaire (BCQ) is a self-rater's scale, the scale was validated in Bengali and used in the current study by following a standard procedure of validation. Assessment will be done by an expert such special educator, psychiatrist. Standard Package of Statistical Software Version 20 was used for statistical analysis of the collected data.

RESULTS

The study comprised of 92 children with ID and their parents (both father and mother).

(1) Description of Children :

The median age of children was 9.00 years, mean 11.34 years and Standard Deviation 5.07. Among the whole population 62% (n=57) of children were male and 38% (n=35) were female. Median, Mean, Standard Deviation of IQ and disability percentage of the children were 80.00, 79.87, 11.28 and 50, 38.83, 11.39 respectively. Among all children, 8 (8.7%) children had 50% or minor disability, 35 (38.04%) children had 75% or moderate disability and 49 (53.06%) children had 90% or severe disability (Table 1).

(2) Behavioral Disturbances of the Children :

Behavioral problem of the children in the current study were assessed by BASIC MR Part A and Part B. The distribution of scores across seven sub domains of BASIC MR Part A and Part B is described in Table 2.

Table 1 — Distribution of Age, IQ and Disability percentage of the Children (N=92)								
Variables	Variables Median Mean Standard Deviation							
Age (in years)	9.00	11.34	5.07					
IQ	80.00	79.87	11.28					
Disability 50 38.83 11.39								
	Table 2 — Distribution of the Scores of BASIC-MR of the Children (n=92)							
-----------------------------	---	----------	----------	----------	----------	----------	----------	--------
Part A (domain wise scores)					Part B			
	Domain 1	Domain 2	Domain 3	Domain 4	Domain 5	Domain 6	Domain 7	score)
Mean	105.79	113.13	118.48	58.69	50.15	58.05	41.99	14.60
Median	108.00	110.50	113.50	52.50	53.00	48.00	29.00	10.00
SD	46.02	51.09	95.87	40.41	32.48	42.15	42.13	12.14

(3) Description of Parents and their Families :

The median age of the fathers who participated in the study was 50.00 years, mean 51.00 years, Standard Deviation 8.85. The median age of mothers was 45.00 years, Mean 44.32 years, Standard Deviation 8.65. Among the families 71.7% (n=66) were nuclear and 28.3% (n=26) joint. Regarding socioeconomic status, 46.7% (n=43) belonged to upper middle class, followed by lower middle (n=35, 38%), upper lower (n=10, 10.9%) and upper (n=4, 4.3%) according to Modified Kuppuswamy's classification.

(4) Coping Mechanism among Parents :

The Brief Cope Scale doesn't yield any summary score. But for convenience of applying statistical tests the scores of 2 specific questions (which denote a specific type of coping mechanism, as per the scales) were added to calculate a score for each coping mechanism. Tables 3 show the distribution of these scores in 12 different areas of coping for both fathers and mothers.

DISCUSSION

A cross-sectional observational study was done in a Tertiary Care Medical College in Kolkata to know the extent of disability and behavioral disturbances of the children with ID and coping mechanism of their parents.

With ID, the children have deficit in several domains in adaptive functions and different types of

form of stereotypes, aggression, and violence that might lead to injury to self, to others. or destruction of property etc^{31,32}. These behaviors are detrimental not only to the person himself who exhibits such behavior but pose a challenge to their teachers and caregivers. A behavior becomes challenging when it is of such intensity, frequency or duration that it threatens the safety of self or others. It is likely to lead to restrictive or aversive responses and exclude social participation. Earlier studies from the west have found that ~10% of persons with ID manifest severe challenging behavior ranging from self-injury, destructiveness, aggression, and disruptive behavior³³. When milder forms of behavioral problems such as verbal aggression and temper

its severity³⁵. In the current study, BASIC MR scores was used to assess the level of behavioral problem in these children. Part-A deals with skills behavior and higher score indicates higher skills behavior. It has 7 domains with 40 items each. The scoring is done from 0 to 5 so a maximum score that can be obtained by a child is 1400. The average BASIC MR-A score of the sample of current study was 569 with maximum score of 1281 and minimum score of 46. So, it can be understood that most of the children had average skill set but definite conclusion cannot be reached as it was a heterogenous group. The BASIC MR, Part-B

tantrums are included, the prevalence goes up to 22.5

to 55%³⁴. A study from India reported that violent and

destructive temper tantrums and self-injurious

behavior among persons with ID vary according to

score is the indicator of behavioral issues or undesirable behavior. Total 75 tests are done and scored from 0 to 2. So, the maximum score that a child scored is 150. In the current study sample the average score was around 15 with maximum score was 50 and minimum score was 0. Which indicates a low level of behavioral problem in this particular group with a wide variation that's not always corroborated with IQ level. But this study still shows that a significant number of behavioral problems is found in children with MR.

In the current study most of the parents had applied positive modes of coping such

						/
Items of Brief Cope Scale	Fathers		Mothers			
	Mean	Median	SD	Mean	Median	SD
Self blame	3.14	2.00	1.43	3.11	3.00	1.48
Religion	3.97	4.00	1.72	3.62	3.00	1.65
Acceptance	6.89	7.00	1.57	6.43	7.00	1.78
Humor	2.23	2.00	0.79	2.13	2.00	0.54
Planning	5.82	6.00	1.65	5.57	6.00	1.83
Positive reframing	5.93	6.00	1.59	5.79	6.00	1.74
Venting	2.97	2.00	1.29	2.97	2.00	1.36
Instrumental support	4.87	5.00	1.85	4.68	5.00	1.88
Emotional support	4.63	4.00	1.72	4.27	4.00	1.76
Substance use	2.04	2.00	0.29	2.09	2.00	0.45
Behavioral disengagement	2.85	2.00	1.38	2.89	2.00	1.29
Active coping	6.13	6.00	1.77	6.04	6.00	1.93
Denial	2.92	2.00	1.82	2.80	2.00	1.64
Self distraction	4.03	4.00	1.76	3.82	4.00	1.56

Tables 3 — Distribution of scores of Brief Cope Scale of the parents (N=92)

37

b e h a v i o r a l disturbances in the

as acceptance, positive refraining and active coping to combat against their stress out of caring their disabled children. Education level of both fathers and mothers of the current sample were reasonably high and most of them belonged higher socio-economic classes in reference to national average level. Similar observation was reported from other Indian study²² where the association between higher education and use of healthy pattern of using coping strategies by the parents was also noted. Few other Indian research tried to study the gender difference among the parents in terms of perceived stress on account of the caregiving of children with developmental disabilities. In general, it was reported in the literature that the Indian fathers would perceive and cope to their stress more positive way than the mothers¹⁷. The difference in social role among the fathers and mothers in Indian society can explain this differential observation. However, in the current study no significant difference was observed among the parents in this regard. Current social reforms in the Bengali community and at large in the Indian society might be the possible explanation.

CONCLUSION

Parents with children with ID having significant disability and behavioral disturbances were using their coping strategies well to combat against their stress, out of caregiving their disabled children, being comparable for both fathers and mothers in Indian community.

Limitations :

It was a hospital-based study done in a metropolitan city on a relatively small study population. The current research was planned as a descriptive study. So, inferential statistics was not attempted which would have greater significance and larger clinical implications. Multicentric study with a large and variable sample is needed in future to explore the issue further.

REFERENCES

- World Health Organization. The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines. Geneva: World Health Organization; 1992. p. 377.
- Berill M, Brown I Quality of Life for people with intellectual disabilities. *Current Opinion Psychiatry* 2006; **19**: 508-13.
- 3 Boromand N, Narimani M, Mosazadeh T Comparing the psychological well being factors among the parents of the mentally retarded children with those of the normal children. *International Letters of Social and Humanistic Sciences* 2014; 21: 1-8.

- 4 Haimour AI, Abu-Hawwash RM Evaluating Quality of Life of parents having a child with disability. *International Interdisciplinary Journal of Education* 2012; **1(2):** 37-43.
- 5 Karasavvidis S, Avgerinou C, Lianou E, Priftis D, Lianou A, Siamaga E — Mental Retardation and Parenting Stress. *International Journal of Caring Sciences* 2011; 4: 21-31.
- 6 Malhotra S, Khan W, Bhatia MS Quality of Life of Parents having Children with Developmental Disabilities. *Delhi Psychiatry Journal* 2012; **15(1)**: 177-76.
- 7 Ravindranadan V, Raju S Emotional Intelligence and Quality of Life of Parents of Children with Special Needs. *Journal of the Indian Academy of Applied Psychology* 2008; 34: 34-9.
- 8 Summers JA, Poston DJ, Turnbull AP, Marquis J, Hoffman L, Mannan H, et al — Conceptualizing and measuring family quality of life. *Journal of Intellectual Diability Research* 2005; 49(10): 777-83.
- 9 Antonopoulou K, Manta N, Maridaki-Kassotaki K, Kouvava S, Stampoltzis A — Parenting and coping strategies among parents of children with and without autism: the role of anxiety and emotional expressiveness in the family. *Austin Journal of Autism & Related Disabilities* 2020; 6(1): 1054.
- 10 Bujnowska AM, Rodríguez C, García T, Areces D, Marsh NV — Parenting and future anxiety: The impact of having a child with developmental disabilities. *International Journal of Environmental Research and Public Health* 2019; **16(4):** 668.
- 11 Cuzzocrea F, Murdaca AM, Costa S, Filippello P, Larcan R Parental stress, coping strategies and social support in families of children with a disability. *Child Care in Practice* 2016; 22(1): 3-19.
- 12 Scherer N, Verhey I, Kuper H Depression and anxiety in parents of children with intellectual and developmental disabilities: A systematic review and meta-analysis. *PloS one* 2019; **14(7)**: e0219888.
- 13 Freedman BH, Kalb LG, Zablotsky B, Stuart EA Relationship status among parents of children with autism spectrum disorders: A population-based study. *Journal of Autism and Developmental Disorders* 2012; **42(4):** 539-48.
- 14 Parish SS, Rose RA, Swaine JG, Dababnah S, Mayr ET Financial well-being of single, working-age mothers of children with developmental disabilities. *American Journal on Intellectual and Developmental Disabilities* 2012; **117(5)**: 400-12.
- 15 Karst JS, Van Hecke AV Parent and family impact of autism spectrum disorders: A review and proposed model for intervention evaluation. *Clinical Child Family Psychological Review* 2012; **15(3)**: 247-77.
- 16 Trute B, Hiebert-Murphy D Family adjustment to childhood developmental disability: A measure of parent appraisal of family. *Journal of Pediatric Psychology* 2002; 27: 271-80.
- 17 Khan MF, Alam MA Coping trends of parents having children with developmental disabilities: A literature review. *European Journal of Special Education Research* 2016; **3(1):** 38-46.
- 18 Lazarus RS Stress, appraisal, and coping. Springer; 1984.
- 19 Vinayak S, Sekhon P Parenting of children with learning disabilities and their siblings. *Journal of the Indian Academy* of Applied Psychology 2012; **38(1):** 84-92.

- 20 Vidyasagar N, Koshy S Stress and coping in mothers of autistic children. *Journal of the Indian Academy of Applied Psychology* 2010; **36(2):** 245-48.
- 21 Chandramuki, Shastry IVK, Vranda MN Attitudes of parents towards children with Specific Learning Disabilities. *Disability, CBR and Inclusive Development* 2012; 23(1): 63-69.
- 22 Kumar VG Psychological stress and coping strategies of the parents of mentally challenged children. *Journal of the Indian Academy of Applied Psychology* 2008; **34(2):** 227-31.
- 23 Hussain A, Juyal I Stress appraisal and coping strategies among parents of physically challenged children. *Journal of the Indian Academy of Applied Psychology* 2007; **33(2):** 179-82.
- 24 Kalgotra K, Warwal JS Impact of disability and coping strategies among the parents of children with intellectual disabilities. *Indian Journal of Applied Research* 2014; 4(6): 1-4.
- 25 Doll EA Vineland Social Maturity Scale: Manual of Directions, Revised Edition. Circle Pines; MN: American Guidance Service; 1965.
- 26 Frankenburg William K, Dobbs JB The Denver Developmental Screening Test. *The Journal of Pediatrics* 1967; **71(2)**: 181-91
- 27 Goel SK, Bhargava M Handbook of Seguin Form Board (SFB); Agra, National Psychological Corporation; 1990.
- 28 Kamat W Binet Kamat Test of Intelligence. Measuring Intelligence of Indian Child, Bombay Oxford University Press, 1967.

- 29 Government of India. Guidelines for Evaluation of Various Disabilities and Procedure for Certiûcation. Ministry of Social Justice and Empowerment Notiûcation. New Delhi: 1st June, 2001. Available from: http://www. socialjustice.nic.in. [Last accessed on 204th June,2023].
- 30 Peshawaria R, Venkatesan S Behavioral assessment scales for Indian children with mental retardation: BASIC-MR. 1992.
- 31 Lowe K, Allen D, Jones E, Brophy S, Moore K, James W Challenging behaviors: prevalence and topographies. *Journal of Intellectual Disability Research* 2007; **51**: 625-36.
- 32 Smith KR, Matson JL Behavior problems: differences among intellectually disabled adults with co-morbid autism spectrum disorders and epilepsy. *Research of Developmental Disability* 2010; **31(5):** 1062-69.
- 33 Emerson E, Kiernan C, Alborz A The prevalence of challenging behaviors: a total population study. *Research on Developmental Disability* 2001; 22(01): 77-93.
- 34 Cooper SA, Smiley E, Morrison J, Williamson A, Allan L Mental ill-health in adults with intellectual disabilities: prevalence and associated factors. *British Journal of Psychiatry* 2007; **190**: 27-35.
- 35 Lakhan R, Kishore MT Behavior problems in children with intellectual disabilities in a resource-poor setting in India - part 1: association with age, sex, severity of intellectual disabilities and IQ. *Journal of Applied Research of Intellectual Disability* 2018; **31(01):** 43-50.



DISCLAIMER



Journal of the Indian Medical Association (JIMA)

The Journal of the Indian Medical Association (JIMA) (ISSN 0019-5847) is published monthly in English language from Editorial Offices at Sir Nil Ratan Sircar IMA House, 53, Sir Nilratan Sarkar Sarani, Kolkata-700014. Telephone No.:+91-33-22378092, (+919477493027); websites: <u>https://onlinejima.com</u> & <u>www.ejima.in</u>; Emails: jima1930@rediffmail.com; jimaeditorial@gmail.com. The Journal of the Indian Medical Association (JIMA) is a publication of Indian Medical Association (IMA). Material printed in JIMA is copyrighted by the Journal of the Indian Medical Association (JIMA). All rights reserved. No part of this reprint may be reproduced, displayed, or transmitted in any form or by any means without prior written permission from the Editorial Board. Please

contact the Permissions Department via email at jimaeditorial@gmail.com. For reprints please email: jimamkt@gmail.com. JIMA does not hold itself responsible for statements made by any contributor. Statements or opinions expressed in JIMA reflect the views of the author(s) and not the official policy of the Indian Medical Association unless so stated. JIMA reprints are not intended as the sole source of clinical information on this topic. Readers are advised to search the JIMA Web site at https://onlinejima.com and other medical sources for relevant clinical information on this topic. Reprints of articles published in JIMA are distributed only as freestanding educational material. They are not intended to endorse or promote any organization or its products or services.

— Hony Editor

Original Article

Neutrophil to Lymphocyte Ratio (NLR) as an Inflammatory Marker in Predicting Severity in Acute Ischemic Stroke Patients

Deepti Sharma¹, Gordhan Lal Nagar², Narendra Fageria³, Anup Kumar Mangal⁴, Pawan Kumar⁴

Background : Inflammation plays an important role in Acute Ischemic Stroke (AIS). Chemokines and cytokines releases from ischemic tissue and recruit peripheral circulating leukocytes mainly neutrophils into the brain which precipitate ischemic brain injury and increases stroke severity. Lymphocytes on the other hand represent the protective or regulatory component of inflammation which decreases due to increased release of cortisol in response to stress.

Aims and Objectives : To find the role of Neutrophil to Lymphocyte Ratio (NLR) in predicting the severity in patients of AIS.

Materials and Methods : This study included 100 patients of AIS admitted in Government Medical College and Hospitals, Kota with all relevant exclusion criteria. Peripheral blood sample was taken before initiating any treatment and CBC test was performed from which NLR was calculated. The ratio was then compared with healthy control group and also among patients of AIS .

Results : Mean value of NLR in AIS patients was 3.44 which was higher than NLR of control group which was 1.89 (p=0.001).

Conclusion : Thus, NLR is a cost effective, easy to calculate and newer inflammatory marker that can help in predicting severity of acute ischemic stroke score.

[J Indian Med Assoc 2024; 122(12): 40-3]

Key words : Acute Ischemic Stroke (AIS), Neutrophil to Lymphocyte Ratio (NLR).

ippocrates, the father of medicine, first recognized stroke over 2,400 years ago. At this time stroke was called apoplexy^{1,2}. Stroke is the most common neurological disorder Worldwide and it is the most frequent of all the neurological disorders. Stroke is also known as Cerebrovascular Accident (CVA) derived from Greek word in the year 1599 which means 'StruckDown'³. It is the disease of developed nations.

According to the World Health Organization, 15 million people suffer from stroke worldwide every year. Of these, 5 million die and another 5 million are permanently disabled⁴.

Neutrophil to Lymphocyte Ratio (NLR) is an effortless and basic parameter that is readily obtained from the complete blood count, even in peripheral hospitals. Inflammation is regarded as a set of interactions between and among immune related cells such as lymphocytes, neutrophils which in turn lead to killing of tissues and destruction which is going on in stroke. One of the inflammatory markers in stroke

¹MBBS, MD (General Medicine), Senior Professor and Head
²MBBS, Junior Resident and Corresponding Author
³MBBS, MD (General Medicine), Senior Resident
⁴MBBS, MD (General Medicine), Senior Resident

Received on : 19/01/2023

Accepted on : 08/12/2023

Editor's Comment :

The Neutrophil to Lymphocyte Ratio (NLR) is a simple, accessible, and cost-effective inflammatory marker. An elevated NLR on admission is associated with greater stroke severity, worse outcomes and increased risk of complications like infarct expansion and mortality. NLR can be used as an adjunct to traditional markers and imaging to help guide early risk stratification and management in acute ischemic stroke patients.

is NLR and also has its relationship between many diseases like, myocardial infarction, COPD, chronic renal failure etc⁵⁻⁷.

Inflammatory cascade following Acute Ischemic Stroke (AIS) is an important pathological process in damaged brain tissue. The inflammatory process in acute ischemic stroke involves cytokines and chemokines released from ischemic tissues, promoting the accumulation of leukocytes to the ischemic areas. Among circulating leukocytes, neutrophils have been regarded as crucial mediators of ischemic injury. Accumulated neutrophils release free oxygen radicals, various inflammatory cytokines, and neurotoxic substances, all of which cause cellular necrosis and apoptosis in ischemic tissues. Lymphocytes, a type of leukocyte, are also known to be involved in inflammatory responses to Acute Ischemic Stroke (AIS). Low lymphocyte counts increase sympathetic activity and baseline cortisol

Department of Medicine, Goverment Medical College, Kota, Rajasthan 324010

levels, which can cause an increase in the production of inflammatory cytokines that aggravate ischemic injury. Therefore, increased total White Blood Cell (WBC) and Leukocyte Counts and reduced lymphocyte counts are associated with poor outcomes in various cardiovascular diseases⁸.

Earlier researchers have studied Neutrophil to Lymphocyte Ratio (NLR) and Mean platelet Volume (MPV) as prognostic marker in acute ischemic stroke separately. But we would like to study the combined effect of NLR and MPV in patients of acute ischemic stroke as prognostic marker.

AIMS AND OBJECTIVES

To study Neutrophil to Lymphocyte Ratio (NLR) in patients of Acute Ischemic Stroke.

MATERIALS AND METHOD

The study was performed at the Government Medical College and attached groups of Hospitals Kota in year 2021-2022, after obtaining the approval of the Internal Ethical Committee of Government Medical College & Hospital, Kota and Informed written consent from all the patients considered for the study was procured before conducting the study.

Study Design : Hospital based prospective and observational study carried out on patients admitted and diagnosed as an Acute Ischemic Stroke.

Study Place : Department of medicine, Government Medical College & Associated Group of MBS Hospital, Kota (Rajasthan).

Study Population : patients with Acute Ischemic Stroke who had symptom onset within 7 days.

Study Duration : 2021-2022.

Inclusion Criteria :

(1) Patients of Age >18 years admitted in MBS Hospital, Kota diagnosed with Acute Ischemic Stroke

Exclusion Criteria:

(1) Patients of Age <18 years.

(2) Patients with other forms of stroke like ICH/ SAH /post traumatic.

(3) Patients with venous sinus thrombosis.

(4) Serious ill/moribund patients.

(5) Patients having hepatic or renal disease.

(6) Systemic acute or chronic inflammatory or autoimmune or infectious disease, connective tissue diseases or prior myocardial infarction.

(7) Individuals with missing data of lipid profile, monocyte count.

(8) Pregnant patients.

(9) Patient refusing to give informed consent for the study.

RESULT

Out of 100 subjects in the control group; 55 were males and 45 were females whereas in the study group of 100 subjects,57 were males and 43 were females. Males were more than females with male to female ratio being 1.3:1. Thus, males are more prone to develop stroke as compared to female. The maximum number of patients were in the range of 50-59 years ie, 38 cases (38%) followed by in 60-69 years 35 cases (35%), in 70-79 years 15 cases (15%), in 40-49 years 8 cases (8%) and in \geq 80 years of age group 4 cases (4%) so in 50-69 years cases 73%.

When the study group was compared on the basis of focal neurological deficit, maximum number of patients had either isolated left hemiparesis or isolated right hemiparesis amounting to 32% each. 17% of patients had right hemiparesis with cranial nerve palsy and 13% had left hemiparesis with cranial nerve palsy. Patients with the features of posterior circulatory stroke like vertigo, blurring of vision etc contributed 6% of the cases. ACS is 94% in compare to PCS is 6%.

Table 1 shows the comparison of control and study group on the basis of mean Neutrophil count at the time of admission. Mean Neutrophil count \pm SD of the cases in study group was $4.472 \pm 1.1(10^3/\text{cu mm})$ which was higher than the mean value of $3.4965\pm 1.7 (10^3/\text{cu mm})$ obtained in the control group and difference between the two was statistically significant with the p-value of 0.0001.

Table 2 shows the comparison of control and study group on the basis of mean total lymphocyte count. Mean Total lymphocyte count \pm SD of the cases in the study group was 1.30 ± 0.54 (10^{3} /cu mm) which was lower than the mean value of 1.85 ± 0.22 (10^{3} /cu.mm) obtained in the subjects of control group and difference between the two values was statistically significant with the p-value of 0.0001.

Table 3 shows the comparison of control and study group on the basis of Neutrophil to Lymphocyte Ratio. Mean NLR \pm SD of all the cases in study group was 3.44 \pm 2.56 which was higher than the mean value of 1.89 \pm 0.61 obtained in the subjects of the control group

Table 1 — Comparison of Total Neutrophil Count in Control and Study Group						
Neutrophil	Control group Acute Ischemic					
(10 ³ /cu mm)	(n=100) Stroke Group (n=100)					
Mean	3.4965	4.472				
SD	1.7	1.1				
Т	6.473					
P-value		0.0001				

Table 2 — Comparison of Total Lymphocyte Count in Control and Study Group					
Total LymphocyteControl groupStudy groupcount (10³/cu mm)(n=100)(n=100)					
Mean	1.85	1.30			
SD	0.22 0.54				
Т	9.432				
P-value	0.000	01			

Table 3 — Comparison of Neutrophil to Lymphocyte Ratio (NLR) Between Acute Ischemic Stroke Subjects and Control Subjects						
Neutrophil-to-	Control Subjects Acute Ischemic					
lymphocyte Ratio	(n=100) Stroke Subjects (n=100)					
Mean	1.89	3.44				
SD	0.61	2.56				
Т	2.32					
P-value	0.021					

and the difference between the two values was statistically significant with the p-value of 0.021.

Table 4 depicts the comparison of Acute Ischemic Stroke group and control group on the basis of demographic and biochemical profile including risk factors. The Mean \pm SD of all subjects in control and Acute Ischemic Stroke study group was obtained including age (61.33 \pm 10.965; 60.79 \pm 13.86) years,

Table 4 — Comparison of Demographic and BiochemicalProfile Between Acute Ischemic Stroke Group and ControlGroup				
Neutrophil-to-	Control Group	Acute Ischemic		
lymphocyte Ratio	(n=100)	Stroke Group		
		(n=100)		
Male (%)	55	57		
Age (year)	61.33±10.965	60.79±13.86		
Hypertension, n(%)	44	50		
Smoker, n(%)	30	40		
Alcoholic , n(%)	26	35		
Diabetes mellitus, n(%)	25	36		
HB (gm/dl), n(%)	11.63±1.22	12.57±1.69		
B Glucose (mg/dL) (R)	148.97±87.75	196.33±92.26		
B Urea (mg/dL)	30.87±6.51	39.57±16.72		
S Creatinine (mg/dL)	0.984±0.162	1.19±0.40		
Total cholestrol (mg/dL)	190.2±31.29	198.48±62.71		
S Triglyceride (mg/dL) 127.66±35.64 145.96±55.04				
Neutrophil count (10 ³ /cu.mm)	3.4965±1.7	4.472±1.1		
Lymphocyte count (10 ³ /cu.mm)	1.85±0.22	1.30±0.54		
Neutrophil to Lymphocyte Ratio	1.89±0.69	3.44±2.56		

Table 5 — Distribution of Neutrophil Count, Lymphocyte Count and Neutrophil to Lymphocyte Ratio in Both Control and Study Group at the Time of Admission					
	Control Group (Mean±SD)	Study group at the Time of Admission (Mean±SD)	t value	p value	
Neutrophil Count (10³/cu.mm) Lymphocyte count (10³/cu.mm) Neutrophil to Lymphocyte Ratio	3.4965±1.7 1.85±0.22 1.89±0.61	4.472±1.1 1.30±0.54 3.44±2.56	6.473 9.432 2.32	0.0001 0.0001 0.021	

Hb (11.63 \pm 1.22; 12.57 \pm 1.69) gm/dl, B Glucose (148.97 \pm 87.75; 196.33 \pm 92.26) mg/dl, B Urea (30.87 \pm 6.51; 39.57 \pm 16.72) mg/dl, S Creatinine (0.984 \pm 0.162; 1.19 \pm 0.40) mg/dl, Total cholesterol (190.2 \pm 31.29; 198.48 \pm 62.71) mg/dl, Triglyceride (127.66 \pm 35.64; 145.96 \pm 55.04) mg/dl, Neutrophil count (3.4965 \pm 1.7; 4.472 \pm 1.1) 10³/cu mm, Lymphocyte count (1.85 \pm 0.22;1.30 \pm 0.54) 10³/cu mm and Neutrophil to Lymphocyte Ratio (1.89 \pm 0.61; 3.44 \pm 2.56) respectively.

Table 5 clearly highlighted that the mean Neutrophil count, mean lymphocyte count and mean Neutrophil to mean Lymphocyte Ratio in study group was $4.472\pm1.1\times10^{3}/cu$ mm, $1.30\pm0.54\times10^{3}/cu$ mm and 3.44 ± 2.56 respectively which was higher than the mean control values of $3.4965\pm1.7 \times 10^{3}/cu$ mm, $1.85\pm0.22\times10^{3}/cu$ mm and 1.89 ± 0.61 respectively and the difference was statistically significant in both control and Acute Ischemic Stroke Study group with the p-value of 0.021.

DISCUSSION

Atherosclerosis, especially intracranial atherosclerosis, is an intrinsic abnormality in the development of ischemic stroke.Inflammation and lipid abnormalities has been proposed as the main constituents of the pathophysiology of atherosclerosis development and progression.

In this study we evaluated mean Neutrophil to Lymphocyte Ratio (NLR) and Mean Platelet Volume (MPV) in patients of Acute Ischemic Stroke by comparing with age and sex matched control subjects. An attempt was made to find out the association between NLR and smoking ,alcohol consumption , hypertension, diabetes mellitus. The Neutrophil to Lymphocyte Ratio (NLR) is a parameter of inflammation that is easy to obtain and has been proposed as an independent useful prognostic marker to predict the mortality and prognosis of some cardiovascular and neurologic diseases. Many studies have demonstrated that peripheral leucocyte levels increase following cerebrovascular ischemia and postulated that the initial peripheral leucocyte count following a stroke can help to predict stroke severity.

> It has been supposed that the leucocyte count at the time of admission is predictive of the likelihood of AIS and the impact of any resulting neurologic disability on daily living activity.

> 100 patients of Acute Ischemic Stroke and 100 age and sex matched

control subjects were included in our study. Although the control subjects were free from Acute Ischemic Stroke, some of them were having risk factors for ischemic stroke and it can be expected that they represented a stage in the initiation and progression of ischemic stroke.

NLR and Acute Ischemic Stroke :

In our study, mean Neutrophil to Lymphocyte Ratio (NLR), which is obtained by dividing absolute Neutrophil count by Lymphocyte count, was 3.44±2.56 in Acute Ischemic Stroke patients, which was significantly higher than the mean NLR value of 1.89±0.61 obtained in control subjects. Another study by Celikbilek, *et al* 2013 showed that high NLR levels are associated with increased infarct volume and mortality.

The Neutrophil to Lymphocyte Ratio (NLR) is a parameter of inflammation⁴ that peripheral leucocyte levels increase following cerebrovascular ischemia, and postulated that the initial peripheral leucocyte count following a stroke can help predict stroke severity. It has been supposed that the leucocyte count at the time of admission is predictive of the likelihood of AIS and the impact of any resulting neurologic disability on daily living activity.

NLR and Hypertension :

Hypertension is the major risk factor for Acute Ischemic Stroke in our study. We have tried to study the correlation between NLR and hypertension in both controls and cases. The NLR values of our cases were 3.44±2.56 in hypertensive patients and were 3.25±1.95 in non hypertensive patients. The NLR ratio in hypertensive controls and non-hypertensive controls were 1.89±0.61 and 1.81±0.52 respectively. This showed that there was a slight rise in values of NLR in hypertensive population but this was statistically non significant in our study.

A study conducted by Celik, *et al*^{θ}, found that the NLR values were significantly higher in the hypertensive group than in the control group. A similar finding was obtained in the study by Aydin, *et al*¹⁰. In their study found that NLR value was higher in hypertensive patients compared to controls and also added to their conclusion that asymptomatic organ damage in primary hypertension was higher in patients with high NLR values.

The statistically insignificant NLR value in hypertensive patients in our study may be due to the

fact that there may be some other risk factors of atherosclerosis like diabetes, smoking etc in the non hypertensive group too, which can cause a rise in their NLR value there by not making a statistically significant difference in NLR values between hypertensive and non-hypertensive group. Also in this study we have not considered the duration and control of hypertension which may also have interfered with the NLR value of the patients. Our study was conducted in a small group of 100 stroke patients, this low number of cases may have further influenced the observation.

CONCLUSION

NLR is a simple, cost effective and easily obtainable novel marker that may help in predicting the severity of disease as evidenced by its increased value in patients of acute ischemic stroke.

REFERENCES

- Jeff K A Stroke in Time. *MERGINET.com*. September 2001 6(9). Archived from the original on 2012-12-08.
- 2 The history of stroke and cerebrovascular disease, MaurizioPaciaroni, Julien Bogousslavsky, Stroke Unit and Division of Cardiovascular Medicine, University of Perugia, Santa Maria dellaMisericordia Hospital, Perug, Italy Department of Neurology and Neurorehabilitation, Genolier Swiss Medical Network, 1823, Glion/Montreux, Switzerland.
- 3 Thompson JE The evolution of surgery for the treatment and prevention of stroke. The Willis Lecture. *Stroke* 1996; 27(8): 1427-34.
- 4 Jeff K A Stroke in Time (September 2001). MERGINET.com
- 5 Templeton AJ, M cNamara MG, Šeruga B Prognostic role of neutrophil-to-lymph ocyte ratio in solid tumors: a systematic review and meta-analysis. J Natl Cancer Inst 2014; 106(6): dju124.
- 6 Yoon NB, Son C, U m SJ Role of the neutrophillymphocyte count ratio in the differential diagnosis between pulmonary tuberculosis and bacterial communit y-acquired pneumonia. *Ann Lab Med* 2013; **33(2):** 105-10.
- 7 Rodak BF, Fritsma GA, Keohane E Hematology: Clinical Principles and Applications. 4th edition.
- 8 National Institute of Health, National Institute of Neurological Disorders and Stroke. Stroke Scale. https://www.ninds.nih.gov /sites/default/files /NIH _ Stroke_ Scale_Booklet.pdf
- 9 Balta S, Celik T, Ozturk C, Kaya MG, Aparci M, Yildirim AO, et al — The relation between monocyte to HDL ratio and noreflow phenomenon in the patients with acute ST-segment elevation myocardial infarction. The American Journal of Emergency Medicine 2016; 34(8): 1542-7.
- 10 Cetin MS, Cetin EH, Kalender E, Aydin S, Topaloglu S, Kisacik HL, et al Monocyte to HDL cholesterol ratio predicts coronary artery disease severity and future major cardiovascular adverse events in acute coronary syndrome. Heart, Lung and Circulation 2016; 25(11): 1077-86.

Original Article

Microvascular Density of Invasive Breast Carcinoma as an Independent Prognostic Factor in Comparison with Already Established Ones : A Study from Northeast India

Junu Devi¹, Asreen Suhana²

Background and Objectives : Breast Carcinoma is the most common cancer in women Worldwide. This study was done to assess the tumour angiogenesis using CD34 immuno-reactivity and thereby see the relationship of the same with already established prognostic factors like age, histological type, tumour size, tumour grade (Modified Bloom Richardson Scoring System) and lymph nodal metastasis.

Materials and Methods : The 19 Breast Carcinoma cases included in this study were diagnosed in the Department of Pathology, Gauhati Medical College and Hospital within 1 year and Tumour Angiogenesis was assessed using CD34 immunoreactivity. The results were then compared with the already established prognostic factors.

Results : The relationship of MVD/HPF with tumour size, MBR tumour grade and lymph nodal metastases was found to be statistically significant at p-value <0.05. MVD/HPF with patient age was not statistically significant. Grade I &II IDC and medullary carcinoma had low MVD/HPF, whereas grade III IDC, metaplastic carcinoma and invasive papillary carcinoma had high MVD/HPF.

Conclusion : Tumour angiogenesis assessed by MVD/HPF following CD34 immunostaining can be used as a prognostic factor in Breast Carcinoma alongwith other known factors with more studies and standardized methods to calculate the MVD/HPF and can also be a potential targeted therapy with anti-angiogenic factors.

[J Indian Med Assoc 2024; 122(12): 44-6]

Key words : Tumour Angiogenesis, Breast Carcinoma, Prognostic Factor.

reast Carcinoma is the most common cancer in women worldwide, second only to lung cancer as a cause of cancer related deaths¹ with increasing prevalence in India². It is seen in all age groups with peak incidence in 45-60 years age group³. For every two women affected by breast cancer in India, one is dying⁴. Breast cancers staged on the basis of tumour size, lymph nodal metastases and distant metastases seem to show different etiology and clinical behaviour even among those presenting in the same stage⁵. A number of prognostic factors, namely histologic type, tumour size, lymph node metastases, distant metastases, locally advanced disease, molecular subtype, ER/PR receptors, HER2 expression and proliferation rate (Ki-76 index) have been extensively studied and established. Recent studies suggest tumour angiogenesis (assessed by microvessel density) as an independent predictive and prognostic factor in breast carcinoma⁶.

Received on : 10/08/2023

Accepted on : 14/05/2024

Editor's Comment :

Microvascular Density (MVD) is an independent prognostic factor in invasive breast carcinoma and it has strong relationship with other prognostic factors. This new prognostic marker can be helpful for targeted therapy using antiangiogenic factor in future.

CD34 is an endothelial marker, sensitive to tumour angiogenesis representing the clear state of neovascularization during tumour growth⁷. Previous studies have suggested the importance of CD34 in cancer prognosis⁸. Some studies have, however, denied this association⁹. So, this study was done to assess tumour angiogenesis using CD34 immunostaining to see the relationship of tumour angiogenesis with other prognostic parameters.

MATERIALS AND METHODS

This hospital-based cross-sectional study was conducted in the surgical specimens obtained in the Department of Pathology, Gauhati Medical College and Hospital, Guwahati in a one-year period.

Inclusion Criteria : All invasive Breast Carcinoma, irrespective of age and sex.

Exclusion Criteria: All in-situ Breast Carcinoma. The specimens were subjected to macroscopic examination, followed by histopathological

Department of Pathology, Gauhati Medical College and Hospital, Guwahati, Assam 781032

¹MD (Pathology), Ph D, Associate Professor

 $^{^2\}mbox{MD}$ (Pathology), Postgraduate Trainee and Corresponding Author

examination and immuno-histochemistry with CD34. Microvessel Density (MVD) was calculated in the tumour sections by the technique developed by Weidner, *et.al* and mean MVD per high power field (MVD/HPF) was calculated which was found to be 11.7. Any value lower than the mean was regarded as "low MVD" and any value above the mean was regarded as "high MVD". The p-value was calculated using Chi-square method and p-value <0.05 was considered as statistically significant.

RESULTS

Most common age group of the patients studied was 50-59 years (42.1%), followed by 40-49 years age group (21%). IDC was the most common histological type (78.9%). Most tumours were of MBR grade III (57.9%) with tumour size >5cm (T3) (57.9%). Lymph nodal metastases was seen in majority of the cases (73.6%).

All the 19 cases were divided into high and low based on microvessel density/ HPF with respect to tumour angiogenesis based on CD34 immunoreactivity. The relationship between tumour angiogenesis with patient age, histological type, MBR score, tumour size and lymph nodal metastases was analyzed.

Most of the cases were IDC (15 out of 19). IDC and Medullary carcinoma mostly showed low MVD/ HPF, whereas the high grade ones like metaplastic carcinoma, sarcoma and invasive papillary carcinoma had high MVD/HPF (Tables 1-5)(Figs 1-4).

DISCUSSION

In this study, statistically significant relation (p<0.05) was noted between MVD/HPF and MBR tumour grade, tumour size and lymph nodal metastases. Similar findings were noted in studies done by Pyakurel, *et* al^{10} , Kwatra, *et* al^{11} , Bhawna Agnani, *et* al^{12} and Horak, *et* al^{13} .

No statistical significance was noted between patient age and MVD/HPF in this study, which could be compared to studies done by Ebru, *et al*¹⁴ and Wang G, *et al*¹⁵.

So, it has been observed that tumour angiogenesis assessed by MVD/HPF following CD34 immunostaining has a strong relationship with tumour size, MBR tumour grade and lymph nodal metastases. Thus,

Table 1 — Microvessel Density (MVD) / HPF versus Patient Age							
	Age Groups						
MVD / HPF	20-29	30-39	40-49	50-59	60-69	70-79	80-89
Low (<11.7)	0	2	3	6	0	0	0
High (>11.7)	1	1	1	2	1	1	1
Total	1	3	4	8	1	1	1
The p-value >0.05							

Table 2 — MVD / HPF versus Histological type						
	Histological Type					
MVD / HPF	IDC	Medullary	Metaplastic	Sarcoma	Invasive papillary	
Low (<11.7)	10	1	0	0	0	
High (>11.7)	5	0	1	1	1	
Total	15	1	1	1	1	

Table 3 — MVD / HPF versus MBR Tumour Grade					
	MBR Grade				
MVD / HPF	Grade I	Grade II	Grade III		
Low (<11.7)	2	6	3		
High (>11.7)	0	0	8		
Total 2 6 11					
The p-value <0.05.					

Table 4 — MVD / HPF versus Tumour Size						
Tumour size						
MVD / HPF	T1 (<2 cm)	T2 (2-5 cm)	T3 (> 5 cm)			
Low (<11.7)	1	4	1			
High (>11.7)	0	3	10			
Total	1	7	11			
p=0.02.						

Table 5 — MVD / HPF versus Lymph Nodal Metastases					
	Lymph nodal metastases				
MVD/HPF	Present	Absent			
Low (<11.7)	3	4			
High (>11.7)	11	1			
Total	14	5			
p=0.01.					



Figs 1 & 2 — Breast Carcinoma with High Microvessel Density after CD34 immuno-staining, IHC Staining CD34, 10X10

tumour angiogenesis can be used as a specific prognostic marker in breast carcinoma.

CONCLUSION

Breast Carcinoma continues to carry its havoc inspite of the tragic fact that it occurs in an exposed organ accessible to self-examination and continuous high clinical surveillance. With larger number of samples, newer prognostic parameters like tumour angiogenesis and more detailed studies on



Figs 3 & 4 — Breast Carcinoma with Low Microvessel Density after CD34 Immuno-staining, IHC Staining CD34, 10X10

targeted therapies with anti-angiogenic factors, we will hopefully be able to treat and manage cases of Breast Carcinoma in better ways in the upcoming future.

REFERENCES

- 1 Kumar V, Abbas AK, Aster JC, Perkins JA Robbins & Cotran Pathologic Basis of Disease. 2014.
- 2 Agarwal S, Singh A, Bagga PK Immunohistochemical evaluation of lymphovascular invasion in carcinoma breast with CD34 and D2-40 and its correlation with other prognostic markers. *Indian Journal of Pathology and Microbiology* 2018; 61: 39-44.
- 3 Renshaw AA Diagnostic Histopathology of Tumors, 3rd Edition. Advances in Anatomic Pathology 2008; 15: 183.
- 4 Cancerindia.org.in/breast-cancer/
- 5 Foulkes WD, Grainge MJ, Rakha EA, Green AR, Ellis IO Tumor size is an unreliable predictor of prognosis in basallike breast cancers and does not correlate closely with lymph node status. *Breast Cancer Res Treat* 2008; **117(1)**: 199-204. Breast cancer research and treatment. 117. 199-204. 10.1007/ s10549-008-0102-6.
- 6 Goulding, H, Rashid A, Robertson NF, Bell JF, Elston JA, Blamey CW, et al — Assessment of angiogenesis in breast carcinoma: an important factor in prognosis? *Human Pathol*ogy 1995; **26 (11):** 1196-200.
- 7 Cao Y, Zhang ZL, Zhou M, Elson P, Rini B, Aydin H, et al Pericyte coverage of differentiated vessels inside tumor vasculature is an independent unfavorable prognostic factor for patients with clear cell renal cell carcinoma. *Cancer* 2013; **119(2):** 313-24. doi: 10.1002/cncr.27746. Epub 2012 Jul 18. PMID: 22811049.
- 8 Zorgetto VA, Silveira GG, Oliveira-Costa JP, Soave DF, Soares FA, Ribeiro-Silva A The relationship between lymphatic vascular density and vascular endothelial growth factor A (VEGF-A) expression with clinical-pathological features and survival in pancreatic adenocarcinomas. *Diagn Pathol* 2013; **8**: 170. doi: 10.1186/1746-1596-8-1.

- 9 Dhakal HP, Nesland JM, Førsund M, Trope CG, Holm R Primary tumor vascularity, HIF-1á and VEGF expression in vulvar squamous cell carcinomas: their relationships with clinicopathological characteristics and prognostic impact. *BMC Cancer* 2013; **13**: 506. doi: 10.1186/1471-2407-13-506. PMID: 24165149; PMCID: PMC3871003.70. PMID: 24138811; PMCID: PMC3816792.
- 10 Pyakurel D, Karki S, Cs A A study on microvascular density in breast carcinoma. *Journal of Pathology of Nepal* 2014; 4: 570-5.
- 11 Kwatra A, Aggarwal D, Gupta R, Chaturvedi A, Kudesia M, Singh S — Correlation of various histopathologic prognostic factors with Nottingham prognostic index and microvessel density in invasive breast carcinoma: A study of 100 cases. *Indian Journal of Cancer* 2015; **52(1):** 110-3.
- 12 Agnani B, Solanki R, Ansari M, Agnani S Prognostic Significance of Microvessel Density as Assessed by anti CD34 Monoclonal Antibody in Invasive Ductal Carcinoma of Breast. 2020.
- 13 Horak E, Klenk N, Leek RD, Lejeune S, Smith K, Stuart N, et al — Angiogenesis, assessed by platelet/endothelial cell adhesion molecule antibodies, as indicator of node metastases and survival in breast cancer. The Lancet 1992; 340: 1120-4.
- 14 Sener E, Sipal S, Gündoçdu C Comparison of Microvessel Density with Prognostic Factors in Invasive Ductal Carcinomas of the Breast. *Turkish Journal of Pathology* 2016; 32: 164-70.
- 15 Wang G Microvessel density recognized by Endoglin as prognostic markers in breast carcinoma. *Journal of Cancer Science & Therapy*, 2015.

Original Article

Comparative Analysis of Sensitivity and Specificity of Creatinine Kinase MB and Cardiac Troponin I for the Diagnosis of Acute Myocardial Infarction

Nidhi Rohan Purandare¹, Astha Goyal², Sarita Shinde³, Shilpa Jain⁴

Background : Early diagnosis is of crucial importance for the successful management of individuals with Acute Myocardial Infarction and is one of the leading causes of mortality and morbidity. We aimed to determine the sensitivity and specificity of two markers commonly used in our institution to diagnose patients with complaints indicative of Acute Myocardial Infarction.

Material and Methods : Patients admitted to the tertiary care center with complaints of chest pain were screened. Creatine Kinase Myoglobin Binding (CKMB) and Cardiac Troponin I (cTnI) were performed by following standard protocols. Depending upon the onset of chest pain subjects were divided into two groups, Group 1 : Within 24 hours of chest pain, Group 2 : Within 24-72 hours of chest pain. They were subdivided based on a diagnosis of MI. Results were used to calculate the specificity and sensitivity of CKMB, cTnI, and CKMB and cTnI together.

Results: Total of 368 patients were studied. Majority of them were >60 years of age. Of these, 133(36%) had confirmed diagnosis of MI. CKMB is more sensitive (82.19%) within 24 hours of MI but after 24 hours sensitivity decreases to (57.53%). CKMB is specific after 24 hours of MI (92.26%) compared to within 24 hours of MI (65.81%) whereas, cTnI is more specific (95.00%) after 24 hours.

Conclusion : For the diagnosis of Acute Myocardial Infarction (AMI), CKMB is more sensitive and cTnI is more specific, therefore it will be better if we perform cTnI rapid card test in combination with CKMB.

[J Indian Med Assoc 2024; 122(12): 47-51]

Key words : CKMB, cTnl, Acute Myocardial Infarction, Sensitivity, Specificity.

Myocardial infarction, the most serious side effect of Coronary Artery Disease, results from an abrupt thrombotic process that alters the perfusion balance between supply and demand within the coronary arteries^{1,2}. Acute Myocardial Infarction (AMI) has a significant chance of being diagnosed early, which could save or extend a patient's life³. The American Heart Association advises that choosing the best treatment plans in the Emergency Department (ED) requires accurate 30-minute preferred turn around time for the identification of cardiac biomarkers play an important role in the detection of Acute MI when the patient's history and Electrocardiogram (ECG) are non-diagnostic or equivocal⁵. Cardiac troponin-I (cTnI), the gold

Accepted on : 16/05/2024

Editor's Comment :

- There is a good probability that Acute Myocardial Infarction will be identified early, which could prolong or preserve a patient's life. When the patient's history and Electrocardiogram are non-diagnostic or ambiguous, cardiac biomarkers play crucial role in identification of Acute Myocardial Infarction.
- The most accurate is cTnI, which is the gold standard for identifying myocardial damage, as CKMB and cTnI are often administered to patients with symptoms suggestive of acute myocardial infarction, this study looked into their sensitivity and specificity for the better diagnosis of Acute Myocardial Infarction.

standard for detecting myocardial injury, is a highly specific biomarker to AMI⁶.

Traditionally, patients presenting with chest pain had their myoc ardial health assessed using the enzymes such as Creatinine Kinase (CK), Lactate Dehydrogenase (LDH) and Aspartate Aminotransferase (AST). These indicators were sensitive to cellular death but not specific to myocardial damage. Later, serum myoglobin levels and the Myocardial Band (MB) isoform of Creatine Kinase (CK-MB) were used to improve diagnosis and increase specificity for heart injury. The most used biomarker for the identification of myocardial necrosis over the past two decades is the

Department of Biochemistry, Dr D Y Patil Medical College, Hospital & Research Centre, Pune, Maharashtra 411018

¹MSc (Medical Biochemistry), Tutor

²MSc (Medical Biochemistry), Tutor and Corresponding Author ³PhD, Medical Biochemistry, Professor

⁴MD (Biochemistry), Professor and Head, Department of Biochemistry, Government Medical College Baroda, Gujarat 390001 *Received on : 26/09/2023*

assessment of serum troponin levels⁷. This study sought to investigate the sensitivity and specificity of CKMB and cTnl for the diagnosis of AMI because they are frequently performed in patients with complaints indicative of AMI.

MATERIALS AND METHODS

This retrospective study was performed at the SSG Hospital, Medical College Baroda between the months of December, 2018 and April, 2019. Patients who complained of chest pain were evaluated based on pre-determined criteria.Patients who underwent CKMB and cTnl between 24 to 72 hours of experiencing chest discomfort were included. They were managed according to standard protocol. Those who agreed to participate signed an informed written consent. CKMB and cTn I were done within 24 hours of the chest discomfort to calculate sensitivity, specificity, predictive value of positive test and negative test of these markers. Ethical Clearance was not applicable as this was the retrospective study done for poster presentation at the 15th Asia-Pacific Federation for Clinical Biochemistry and Laboratory Medicine Congress (APFCB) in 2019 and it was not mandatory at that period.

Thorough medical history of the individual with CKMB and cTnl data was obtained. CKMB and cTnl was performed within 24 hours of onset of chest pain and between 24 to 72 hours of chest pain by Immuno-Inhibition by blend of monoclonal antibody and chromatographic immunoassay fast card test respectively.

Depending upon the final diagnosis, patients were divided into two groups.

i) Group 1: Within 24 hours of chest pain

ii) Group 2: Within 24-72 hours of chest pain

They were further subdivided based on onset of chest pain.

i) Patients with confirmed diagnosis of MI (n=133)

ii) Patients without MI (n=235)

Statistical Analysis :

Data of CKMB and cTnI of all MI positive and MI negative patients were collected and

Sensitivity, Specificity, Positive Predictive Value and Negative Predictive Value of test (NPV) were calculated using following formula;

Sensitivity = TP/(TP+FN)*100

Specificity = TN/ (TN+FP)*100

Predictive Value of Positive test (PPV) = TP/ (TP+FP)*100

Predictive Value of Negative test (NPV) = TN/ (FN+TN)*100

OBSERVATION AND RESULTS

A total 368 patients were enrolled with the complaint of chest pain in emergency. Majority of them were >60 years of age. The Age and Gender distribution is shown in Table 1.

In our study, we found that the complaint of chest pain was more in male as compared to females. Also the complaint of chest pain increased with the age, subjects upto 40 years were lower compared to subjects more than 60 years.

Of the enrolled patients, 133 had confirmed diagnosis of MI. CKMB and cTnI were performed in these subjects and the results were plotted in 4x2 table.

Table 2 represents the values of these tests within 24 hours of onset and Table 3 represents the values of these tests within 24 to 72 hours on onset.

Table 2 shows subjects within 24 hours of onset of chest pain; only 73 were with MI positive and 155 were MI negative. Out of these CKMB found positive in 113 subjects whereas in 115 subjects CKMB was negative. In addition to this cTnI found positive in 54 subjects and negative in173 subjects.

Table 3 shows only 60 were with MI positive and 80 were MI negative. Out of these CKMB found positive in 85 subjects wereas in 55 subjects CKMB was negative. In addition to this cTnI found positive in 19 subjects and negative in121 subjects.

Further, we calculated the sensitivity and specificity performed in both study Groups and represented in Table 4.

Table 4 shows the comparision of sensitivity, specificity, PPV and NPV of CKMB and cTnI within 24 hours of MI and within 24-72 hours of MI with 95% of CI.

Fig 1 shows that CKMB is more sensitive (82.19%) within 24 hours of MI but after 24 hours the sensitivity decreases to (57.53%). In addition to this CKMB is specific after 24 hours of MI (92.26%) compared to within 24 hours of MI (65.81%). On other hand cTnI is not sensitive but more specific (95.00%) after 24 hours.

DISCUSSION

Myocardial infarction is the main cause of

Table 1 — Age and Gender Distribution				
Age	Males	Females		
Upto 40 years	9	3		
40-50 years	23	18		
51-60 years	34	28		
> 60 years	136	117		
Total (n=368)	202	166		

Table 2 –	Table 2 — Group 1 - Number of subjects with MI positive and MI negative within 24 hours onset of Chest pain					
MI Positive	Onset	CKMB>24 IU/L (TP)	CKMB<24 IU/L (FN)	cTnI Positive (TP)	cTnI Negative (FN)	
n=73	Within 24 hours	60	13	42	31	
MI Negative n=155	CKMB>24 IU/L CKMB<24 IU/L cTnl Positive cTnl Negative MI Negative Within (FP) (TN) (FP) (TN) n=155 24 hours 53 102 12 143					
MI : Myocardial Infraction, ICCU : Intensive Coronary Care Unit, CKMB : Creatine Kinase Myoglobin Binding, cTnI : Cardiac Troponin I, TP : True Positive, FP : False Positive, TN : True Negative, FN: False Negative						

Table 3 — Group 2 - Number of subjects with MI positive and MI negative within 24-72 hours of onset of chest pain						
MI Positive	Onset	CKMB>24 IU/L (TP)	CKMB<24 IU/L (FN)	cTnI Positive (TP)	cTnI Negative (FN)	
n=60	24-72 hours	38	22	15	45	
CKMB>24 IU/L CKMB<24 IU/L cTnl Positive cTnl Negative MI Negative 24-72 (FP) (TN) (FP) (TN) n=80 hours 47 33 4 76						
ITEOD Hours 47 53 4 76 MI : Myocardial Infraction, ICCU : Intensive Coronary Care Unit,						

morbidity and mortality Worldwide. An acute ST-Elevation Myocardial Infarction (STEMI) affects more than 3 million people annually, while non-ST-elevation Myocardial Infarction (NSTEMI) affects more than 4 million. NSTEMI and unstable angina are identical. Cardiovascular indicators, however, are not increased¹⁰⁻¹². The WHO states that at least two of the following three criteria must be met in order to diagnose MI: (1) a history of chest discomfort of the ischemia variety (2) progression on successive electrocardiograms (3) changes in the serum cardiac marker levels. The examination of serial cardiac indicators is currently the cornerstone of these diagnostic criteria¹³. The Acute Myocardial Infarction consequences peak in the first few hours¹⁴. Early MI diagnosis and treatment are essential for preventing myocardial damage and maintaining heart function¹⁵. Cardiac troponin and creatine kinase-MB isoform are two popular biomarkers used to diagnose Acute Myocardial Infarction.

Cytoplasmic CK is a M and/or B subunit-containing dimer that associates to generate the isoenzymes CK-MM, CK-MB and CK-BB¹⁶. Following myocardial injury, serum total CK activity and CK-MB concentration rise together, beginning to climb 4-6 hours after the damage, reaching peak serum concentrations after 12-24 hours, and reverting to baseline after 48-72 hours. Compared to serum total CK, which may be increased in many diseases where skeletal muscle is injured, serum CK-MB is far more specific for myocardial injury¹⁷. Therefore, CK should only be used in conjunction with other more precise cardiac markers to diagnose myocardial damage.^[18]

The regulation of striated and cardiac muscle contraction is carried out by the troponin

complex components; TnC, TnI, and TnT¹⁶. Troponin C is not highly selective for myocardial damage since its isoforms in skeletal and cardiac muscle are similar^{19,20}. Troponin I has not been isolated from skeletal muscle and is very selective for the cardiac muscle. It is the excellent marker of myocardial damage because of its extreme specificity²¹. After myocardial damage, they are released into the bloodstream 6-8 hours later, reach their peak at 12-24 hours and then remain high for 7-10 days²². The only drawback is the late clearance of cTn, which makes it challenging to detect a recurrence of myocardial infarction²³.

Finding the best and clear cardiac biomarker for evaluating AMI has always been an actively debated issue in research. Despite being more cardiospecific, new diagnostic assays were shown to have lower diagnostic efficacy when compared to the gold standard CK-MB⁸. The purpose of the current study

Table 4 — Sensitivity and specificity of CKMB and cTnI in subjects within 24 hours of onset of Chest pain and within 24-72 hours of onset of chest pain								
Statistics	Statistics Within 24 hours 24-72 hours						S	
	CKMB	95% CI	cTnl	95% CI	CKMB	95% CI	cTnl	95% CI
Sensitivity	82.19%	71.47% to 90.16%	57.53%	45.41% to 69.03%	63.33%	49.90% to 75.41%	25.00%	14.72% to 37.86%
Specificity	65.81%	57.77% to 73.23%	92.26%	86.87% to 95.94%	41.25%	30.35% to 52.82%	95.00%	87.69% to 98.62%
PPV	53.10%	43.48% to 62.55%	77.78%	64.40% to 87.96%	44.71%	33.91% to 55.89%	78.95%	54.43% to 93.95%
NPV 88.70% 81.45% to 93.84% 82.18% 75.68% to 87.56% 60.00% 45.91% to 72.98% 62.81% 53.56% to 71.42%								
CKMB : Cr	eatine Kin	ase Myoglobin Bindin	g, cTnl : C	ardiac Troponin IPP	/: Positive	Predictive Value, NF	V : Negat	ive Predictive Value



Fig 1 — Comparison of sensitivity, Specificity, PPV, and NPV between CKMB, cTn I and Combined CKMB with cTnI

is to evaluate the diagnostic performance of cardiac Troponin-I and CKMB for detecting acute MI in patients as soon as they arrive in the emergency room.

In 2000, Cardiac Troponin took CK-MB's status as the preferred biomarker for identifying myocardial infarctions²⁴. The early clearance aids in the identification of reinfarction, is the only advantage of CKMB over Troponin. To diagnose myocardial infarction, the serum troponin level and CK-MB fraction level are therefore measureds²⁵.

In this study, the serum troponin level is measured for the diagnosis of myocardial infarction, the level of the CK-MB fraction is measured. We found CKMB is more sensitive (82.19%) while cTnI is more specific (92.26%). Compare to the study done by PS Mahalakshmi and PS Babu found that for cTnI the sensitivity and specificity was 96% and 98%⁷ which was higher than our results and for CKMB the sensitivity and specificity was 62% and 68% which was lower than our results. Sharbari Basu,et.al. found that cTnI was 100% specific⁶, in our study it is 92.25% specific while CK-MB was a more sensitive marker in diagnosis of AMI.

CONCLUSION

In developing countries early diagnosis is crucial since AMI is the primary cause of mortality and morbidity. Serum cardiac biomarkers analysis is currently the mainstay of MI diagnosis. A serum cardiac marker should be very sensitive and highly specific in order to accurately diagnose MI.

When compared to cTnI, the current investigation discovered a statistically extremely significant rise in CKMB levels in participants who had MI at the time of admission to the emergency department. High specificity (92.26%) and PPV (77.78%) were displayed by cTnI. It is the earliest marker for the confirmation and exclusion of acute MI within 24 hours after the infarction, in addition to the specificity increases after 24-72 hours of MI (95.00%).

In conclusion, It was found that CKMB, when compared to cTnI, is a more sensitive marker and a superior diagnostic accuracy for the identification of AMI, especially in the first 24 hours after an episode of AMI whereas cTnI is more specific.

REFERENCES

- 1 Amsterdam EA, Wenger NK, Brindis RG, Casey Jr DE, Ganiats TG, Holmes Jr DR, et al — 2014 AHA/ACC guideline for the management of patients with non–ST-elevation acute coronary syndromes: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014; **130(25)**: 2354-94.
- 2 Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, et al — Heart disease and stroke statistics— 2019 update: a report from the American Heart Association. *Circulation* 2019; **139(10)**: e56-28.
- 3 Motamed H, Mohammadi M, Tayebi Z, Rafati Navaei A The diagnostic utility of creatine kinase-MB versus total creatine

phosphokinase ratio in patients with non-ST elevation myocardial infarction from unstable angina. SAGE Open Medicine 2023; **11**: 20503121221148609.

Journal of the Indian Medical Association

- 4 Goswami PP, Deshpande T, Rotake DR, Singh SG Near perfect classification of cardiac biomarker Troponin-I in human serum assisted by SnS2-CNT composite, explainable ML, and operating-voltage-selection-algorithm. *Biosensors and Bioelectronics* 2023; **220**: 114915.
- 5 Rains MG, Laney CA, Bailey AL, Campbell CL Biomarkers of acute myocardial infarction in the elderly: troponin and beyond. *Clinical Interventions in Aging* 2014; **11**: 1081-90.
- 6 Basu S, Rani UP, Srinivasan AR Association of creatine kinase MB and troponin I with electrocardiographic changes in acute myocardial infarction. *Biomed Research* 2009; **20(2)**: 84-6.
- 7 Petimani MS, Babu PS Study of cardiac troponin-i as a diagnostic marker in comparision with creatine KINASE-MB in myocardial infarction.
- 8 Daubert MA, Jeremias A The utility of troponin measurement to deyect myocardial infarction : review of the current findings. *Vascular Health and Risk Management* 2010; 691-9.
- 9 Warell DA, Cox TM, Firth JD Clinical cardiology. In: Oxford textbook of medicine 4th edition. Oxford University Press New York 2003; 2: 388.
- 11 Nascimento BR, Brant LCC, Marino BCA, Passaglia LG, Ribeiro ALP — Implementing myocardial infarction systems of care in low/middle-income countries. *Heart* 2019; **105(1)**: 20-6.
- 11 Barberi C, van den Hondel KE The use of cardiac troponin T (cTnT) in the postmortem diagnosis of acute myocardial infarction and sudden cardiac death: A systematic review. *Forensic Sci Int* 2018; **292:** 27-38.
- 12 Alaour B, Liew F, Kaier TE Cardiac Troponin diagnostic problems and impact on cardiovascular disease. Ann Med 2018; 50(8): 655-65.
- 13 Alexander JH, Sparapani RA, Mahaffey KW, Deckers JM, Newby KL, Ohman ME, et al — Association between minor elevations of creatine kinase-MB level and mortality in patients with acute coronary syndromes without ST-segment elevation. JAMA 2000; 283: 347-53.

- 14 Baheti R, Laddha P, Gehlot RS Value of Troponin-T test in the diagnosiss of acute myocardial infarction. J Indian Academy of Clinical Medicine 2002; 3(1): 55-8.
- 15 Daubert MA, Jeremias A The utility of troponin measurement to deyect myocardial infarction : review of the current findings. Vascular Health and Risk Management 2010; 691-9.
- 16 Kemp M, Donovan J, Higham H, Hooper J Biochemical markers of myocardial injury. *British Journal of Anaesthesia* 2004; **93(1):** 63-73.
- 17 Homburg JJ, Friedman DL, Perryman MB Metabolic and diagnostic signi®cance of creatine kinase isoenzymes. *Trends Cardiovasc Med* 1991; 1: 195±200
- 18 Jaffe AS, Ravkilde J, Roberts R Its time for a change to a troponin standard. *Circulation* 2000; **102:** 1216±20.
- Lewandrowski K, Chen A and Januzzi J Cardiac markers for myocardial infarction. A brief review. *Am J Clin Pathol* 2002; 118 (Suppl 1): S93-S99.
- 20 Ruseva A Laboratory diagnosis of acute myocardial infarction. *Trakia J Sci* 2005; 3: 8-14.
- 21 Higgins JP and Higgins JA Elevation of cardiac troponin I indicates more than myocardial ischemia. *Clin Invest Med* 2003; **26**: 133-47.
- 22 Tucker JF, Collins RA, Anderson AJ, Hauser J, Kalas J, Apple FS — Early diagnostic efficiency of cardiac troponin I and Troponin T for acute myocardial infarction. *Acad Emerg Med* 1997; 4: 13-21.
- 23 Mythili S, Malathi N Diagnostic markers of acute myocardial infarction. *Biomedical Reports* 2015; 3(6): 743-8.
- 24 Alpert JS, Thygesen K, Antman E, Bassand JP Myocardial infarction redefined—a consensus document of The Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of myocardial infarction. J Am Coll Cardiol 2000; 36: 959-69. 10.1016/S0735-1097(00)00804-4.
- 25 Gerhardt W, Katus H, Ravkilde J S-troponin T in suspected ischemic myocardial injury compared with mass and catalytic concentrations of S-creatine kinase isoenzyme MB. *Clin Chem* 1991; **37:** 1405-11.

JIMA now publishes Articles, submitted **ONLINE** only through https://onlinejima.com

Original Article

Prevalence of Depression and Anxiety among Elderly Patients Visiting Tertiary Care Hospital

Samreen Kour¹, Manmeet Singh², Parul Gupta³, Raghav Arora⁴, Urvi Gupta⁵

Background : Among the elderly population, the most common psychiatric conditions include depression and anxiety, which often go untreated or unrecognized, owing to the lack of knowledge and misperceptions about these conditions. As the rise in ageing population is the current trend all over the World, these morbidities may result in decreased Quality of Life among elderly population if left untreated.

Materials and Methods : The study was conducted at Tertiary Care Hospital ASCOMS on 90 cases above 60 years of age. Data was collected by doing face to face interviews using Geriatric Depression Scale -10 and Geriatric Anxiety Scale-10.

Results : The mean age of the participants was 71.8 ± 8.2 years. Most of the participants were Males (54.4%), 85.6% were physically independent and 81.1% had some physical problems. Regarding depression, 16.7% had mild Depression, 18.9% had moderate depression followed by 5.5% with severe Depression. Whereas, 7.8% had severe symptoms of anxiety. The anxiety and depression were significantly correlated with companion of living, education status, physical dependency and co-morbid physical problem.

Conclusion : Elderly cases have symptoms of depression and anxiety. Having physical health problems, being physically dependent are likely to result in poor psychological health in elderly. There is a need to recognize the mental health problem of elderly in community settings.

[J Indian Med Assoc 2024; 122(12): 52-6]

Key words : Anxiety, Depression, Elderly, Psychiatric Conditions.

A geing is an inevitable developmental phenomenon bringing along a number of changes in the physical, psychological, hormonal and the social conditions. The aged become increasingly dependent on others. As man grows, his reduced activities, income and consequent decline in the position of the family and society makes his life more vulnerable. An old person begins to feel that even his children do not look upon him with that degree of respect, which he used to get some years earlier. The old persons feel neglected and humiliated. This may lead to the development of psychology of shunning the company of others. Ageing begins with conception and terminates with death¹.

Senior citizens are at risk of developing mental and physical health problems and are particularly vulnerable to major depressive episodes and also suffer from co-morbid anxiety disorders. Depression

⁴MBBS, MD, Associate Professor

⁵BA (Honours), Psychology Student, Department of Psychiatry, Model Institute of Education and Research, Jammu and Kashmir 180001

Received on : 12/02/2024

Accepted on : 25/02/2024

Editor's Comment :

Anxiety and Depression is common among elderly patients. It is important to screen the patients with mental health issues in other medical speciality so that they can be appropriately referred to mental health professional for better care and management.

and anxiety are the most common mental health problems and were identified as leading contributors to global disability². Together with modernization, elderly in Asian countries also is living alone and are vulnerable to mental health. In such context, identification and treatment of mental health problems in older adults has become increasingly important³.

Depression is a major mental health problem, which is yet to be recognised as an important public health challenge. About 322 million persons are affected by Depression Worldwide⁴. Depression is single largest contributor to the global disability (7.50%, 2015) and suicides (about 8,00,000 annually)⁵.

In India, the elderly persons (above 60 years) constitute 8.60% of total population (Census 2011), which is expected to reach 19.0% by 2050⁶. Therefore, Depression among the elderly population is a major cause of the disease burden in future. Female gender, chronic morbidity, ignorance of elderly in household decision making, day-time spending without the work or hobbies and death of the close

Department of Psychiatry, Acharya Shri Chander College of Medical Sciences and Hospital, Jammu, Jammu and Kashmir 180017 ¹MBBS, Postgraduate Student

²MBBS, MD, Professor and Head and Corresponding Author ³MBBS, MD, Assistant Professor

relatives were identified as the significant risk-factors of Depression in elderly⁷. With this background, this study has been taken up with the objective to assess and compare Depression and anxiety among the elderly people.

MATERIALS AND METHODS

It is a cross-sectional study. This study included 90 patients with age above 60 years both, Males and Females and providing informed consent were included in the study.

The present study was conducted at tertiary care hospital. Data was collected from elderly patients attending Tertiary Care Hospital and was screened for Depression and Anxiety.

Convenience sampling technique was used to obtain the sample. Appropriate ethical clearance was obtained from the Institutional Ethics Committee and measures were undertaken to maintain confidentiality of caregivers throughout the study and also during the analysis of data. All participants were fully informed about the purpose of the study. Written informed consent was obtained from each participant after the consent form was read by the participants. The consent form was in Hindi and in English and it stated that the participant could withdraw at any time from the study. Confidentiality was maintained throughout the study.

Inclusion Criteria :

- Patients above 60 years of age.
- Both genders and providing informed consent. **Exclusion Criteria :**

 Patients with history of psychiatric illness / Dementia.

• Patients not giving informed consent and were too ill to be assessed.

Socio Demographic Data :

This arranged format was used to assess the variables of socio-demographic profile such as sex, age, marital status, religion, income, locality and type of family.

Geriatric Depression Scale (GDS-10) as mentioned by Yesavage J, *et al*⁶ was used as the study proforma. It was transformed into local vernacular language. The severity of anxiety was assessed using Geriatric Anxiety Scale (GAS). GAS was developed by Segal and colleagues in 2010. The GAS investigates three dimensions of anxiety, namely Somatic, Cognitive, and Affective ones and this factorial structure has been confirmed in recent research¹⁰.

Statistical Analysis :

Microsoft Excel was used in creating the database and producing graphs, while the data was analysed using the Statistical Package for the Social Sciences (SPSS) version 23 for Windows¹¹. Mean and Standard Deviation (±SD) were used to describe quantitative data meeting normal distribution. Continuous two independent groups were compared by parametric independent Student's t test. Discrete (categorical) groups were compared by chi-square (χ^2) test was used. P values less than 0.05 (p<0.05) was considered statistically significant.

OBSERVATIONS AND RESULTS

In present study the majority of the studied cases were in the age group ranging from 61-70 years (57.8%) with mean age 71.8 \pm 8.2 years and slight male pre-dominance (54.4%).

In 16.7% were having mild Depression, 18.9% with moderate Depression and 5.5% with severe Depression. 78.8% were having minimal to mild anxiety followed by 13.3% with moderate anxiety level and 7.8% were having severe anxiety. Depression and anxiety were significantly higher in those elderly people who were living alone, having physical health issues, need support and feel difficult to walk even with support and whose educational status is very low (p<0.05)(Tables 1-4 & Figs 1-2).

Table 1 — Demographics and Clinical Characteristics of study participants				
Variables	No of cases (n=90)	Percentage		
Age in years :				
61-70	52	57.8		
71-80	31	34.4		
>80	7	7.8		
Mean Age	71.8	8±8.2		
Gender :				
Male	49	54.4		
Female	41	45.6		
Living Companion :				
Family	61	67.8		
Husband Wife only	22	24.4		
Alone	7	7.8		
Having Physical health issues :				
Yes	73	81.1		
No	17	18.9		
Physical Dependency :				
Independent	77	85.6		
Need support	9	10.0		
Difficult to walk even with sur	port 4	4.4		
Educational Status :				
Illiterate	10	11.1		
Primary	21	23.3		
Secondary	29	32.2		
Higher School	11	12.2		
Graduate	19	21.1		

Table 2 — Prevalence of Depression and Anxiety in elderly patients						
Mental Health		No. of cases (n=90)	Percentage			
Depression	Normal	53	58.8			
	Mild	15	16.7			
	Moderate	17	18.9			
	Severe	5	5.5			
Anxiety	Minimal	4	4.4			
	Mild	67	74.4			
	Moderate	12	13.3			
	Severe	7	7.8			

Table 3 — The association between Depression and Independent variables among the study participants					
	Depre	ssion	p-value		
Variables	No (n=53)	Yes (n=37)			
Age in years :					
61-70	36	16	0.083		
71-80	13	18			
>80	4	3			
Gender :					
Male	33	16	0.075		
Female	20	21			
Living Companion :					
Family	39	22	0.039		
Husband Wife only	13	9			
Alone	1	6			
Having Physical health is	sues :				
Yes	47	26	0.028		
No	6	11			
Physical Dependency :					
Independent	50	27	0.017		
Need support	2	7			
Difficult to walk even					
with support	1	3			
Educational Status :					
Illiterate	2	8	0.002		
Primary	7	14			
Secondary	13	16			
Higher School	3	8			
Graduate	16	3			



Fig 1 — Prevalence of depression in study participants

Table 4 — The association between Anxiety and Independent			
	ay partic	apants	
variables	Ainimal	Madarate	p-value
	to Mild		;
	(n-71)	(n=10)	5
. .	(1=71)	(11=19)	
Age in years :			
61-70	45	11	0.289
71-80	22	9	
>80	4	3	
Gender :			
Male	39	10	0.858
Female	32	9	
Living Companion :			
Family	58	3	<0.001
Husband Wife only		11	
Alone	2	5	
Having Physical health issues :			
Yes	55	18	0.048
No	16	1	
Physical Dependency :			
Independent	67	10	<0.001
Need support	3	6	
Difficult to walk even with support	rt 1	3	
Educational Status :			
Illiterate	4	6	0.014
Primary	12	9	0.0.1
Secondary	27	2	
Higher School	10	1	
Graduate	20	1	
Ciududio	20		



Fig 2 — Prevalence of Anxiety in study participants

DISCUSSION

Psychiatric morbidity is one of the domains which can significantly influence the Quality of Life of elderly people. Because depression and anxiety are the most common psychiatric morbidity among elderly persons, understanding this issue is vital for comprehensive geriatric assessment and care^{12,13}.

In present study the majority of the studied cases were in the age group ranging from 61-70 years (57.8%) with mean age 71.8 \pm 8.2 years and slight male predominance (54.4%). 67.8% were living with their family and 81.1% were having physical health problems. Our findings were in accordance with the findings of Sharma M, *et al* who reported that among

the 245 senior citizens, mean age was 78.3±6 years. Majority respondents (52.2%) were female and 12(4.9%) were living alone in their home. Majority (54.2%) could not read and write, (68.2%) had some sort of physical health problems and (66.9%) were taking some types of regular medications for their health conditions². According to Kumar P, et al majority of the elderly people were in the age group of 60-69 years. Overall mean age of the elderly population was 68.55 ± 6.54 years¹⁴. Joshi OS and Ghorpade N reported that in urban group, 50.0% were in age group 60-70 years, 45.0% in 71-80 years. In rural group 50.0% were in age group 60-70 years, 45.0% in 71-80 years. In urban group according to gender, 55.0% were males and 45.0% were females and similar data was in rural also. In urban group according to education, 11.0% were illiterate, 22.0% were educated up to primary, 33.0% up to secondary, 13.0% up to higher secondary and 21.0% were graduates. In rural group 16.0% were illiterate, 35.0% were educated up to primary, 25.0% up to secondary, 13.0% up to higher secondary and 11.0% were graduates.

In our study, the 41.1% had depression, 16.7% were having mild depression, 18.9% with moderate depression followed by 5.5% having severe depression. 78.8% were having minimal to mild anxiety followed by 13.3% with moderate anxiety level and 7.8% were having severe anxiety. Our findings were consistent with the findings of Sahni B, et al who reported 40.7% of the subjects reported depressive symptoms. However, out of the 66 depressed senior citizens in the total sample, 65.0% had only mild depression while moderate to severe depression was observed in 14.2%¹⁶. Kumar P, et al reported the overall study of depression and anxiety was 66.1% and 93.8%, respectively which was higher than our study which may be because they have conducted the study on elderly living in old age homes¹⁴. Sharma M, et al reported that overall anxiety score, 64.5% senior citizens in this study had minimal, 15.5% had mild, 11.8% had moderate, and 8.2% had severe symptoms of anxiety.

Depression and anxiety were significantly higher in those elderly people who were living alone, having physical health issues, need support and feel difficult to walk even with support and whose educational status is very low (p<0.05). Our findings were comparable to the findings of previous study among similar population by Beekman AT, *et al* had found concurrent anxiety and depressive disorder in 47.5% and 21.0% of elderly. Additionally, living without partner (p=0.001) and having lower education status (p=0.000) would likely to 3 times increase respondents' chances of having anxiety. Moreover, anxiety was associated with having physical health problems and physical dependence of the respondents¹⁷. Similar to this finding, Sousa RD, *et al* found significant association of female sex and low educational level with anxiety symptoms¹⁸. Previous studies also reported higher prevalence of anxiety among women than male. However, anxiety was not significantly associated with the sex of senior citizens in this study^{19,20}.

According to Sousa RD, *et al*¹⁸ low educational levels significantly associates with Depression of elderly. On the other hand, education level, marital status, living companion and physical dependency were associated with depression in their study. A study by Perkovic R, *et al* found a significantly higher incidence of depression in male (63.0%) and anxiety among female (42.0%) respondents²¹.

Nautiyal A, *et al* stated that out of 53 depressive participants, 44 people were found to be males, reflecting depression conditions more in males²².

Another study by Rao AV, *et al* also showed that geriatric depression was more common among males²³. In contrast, Jain RK, *et al* in their study reported that prevalence of depression was found to be more in females (57.8%) than in males (45.9%)²⁴.

Elderly people living alone/living with their children without their spouse had increased rate of depression. These factors show that depression among geriatric people could be due to financial dependency on others/ loss of their partner. Similar results were reported by Nautiyal A, *et al*, *Jariwala V*, *et al*, Rajkumar AP, *et al* and Jones RN, *et al*^{22,25-27}.

Swarnalatha N study states that the prevalence of Depression was found to be inversely proportional to the literacy status. Literacy reflects economic dependence²⁸. Similar findings were observed by Rajkumar AP, *et al* and Ramachandran V, *et al* in India^{26,29}.

Based on the first wave of the Survey of Health, Ageing and Retirement in Europe (SHARE), depressive symptoms of 28,538 persons aged 50 to 89 from eleven European countries and Israel are analysed using a negative binomial regression model. The results indicate that the number of depressive symptoms measured by EURO-D scores increase with age and are higher among women than among men. When including socio-demographic characteristics, health conditions and economic strains, the association between depressive symptoms and age vanishes for men and even reverses for women. Thus, the association between age and mental health is mediated by the health and living conditions of older persons; age by itself has no explanatory power.

Limitations :

The small sample size affects generalizability.

• Being a cross sectional study, causality cannot be established.

• The data cannot be generalized for larger area as the study was carried out in Tertiary Care Hospital.

CONCLUSION

Senior citizens with Anxiety tend to have Depression too and vice versa. There are chances of increased anxiety level with lower education status and living without a partner. Having physical health problems increases the chances of having Depression. The senior citizens, living without companions and having physical health problems need to be assessed and cared for mental health status like Anxiety and Depression. Results from this study highlight the need for proper assessment and confirmation of Depression and Anxiety, especially among elderly persons.

Financial Support and Sponsorship : Nil

Conflicts of Interest : There are no conflicts of interest.

ACKNOWLEDGEMENTS

We would like to express our gratitude and acknowledge the support of Acharya Shri Chander College of Medical Sciences & Hospital. Additionally, we extend our sincere appreciation to Dr Manmeet Singh, Professor and Head of Department of Psychiatry, for his invaluable support throughout our endeavours.

REFERENCES

- Shullai BML, Mahananda M Prevalence of Anxiety and Depression among the Elderly. *International Journal of Science and Research (IJSR)* 2020; 9(1): 1725-25.
- 2 Sharma M, Bhattarai T, Sharma P Anxiety and Depression among Senior Citizens. J Nepal Health Res Counc 2021; 19(51): 305-10.
- 3 Rodda J, Walker Z, Carter J Depression in older adults. *BMJ* 2011; **343:** d5219.
- 4 Friedrich MJ Depression is the leading cause of disability around the world. *JAMA* 2017; **317(15):** 1517.
- 5 World Health Organization Depression and other common mental disorders: Global health estimates. World Health Organization. Geneva; 2017.
- 6 Agarwal A, Lubet A, Mitgang E, Mohanty S, Bloom DE Population aging in India: Facts, issues, and options [Internet]. Bonn; 2016. (Discussion Papers). Report No.: 10162.
- 7 Pilania M, Bairwa M, Khurana H, Kumar N Prevalence and predictors of depression in community-dwelling elderly in rural Haryana, India. *Indian J Community Med* 2017; 42: 13-8.
- 8 Yesavage J, Brink T, Rose T, Lum O, Huang V, Adey M, et al — Development and validation of a geriatric depression screening scale: a preliminary report. J Psychiatric Res 1982; 17(1): 37-49.
- 9 Segal DL, June A, Payne M, Coolidge FL, Yochim B Development and initial validation of a self-report assessment tool for anxiety among older adults: The Geriatric Anxiety Scale. *Journal of Anxiety Disorders* 2010; 24: 709-14.

- 10 Picconi L, Balsamo M, Palumbo R, Fairfield B Testing factor structure and measurement invariance across gender with Italian geriatric anxiety scale. *Front Psychol* 2018; 9: 1164.
- 11 IBM Corp. Released 2015. IBM SPSS Statistics for Windows, Version 23.0. Armonk, NY: IBM Corp.
- 12 Nair SS, Hiremath SG, Ramesh, Pooja, Nair SS Depression among geriatrics: prevalence and associated factors. *IJCRR* 2013; 5(8): 110-2.
- 13 Sharan P, Rai S Anxiety and Depression in Women in India. Available at: http://www.womenshealthsection.com/content/print.php3?title=gynmh001&cat=54&Ing=english. Last accessed on 20-02-2016.
- 14 Kumar P, Udayar SE, Sravan S, Arun D Depression and anxiety among the elderly persons from institutional and noninstitutional settings in the field practice area of a tertiary-care institute, Andhra Pradesh: a comparative study. Int J Med Sci Public Health 2015; 4.
- 15 Joshi OS and Ghorpade N A comparative study to assess the level of anxiety and depression among senior citizens living in urban area and rural area of Sangli district. Journal of Critical Reviews 2020; 15: 4309-12.
- 16 Sahni B, Bala K, Kumar T, Narangyal A Prevalence and determinants of geriatric depression in North India: A crosssectional study. J Family Med Prim Care 2020; 9: 2332-6.
- 17 Beekman AT, De Beurs E, Van Balkom AJ, Deeg DJ, Van Dyck R, Van Tilburg W — Anxiety and depression in later life: co-occurrence and communality of risk factors. American Journal of Psychiatry 2000; 157(1): 89-95.
- 18 Sousa RD, Rodrigues AM, Gregório MJ, Branco JD, Gouveia MJ, Canhão H, et al —Anxiety and depression in the Portuguese older adults: Prevalence and associated factors. Frontiers in Medicine 2017; 4: 196.
- 19 Afifi M Gender differences in mental health. Singapore Med J 2007; 48(5): 385-91.
- 20 Steel Z, Marnane C, Iranpour C, Chey T, Jackson JW, Patel V, et al The global prevalence of common mental disorders: a systematic review and meta-analysis 1980- 2013. Int J Epidemiol 2014; 43(2): 476-93.
- 21 Perkovic R, Sucic A, Vasilj I, Kristo B The incidence of depression and anxiety among the elderly in the area of Livno, Bosnia and Herzegovina. *Materia Socio-medica* 2018; **30(3)**: 176
- 22 Nautiyal A, Madhav NV, Ojha A, Sharma RK, Bhargava S, Kothiyal P — Prevalence of depression among geriatric people in Dehradun City of Uttarakhand, India. *J Depress Anxiety* 2015; **4**: 208.
- 23 Rao AV, Madhavan T Depression and suicide behaviour in the aged. Indian J Psychiatry 1983; 25(4): 251-9.
- 24 Jain RK, Aras RY Depression in geriatric population in urban slums of Mumbai. *Indian J Public Health* 2007; **51(2)**: 112-3.
- 25 Vishal J, Bansal RK, Swati P, Bimal T A study of depression among aged in Surat city. *National J Commun Med* 2010; 1(1): 47-9.
- 26 Rajkumar AP, Thangadurai P, Senthilkumar P, Gayathri K, Prince M, Jacob KS — Nature, prevalence and factors associated with depression among the elderly in a rural south Indian community. *Int Psychogeriatric* 2009; **21**: 372-8.
- 27 Jones RN, Marcantonio ER, Rabinowitz T Prevalence and correlates of recognized depression in US Nursing homes. J Am Geriatr Soc 2003; 51(10): 1404-9.
- 28 Swarnalatha N The prevalence depression the rural elderly in Chittoor district, Andra Pradesh. J Clin Diagn Res 2013; 7(7): 1356-60
- 29 Ramachandran V, Sarada MM, Arunagiri S Sociocultural factors in late onset depression. *Indian J Psychiatry* 1982; 24(3): 268-73.

Original Article

Evaluating Farmers' Knowledge, Attitude and Practice Regarding Pesticide Use and Its Impact on Human Health in Northern Karnataka — A Cross Sectional study

Nagendra Angirekula¹, Shailaja S Patil²

Background : In India, a significant agricultural workforce operates with many relying on Pesticides for Farming. However, handling pesticides during mixing and spraying poses high exposure risks. Detecting these hazards early falls under occupational hygiene. Our study aims to assess rural Farmers' understanding, attitudes and behaviours concerning pesticide usage. This exploration is crucial for developing strategies to promote safer agricultural methods and protect Farmers' health.

Aims and Objectives : Assessing the Knowledge, Attitude and Practice of Farmers concerning pesticide usage and its harmful effects on human health.

Methodology : In Vijayapura district, a cross-sectional survey of Farmers was conducted. Convenience sampling was used to select study participants. Data was gathered from June, 2023 to October, 2023 using a semi-structured, pre-tested questionnaire. The data was then imported into Excel. Data analysis was done using SPSS V.26.

Results : Most of the Farmers commonly report symptoms like skin irritation, headaches and burning sensation of eyes after pesticide usage. In 51% lacked knowledge of the chemical names of pesticides and this was found to be statistically associated with their level of education. In Practice, 87.3% not using long gloves it was statistically associated with income. Over 61.8% of Farmers indicated that they would rinse their eyes with water in case of accidental pesticide spillage into their eyes. Farmers noted that Grapes, Toor dal and cotton require higher pesticide application compared to Jowar and Wheat, which requires less. While over 65% of participants do not use Personal Protective Equipment (PPE), a majority adhere to changing their clothes after Pesticide usage.

Conclusion : The majority of Farmers demonstrated a positive attitude and inadequate pesticide usage procedures; they did not wear PPE when mixing or spraying. Therefore, ongoing instruction in safe mixing or spraying will broaden their understanding and aid in preventing negative health effects.

[J Indian Med Assoc 2024; 122(12): 57-60]

Key words : Occupational Health, Pesticides, Farmers Knowledge, Health Awareness, Occupational Hazards.

The highest concentration of agricultural labourers exists in Asia, notably in India, which harbours over 20% of the global agricultural workforce¹. A significant portion of these workers in India regularly uses pesticides. Over the last decade, there has been a significant surge in pesticide usage, leading to enhanced crop yields and decreased postharvest losses. Nonetheless, this extensive application has raised apprehensions about potential adverse effects on human health².

In India, inadequate safety measures taken by those who use or manufacture pesticides, incorrect product labelling, improper home storage of chemicals and industrial pollution of the environment, often as a result of improper waste disposal, are the

Received on : 13/12/2023 Accepted on : 29/01/2024

Editor's Comment :

- Farmers face health risks because of low awareness, limited income and not using safety gear, showing the need for better support and education.
- Promoting organic farming, safer pesticide options, and regular training can protect farmers and the environment.
- Stronger laws and enforcement are crucial to improve safety and encourage sustainable farming practices.

main causes of Occupational ^hazards². Mixing and spraying are the tasks associated with the highest intensity of pesticide exposure because in this stage, Farmers come into contact with the concentrated product and often have high exposure episodes³.

The significance of spray deposition on the bodies of those applying pesticides is highlighted in studies on pesticide dermal contamination. These studies indicated that the Pesticides used by farmers posed significant risks to all individuals involved⁴. Occupational hygiene encompasses the early identification and evaluation of workplace hazards⁵. Therefore, it becomes imperative to understand Farmers' perceptions of Pesticides and their safety

Department of Community Medicine, Shri B M Patil Medical College Hospital and Research Centre, Vijayapura, Karnataka 586103 ¹MBBS, Junior Resident

 $^{^{2}\}mbox{MD}$ (Community Medicine), Professor and Corresponding Author

procedures in order to provide critical information to avoid or lessen the dangers to the environment and human health associated with Pesticide usage.

The purpose of this study was to evaluate Farmers' Knowledge, Attitudes and Practices about the usage of Pesticides and their detrimental consequences on human health.

MATERIALS AND METHODS

Study Area : Vijayapura, Northern Karnataka District.

Study Population : Farmers who have utilized pesticides in their agricultural practices previously.

Study Period : July, 2023 to October, 2023

Study Design : Cross-sectional survey

Study Technique : In the district of Vijayapura, a cross-sectional study was done on farmers. The research participants were chosen through the use of convenience sampling technique. After acquiring ethical clearance from the Institutional Ethics Committee, the questionnaire was initially created in English, then translated into the local Kannada language and subsequently administered following pilot testing. Socio-demographic details, Knowledge, Attitude and Practices about the pesticide usage and its impact on farmers health related data was obtained using pre-tested semi-structured questionnaire by conducting face to face interview.

Sample Size : Given that the anticipated proportion of farmers' knowledge on the frequency of pesticide spraying is 96.6%⁶, the study would need a minimum sample size of 180 with a 96% confidence level and 3% absolute precision.

Formula used -

$$n = \frac{z^2 p^* q}{d^2}$$

Dropout rate of 10% = 180 + 18 = 198

Sampling Technique : We included Farmers who visited the District Hospital and Rural Health Centre using convenience sampling. This method was specifically chosen because it made the process convenient and accessible. During the data collection period, we focused on selecting participants who were already present at these healthcare facilities. This approach allowed us to gather valuable insights from individuals easily reachable in these particular settings.

Inclusion Criteria : Among Farmers who had used Pesticides at least once, those who gave their oral consent to participate were included.

Exclusion Criteria : Farmers who declined to give consent were excluded from the study.

Statistical Analysis :

• Microsoft Excel was used to enter the data, and SPSS (Version 26) was used for statistical analysis.

• Diagrams, percentages and frequency were used to display the results.

• Chi square test was used to determine whether Categorical Variables were associated.

RESULTS

The study enrolled 204 Farmers, with 50.5% falling within the 26-49 age range. Over 95% of participants identified as Hindu, while 46% belonged to the general category. Among the participants, 49% had no formal education and the majority were married. Approximately 70.6% reported an annual income of less than 2 lakhs. Additionally, 27.9% had been using pesticides for over 15 years

The majority of Farmers noted that Grapes, Toor dal, and Cotton require higher pesticide application, while Jowar and Wheat demand less in this region. Regarding Farmers' awareness, 51% lacked knowledge of the chemical names of Pesticides and this was found to be statistically associated with their level of education. About 74% of farmers were aware that Pesticides should be stored separately and handled with care. Only 50% of Farmers believed it was essential to read and comprehend the Pesticide container label before use.

Regarding attitudes, 71.6% believed high-quality Pesticides pose no health risks, while 87.7% deemed Pesticide use necessary. Furthermore, 82.2% indicated that immediate bathing after Pesticide use reduces poisoning. In practice, only 33.8% used masks, statistically associated with their education level. 87.3% not using long gloves, it is statistically associated with income. While over 65% of participants do not utilize Personal Protective Equipment (PPE) and 26.5% didn't change clothes postapplication. In response to accidental Pesticide exposure in the eyes, most Farmers washed their eyes with water.

In 65.2% of Farmers clean their sprayer tanks in the fields' waterways. The majority of Farmers noted that Pesticides can penetrate the body through the nose, skin and eyes. Headache, nausea, skin irritation and a burning sensation of the eyes were the most frequently mentioned symptoms after pesticide use (Tables 1-3)(Figs 1-3).

DISCUSSION

Farmers expressed their intention to take a Pesticide-poisoned person to the hospital when

Table 1 — Socio-demographic Characteristics of Farmers					
Characteristics		Frequency	Percentage		
Age group	26-49	103	50.5		
	50-69	101	49.5		
Education level	Never attended	100	49		
	Pre/primary	62	30.4		
	High school	27	13.2		
	PUC/Diploma	15	7.4		
Religion	Hindu	195	95.6		
	Muslim	9	4.4		
Cast	General	94	46.1		
	OBC	80	39.2		
	SC	20	9.8		
	ST 10 4.9				
Marital status	Married	197	96.6		
	Un married	7	3.4		
Income	<2 lakhs	144	70.6		
	> 2 lakhs	60	29.4		
Duration of	<15 years	147	72.1		
pesticide usage	> 15 years	57	27.9		

encountered, Referencing guidelines from a World Health Organisation report². Most Farmers in the study are males aged between 26 to 50 years old and married. Similar Socio-demographic traits were observed in a study conducted in Puducherry⁷.

In our research, the majority of Farmers are aware of the need to store Pesticides separately, which aligns with the findings of a study conducted in Chikkaballapur District, South Karnataka⁸. In this study, most farmers indicated that Pesticides can penetrate the body through the nose and skin. Similar research carried out in Sweden confirmed that pesticides can penetrate the body through the skin⁹.

In line with study from Uganda, our research showed that 37.3% of Farmers said that Headache was a common side effect of using Pesticides¹⁰.

Table 2 — Farmers responses about Harmful effects of pesticide Usage				
Variables		n (2	204)	
		Frequency I	Percentage	
Through which route can	Nose	95	46.6	
can pesticides enter the	Skin	64	31.4	
human body	Eyes	45	22.1	
What are the symptoms	Headache	76	37.3	
have you experienced	Nausea	44	21.6	
following pesticide usage	Skin irritation	32	15.7	
	Burning sensation of eyes	52	25.5	
If you witnessed someone	Take the person to hospita	al 182	89.2	
how would you respond	it's happened	22	10.8	
How would you respond if	Wash the exposed area			
you accidentally got	with soap and water	58	28.4	
Pesticide on your skin	Wipe it off the cloths	146	71.6	
What would you do, if	Take to nearest hospital	78	38.2	
Pesticides accidentally spilled into your Eyes	Wash the eyes with water	126	61.8	
Where you will clean	Water way in the field	133	65.2	
the sprayer tank	In the Home	71	34.8	











Fig 3 — Practice of the Farmers regarding pesticide usage

Some of the respondents among the 204 Farmers mentioned skin irritation and a burning sensation in the eyes as symptoms of Pesticide's adverse effects on health, which were similar to findings in Kuwait¹¹.

Over 61.8% of Farmers indicated that they would rinse their eyes with water in case of accidental Pesticide spillage into their eyes. Comparable results were reported by the Department of Agriculture Development in Greece in a review study¹².

In this region, most Farmers cultivate crops such as Jowar, Wheat, Toor dal, Groundnut, Cotton, Sugarcane and Grapes. Regarding application frequency, the majority mentioned that grapes require Pesticide application every alternate day, while Jowar and Wheat necessitate a single application or none at all. Commonly used Pesticide brands in this area include DAP,

KINGDOXA and PROCLAIM with a prevalent usage of DAP among the majority.

Following the data collection process, Farmers received in-depth instruction and awareness-raising sessions regarding the use of Pesticides, with a special emphasis on Pesticide handling and the significance of wearing Personal Protective Equipment (PPE). These education and sensitization sessions play a vital role in advocating for safer Pesticide handling methods to reduce health risks.

Limitation :

Implementing a convenient sampling technique with a limited sample size constrained our ability to precisely depict the wider and diverse Farming Community. As a result, this constraint hinders the universal application of the findings or the derivation of broad conclusions relevant to diverse farming communities or regions.

Table 3 — Statistical association Level of Education versus Do you know the chemical name of the Pesticide Do you know the chemical Chi-square P value name of pesticide No Yes Total Level of Never Attended 60 40 100 Education Pre/Primary 30 32 62 High School 9 18 27 8.65 0.03 PUC/Diploma 5 10 15 Total 104 100 204 Level of Education versus Do you wear Mask when spraying Pesticides Do you wear Mask when spraying Pesticides No Yes Total Level of Never Attended 76 24 100 Education Pre/Primary 38 24 62 High School 14 13 27 9.99 0.01 PUC/Diploma 7 8 15 Total 135 69 204 Annual Income versus Do you wear long Gloves when spraying Pesticides Do you wear long Gloves when spraying pesticides No Yes Total Annual <2 Lakhs 130 14 144 4.02 0.04 income >2 Lakhs 48 12 60 Total 178 26 204

CONCLUSION

Majority of the Farmers are not using the Personal Protective Equipment (Mask, Gloves, Goggles, Boots) while mixing or spraying. They displayed a positive attitude but implemented inadequate practices concerning Pesticide usage. Just a small percentage of them are aware of the negative effects of pesticides and what to do in the event of accidental spills. In conclusion, Pesticide usage among Farmers is a complex issue with both benefits and drawbacks.

Along with addressing the health risks associated with Pesticide usage, initiatives should be made to promote responsible Pesticide use, provide access to safer alternatives such as natural chemicals and support organic Farming. Sustainable agricultural practices, education and Government regulations will continue to play vital role in shaping the future of Pesticide usage in agriculture.

REFERENCES

- 1 International Labor Office. Safety and Health in Agriculture. 1-24.
- 2 WHO Guidelines on the prevention of toxic exposures -Education and public awareness activities [Internet]. International Programme on Chemical Safety. 2004. http://www. who.int/ipcs/features/prevention_guidelines.pdf
- 3 Damalas CA, Koutroubas SD Farmers' exposure to pesticides: Toxicity types and ways of prevention. *Toxics* 2016; 4(1): 1-10. PMid: 29051407 PMCid: PMC5606636. https:// doi.org/10.3390/ toxics4010001.
- 4 Tamrin PDSBHM Occupational safety and health in commodity agriculture: Case studies from Malaysian agricultural

perspective. 2015; 1-433.

- 5 Gittleman J Early detection of occupational diseases. Ind Labor Relations Rev 1988; 42(1): 126. https://doi.org/10.1177/ 001979398804200113
- 6 Devegappanavar G Occupational Health: Farmers Knowledge on pesticide usage and its harmful effects on human health in rural areas of South India. *Journal of Ecophysiology and Occupational Health* 2020; **20(1&2):** 108-13. doi:10.18311/jeoh/2020/25130.
- 7 Mohanty MK, Behera BK, Jena SK, Srikanth S, Mogane C, Samal S, et al — Knowledge attitude and practice of pesticide use among agricultural workers in Puducherry, South India. Journal of Forensic and Legal Medicine 2013; 20: 1028-31. PMid: 24237813.
- 8 Satya Sai MV, Devi Revati G, Ramya R, Swaroop AM, Eswaran Maheswari MMK— Knowledge and perception of farmers regarding pesticide usage in a rural farming Village, Southern India. *Indian J Occup Environ Med* 2019; 23(1): 32-6
- 9 Blanco LE, Aragon A, Lundberg I, Liden C, Wesseling C, Nise G — Determinants of dermal exposure among Nicaraguan subsistence farmers during pesticide applications with backpack sprayers. Ann Occup Hyg 2005; 49(1): 17-24
- 10 Oesterlund AH, Thomsen JF, Sekimpi DK, Maziina J, Racheal A, Jørs E Pesticide knowledge, practice and attitude and how it affects the health of small-scale farmers in Uganda: A crosssectional study. *Afr Health Sci* 2014; **14(2):** 420-PMid: 25320593 PMCid: PMC4196420
- 11 Jallow MFA, Awadh DG, Albaho MS, Devi VY, Thomas BM Pesticide knowledge and safety practices among farm workers in Kuwait: Results of a survey. *Int J Environ Res Public Health* 2017; **14(4)**: 340. PMid: 28338612 PMCid: PMC5409541. https://doi.org/10.3390/ijerph14040340
- 12 Damalas CA, Eleftherohorinos IG Pesticide exposure, safety issues and risk assessment indicators. Int J Environ Res Public Health 2011; 8(5): 1402-19. PMid: 21655127 PMCid: PMC3108117. https://doi.org/10.3390/ijerph8051402

<u>Review Article</u>

Cardiac Steatosis — An Emerging Entity

Aradhya Sekhar Bagchi¹, Ananda Bagchi²

In this era of obesity pandemic and increased global prevalence of Type 2 DM (T2DM) and Metabolic Syndrome (Met S), cardiovascular diseases are also increasing globally and is a real concern for morbidity and mortality. In modern lifestyle with sedentary habits, excessive intake of calories and with unfavorable genotypes leads to lipid overflow resulting in failure of Subcutaneous Adipose Tissue (SAT) to expand and store the excess circulating Free Fatty Acids (FFA). This excess FFA is being deposited as Visceral Adipose Tissue (VAT) in the major organs like Liver, Heart, Pancreas, Skeletal muscles causing Hepatic Steatosis, Cardiac Steatosis etc. Nowadays Cardiac steatosis is considered as an important predisposing factor for Diastolic Dysfunction, AV block and Sudden Cardiac Death.

[J Indian Med Assoc 2024; 122(12): 61-5]

Key words : Triglyceride, Subcutaneous Adipose Tissue (SAT), Free Fatty Acids (FFA), Visceral Adipose Tissue (VAT), Non-alcoholic Fatty Liver Disease (NAFLD).

Cardiac Steatosis (the fatty heart), also known as Clipomatosis cordis, where there is deposition of adipose tissue in the myocardium leading to fibrosis. This excessive adipose tissue deposition alters the normal physiology of the myocardium. Cardiac adiposity or steatosis has recently emerged as an important risk factor for the development of cardiovascular diseases, including diastolic dysfunction of heart and sudden cardiac death. The hypothesis is that accumulated fat impairs cardiac performance and induces structural remodeling as a result of lipotoxicity.

Now we are in the era of obesity pandemic and abdominal obesity in particular, which is associated with Insulin resistance and Type 2 DM (T2DM) causing serious metabolic derangements in our body leading to Metabolic Syndrome. Excess caloric intake and sedentary lifestyle combined with unfavorable genotype and several environmental factors result in lipid overflow, due to a failure of Subcutaneous Adipose Tissue (SAT) to expand and store the excess of circulating Free Fatty Acids (FFA)¹. Consequently, the excessive fat is accumulated into different visceral organs like Liver, Heart, Pancreas and Skeletal muscles etc.

Ectopic fat deposits have been subdivided into those with local and those with systemic effect.

²MD (Medicine), Senior Consultant, Department of Medicine, Dum Dum Municipal Specialised Hospital and Cancer Research Center, Kolkata, West Bengal 700028 and Corresponding Author *Received on : 31/08/2024 Accepted on : 18/09/2024*

Editor's Comment :

The association of Non-alcoholic Fatty Liver Disease (NAFLD) with the Cardiovascular system is quite complex and incompletely understood. NAFLD is suggested to be implicated in cardiac steatosis in many ways and seems to worsen the prognosis of both the diseases. Moreover, as there is no definitive treatment for NAFLD as yet, many CV drugs show promising results in NAFLD treatment both biochemically and histologically with fibrosis regression. So the coupling between NAFLD and Cardiac steatosis exists in true sense and despite the suggested results, further studies are needed for better understanding of the two-way liver-heart interplay and the roles of drugs in the pathophysiology and treatment of NAFLD and cardiac steatosis.

Accordingly, perivascular, myocardial, and epi/ pericardial fat have mainly local unfavorable effects, whereas visceral adipose tissue, or fat in the liver, heart or skeletal muscles have systemic effects due to the fundamental role of these organs in glucose, insulin, and lipid metabolism. Here both location and the amount of adipose tissue has got a bearing on the development of cardiovascular morbidity and mortality.

Cardiac Steatosis :

Though it is well recognised that, adipose tissue accumulation occurs in all three cardiac sites like pericardium, myocardium and epicardium, they differ in their capacity of fat accumulation. Pericardial site has got the highest capacity and cardiomyoctes have got the least. One study by Kristopher Nyman, *et al*, has shown that increased pericardial fat is more important than epiicardial fat in causing LV Diastolic dysfunction. though the exact cause of it is not known². Pericardial fat has also been reported to be associated with Insulin Resistance (IR) and also 10

¹MD (Medicine), Senior Resident, Department of Neurology, National Institute of Mental Health and Neuro Sciences, Bangalore, Karnataka 560029

years CAD risk more strongly than Epicardial fat. In Type 2 DM patients, increased myocardial Triglyceride (TG) concentration, which is an important marker for cardiac steatosis, is being related to the development of LV Diastolic Dysfunction though subclinical LV Dysfunction may also be associated with obesity and metabolic syndrome⁷. Increased myocardial fat has also adverse metabolic consequences like impaired lipid oxidation, oxidative stress and mitochondrial defects and this cardiac lipotoxicity is one of the important causes of LV Dysfunction and promotes cardiac fibrosis and apoptosis.

A recent study by Marit Graner, *et al,* has focused on different components of cardiac steatosis and its relationship with intra abdominal ectopic fat deposits and their association with different cardio-metabolic risk factors in non-diabetic individuals with or without metabolic syndrome.

In another large cohort study of 579 men with HIV with 353 men without HIV in the age group of 40 to

70 years, epicardial adipose tissue volume was more with HIV patients which is irrespective of BMI but was associated with increasing duration of combination ART.

Assessment of Cardiac Fat :

Assessment of cardiac fat can be done by using Cardiac proton Magnetic Resonance Spectroscopy (MRS) which is a reproducible and non-invasive technique for measurement of myocardial triglyceride content by measuring their unique resonance frequency when passed through a magnetic field. Both MRS & CMR techniques, allow determination of cardiac function, its lipid accumulation status and fibrosis.

Cardiac Steatosis, Non-alcoholic Fatty Liver Disease (NAFLD) & T2DM — Are they Interrelated ?

There is a vicious spiral of worsening disease when NAFLD coexists with T2DM. Current evidence suggests that, with NAFLD, both cardiac steatosis and liver fibrosis are associated with an increased risk of incident Cardiovascular Disease (CVD), although it is unclear whether any specific NAFLD histological type contributes to a differentially greater risk of CVD. Presence of either NAFLD or T2DM increases the risk and development/progression of the other: liver fat raises the risk of incident T2DM; and further progression of liver disease to liver fibrosis further increases risk of T2DM. Coexistent T2DM and NAFLD often makes it difficult to achieve good glycaemic control due to marked hepatic and peripheral insulin resistance. T2DM also increases progression of liver disease with an increased risk of NASH, advanced fibrosis, cirrhosis and/or hepatocellular carcinoma. Both NAFLD and T2DM independently increase risk of Cardiac steatosis and other cardiac diseases. Fig 1 is a schematic illustration of the vicious spiral of worsening disease that manifests when NAFLD is accompanied by T2DM, thereby increasing the risk of Cardiac steatosis and other cardiac diseases (Fig 1A), and a summary of some of the potential CVD and cardiac risk factors in NAFLD (Fig 1B)⁷.



Atherogenic dyslipidaemia
T2DM
Other metabolic syndrome features such as hypertension, abdominal obesity
Proinflammatory cytokines such as C-reactive protein, interleukin-6, fetuin-A
Reactive oxygen species
Steatotic hepatocyte-derived extracellular vesicles
Plasminogen activator inhibitor-1
Gut-derived factors (such as modified bile acids, trimethylamine oxide, lipopolysaccharide, aromatic
acid metabolites, p-cresyl sulphate, indoxyl sulphate, short-chain fatty acids, incretins)
NAFLD-related genotypes such as GCKR variants

Fig 1(B) — Risk factors for cardiovascular diseases in NAFLD

Factors that Correlate with an Increased Incidence of Cardiac Steatosis :

It includes Aging, Female Sex, Obesity, Insulin Resistance, Diabetes Mellitus, CAD, Human Immunodeficiency Virus-Related Therapies. In many Endocrine and metabolic derangements Cardiac steatosis can be a part of redistribution of visceral fat. It is possible that the patient's truncal obesity, moon facies and abdominal striae were secondary to an under recognized Cushing syndrome. Cushing syndrome, either iatrogenic due to concomitant use of mirtazapine and budesonide or a cortical adenoma, could have deepened the intra-myocardial accumulation of fat.

NAFLD & Cardiac steatosis : Non-alcoholic Fatty Liver Disease (NAFLD) is associated with an increased risk of cardiovascular disease. In NAFLD where hepatic steatosis and fibrosis are cardinal features are also associated with subclinical myocardial dysfunction. This association is linked to altered myocardial glucose uptake and subsequent deposition.

Emerging data on the interplay between NAFLD and Cardiac steatosis show a complex two-way relationship between the two conditions. On one hand, as previously stated, NAFLD was suggested to be a major risk factor for Cardiac steatosis³. On the other hand, reverse relationship is also true regarding the effect of cardiac pathology, mainly acute and chronic heart failure, on hepatic disease.

Prevention Strategies :

Due to the strong association between NAFLD and Cardiac steatosis, and the increased risk of mortality from CV events in patients with NAFLD⁶, a number of primary and secondary prevention strategies were recommended by the American College of Cardiology, the American Heart Association, the European Association for the Study of the Liver and the Italian Association for the Study of the Liver. Both NAFLD and CVD share and target the common traditional risk factors like healthy dietary pattern, moderate exercise and optimal body weight management for their Primary prevention strategies⁴. For instance, in one prospective study on 293 patients with NASH, lifestyle modifications that are beneficial for CVD (decrease in calorie intake and increase in exercise) achieved regression of fibrosis and resolution of steatohepatitis in 19% and 25% of patients, respectively reducing the incidence of cardiac steatosis also.

Modifying NAFLD-related CVD Risk : Pharmacotherapy :

At present, there are no approved pharmacological treatments for NAFLD, and the cornerstone of NAFLD management remains lifestyle modifications which includes dietary restrictions and exercise. No FDA approved treatments are available for NAFLD but guidelines recommend : Vit E (if no Type 2 DM), Pioglitazone (if DM/pre DM), Metformin and Aspirin. For Diabetes, treatment of body weight as a coprimary outcome in obese patients with GLP1RA have emerging evidence of NASH resolution and also CV risk reduction and SGLT2 Inhibitors, which have emerging evidence for reducing liver fat and enzymes as well as CV risk reduction. Therefore, this section specifically focuses on the current pharmacological treatments for modifying NAFLD-related CVD risk⁵.

Dietary Restrictions : Carbohydrate restriction has rapid benefits in hepatic steatosis. Diet with <30 gm carbohydrates will cause weight loss of 1.8% and mean reduction of liver fat of 43.8% which will return to baseline within 1 to 3 months. So it is advisable to stick to low calorie, low carbohydrate, low fat and low dietary sugar content.

Metformin Therapy :

In the management of NAFLD drugs including metformin, Thiazolidinediones (TZD) and aspirin have shown promising results. Though metformin has been very useful in improving liver functions, it failed to improve the histological features as shown in one meta-analysis of Randomized Controlled Trials (RCTs) involving a total of 417 participants Moreover, metformin was shown to improve weight loss, improved insulin sensitivity and lipid profiles (decreased LDL and increased HDL levels) in NAFLD patients, which might in turn decrease the CV risk related to NAFLD.

Thiazolidinediones (TZD) Therapy :

In another meta-analysis including 8 RCTs and a total of 516 patients evaluating TZD effect on histology of biopsy-proven NASH, TZD treatment (5 RCTs evaluating pioglitazone; 3 evaluating rosiglitazone) was associated with improved fibrosis and NASH resolution – a trend seen in patients with or without T2DM. Side effects of TZD therapy was weight gain and limb edema and because of the short duration of the trials with small sample sizes, reports of congestive heart failure or increased CV mortality could not be documented.

Aspirin Therapy :

In a recent prospective cohort study with 361 adults with biopsy-confirmed NAFLD with nine-year followup, daily aspirin use was associated with significantly lower odds for NASH and fibrosis with greatest benefit with at least four years of aspirin use – an association not seen with other non-aspirin NSAIDs. Furthermore, a cross-sectional study including 11,416 patients showed an inverse correlation between regular aspirin use (defined as 15 times in the prior month) and prevalent NAFLD, although this was limited to older men (>60 years). Aspirin, and not ibuprofen use, reduces liver fibrosis in adults with NAFLD.

GLP-1 Receptor Agonists :

GLP-1 Receptor Agonists (GLP-1R) were investigated as potential agents for the management of NAFLD. Early studies demonstrated decreasing and normalizing AST levels in T2DM patients receiving exenatide with elevated AST levels at baseline, compared to those who did not receive it. Recent meta-analyses showed association of GLP-1Ras, mainly exenatide and liraglutide and also semaglutide, with reduced Body Mass Index (BMI) and Waist Circumference (WC) and liver fat fraction.

In a recent clinical trial, the Liraglutide Efficacy and Action in NASH (LEAN) by Armstrong, *et al*, 52 overweight patients with NASH were randomized to receive liraglutide or placebo (26 in each group)⁸. The results of this small pilot study showed that patients in the experimental treatment arm had 4.3 times higher chance of histologically-proven NASH resolution along with documented CV mortality benefit (Fig 2).

SGLT2 Inhibitors+ GLP-1 RAs :

SGLT2 inhibitors are another class of anti-diabetic drugs with proven CV mortality benefit even in nondiabetics as well with recent beneficial evidences in NAFLD treatment. Recent phase III randomized controlled trial by Frías, *et al*, 695 patients with T2DM were randomized into 3 groups - First group received exenatide plus placebo, Second group received dapagliflozin plus placebo Third group received a combination of the two drugs and followed up for 28 weeks. The results of this study showed that all groups had a decrease in the traditional CV risk factors, such as Blood Pressure, HbA1c and glucose levels, that were more pronounced and superior in the group receiving both drugs. this was a very useful trial.

The trial showed that combination treatment had stronger effects than each drug alone in ameliorating markers of hepatic and cardiac steatosis and fibrosis in patients with T2DM. Another study showed that NAFLD patients with T2DM who received dapagliflozin alone had a significant decrease in Controlled Attenuation Parameter (CAP), liver stiffness and AST and GGT levels compared with controls⁸. Other



* Lipophilic statins may have additional benefit for hepatocellular carcinoma risk reduction ** Preferred for their possible antifibrotic hepatic effects

Fig 2 - Suggested treatment options for NAFLD with Cardiac Steatosis

studies also demonstrated the potential of dapagliflozin monotherapy to reduce liver fat assessed by MRI, liver injury biomarkers such as enzyme levels, and achieve histological improvement with fibrosis regression in NAFLD patients with T2DM and improvement in cardiac status.

RAAS Inhibitors :

RAAS activation was shown to be upregulated in NAFLD and to play a role in development of inflammation and insulin resistance, both of which are possible risk factors for NAFLD and Cardiac Steatosis...In an observational cohort study by Peluci, *et al* which included 118 diabetic pts with a median follow up period of 36 months have shown that ACEI or ARB reduces the histological fibrosis progression in NAFLD patients.There are some other studies with ARB which has also shown reduction of ASTlevels as well histological improvement.

Bariatric Surgery :

An effective treatment strategy for severe obesity – has been shown to cause a significant decrease in liver transaminases and histology improvement in NAFLD patients. Many studies have shown that morbidly obese pts with NASH improve remarkably when they underwent B ariatric surgery, NASH disappeared in 85% of them and levels of liver transaminases significantly decreased. Moreover, bariatric surgery was shown to improve the traditional CV risk factors reducing cardiac steatosis as well.

Vitamin E :

Vitamin E was also studied as a potential treatment for NAFLD due to its anti-oxidative properties. In patients with NASH, Vitamin E supplementation in one meta-analysis of three trials analyzing 242 patients with NASH, resulted in improved ALT levels, steatosis, lobular inflammation and ballooning but not fibrosis of the liver. However, Vitamin E supplementation was shown in other studies to have doubtful apparent effects on CV outcomes.

Obeticholic Acid, Saroglitazar and Elafibranor :

Other drugs, including obeticholic acid, saroglitazar, and elafibranor are currently being investigated for NAFLD in large clinical trials. Obeticholic acid is a bile acid derivative that can bind to and activate farnesoid X receptors, which in turn can increase insulin sensitivity, decrease hepatic gluconeogenesis, and protect against cholestasis liver injury. Elafibranor is a dual PPAR- α/δ agonist - both receptors being implicated in the activation of inflammatory changes within the liver. Saroglitazar is a dual PPAR α/γ agonist indicated mainly for the treatment of Diabetic Dyslipidemia and hypertriglyceridemia not controlled by statins. This drug is currently being investigated as a potential treatment for NAFLD in an ongoing phase 2 trial with a very promising result. Further studies are needed before more stringent recommendations can be done on the use of obeticholic acid, saroglitazar and elafibranor.

REFERENCES

- 1 Bugianesi E, Gastaldelli A Hepatic and cardiac steatosis: are they coupled? *Heart Fail Clin* 2012; **8(4):** 663-70.
- 2 Targher G, Corey KE, Byrne CD NAFLD, and cardiovascular and cardiac diseases: Factors influencing risk, prediction and treatment. *Diabetes Metab* 2021; **47(2)**: 101215.
- 3 Tuzzolo A, Febres-Aldana CA, Poppiti R Severe Myocardial Steatosis: Incidental Finding or a Significant Anatomic Substrate for Sudden Cardiac Arrest? Am J Forensic Med Pathol 2020; 41(1): 42-47.
- 4 Nyman K, Granér M, Pentikäinen MO, Lundbom J, Hakkarainen A, Sirén R, et al — Cardiac steatosis and left ventricular function in men with metabolic syndrome. J Cardiovasc Magn Reson 2013; 15(1): 103.
- 5 Lee YH, Kim KJ, Yoo ME, Kim G, Yoon HJ, Jo K, et al Association of non-alcoholic steatohepatitis with subclinical myocardial dysfunction in non-cirrhotic patients. J Hepatol 2018; 68(4): 764-72.
- 6 Ramadan MS, Russo V, Nigro G, Durante-Mangoni E, Zampino R Interplay between Heart Disease and Metabolic Steatosis: A Contemporary Perspective. J Clin Med 2021; **10(8)**: 1569.
- 7 Rijzewijk LJ, van der Meer RW, Smit JW, Diamant M, Bax JJ, Hammer S, et al — Myocardial steatosis is an independent predictor of diastolic dysfunction in type 2 diabetes mellitus. J Am Coll Cardiol 2008; 52(22): 1793-9.
- 8 McGavock JM, Lingvay I, Zib I, Tillery T, Salas N, Unger R, et al Cardiac steatosis in diabetes mellitus: a 1H-magnetic resonance spectroscopy study. *Circulation* 2007; **116(10)**: 1170-5.

JIMA Publishes only ONLINE submitted Articles through https://onlinejima.com

Case Report

Scrub Typhus with Unilateral Parotitis and Encephalopathy : A Rare Case Report and Literature Review

Umakanta Mahapatra¹, Soham Das Bakshi², Soumyadip Mukhopadhyay², Biplab Kumar Gayen³

Scrub Typhus is an acute febrile illness recently re-emerging. Scrub Typhus is a zoonosis caused by Orientia tsutsugamushi (formerly Rickettsia). Though in majority of the cases Scrub Typhus has self limiting course, it may involve any organ resulting in multiorgan dysfunction. Scrub Typhus has to be thought of in the differential diagnosis of febrile illness. Here, we report a case of Scrub Typhus presenting as unilateral parotitis and multiorgan dysfunction. Doxycycline administration rapidly altered the clinical course. Knowledge about the non-specific clinical presentations and complications of Scrub Typhus, will help in early recognition and treatment so that complications can be avoided. [*J Indian Med Assoc* 2024; **122(12):** 66-8]

Key words : Scrub Typhus, Parotitis, Multiorgan Dysfunction.

Scrub typhus is caused by *Orientia* tsutsugamushi, a mite born obligate, intracellular gram- negative coccobacilli, which is widely distributed throughout the South Pacific Asia¹. The original name of Scrub Typhus, that was given by Hashimoto in 1810, is "tsutsugamushi disease." It is transmitted by the bite of infected larvae of trombiculid mite (Chiggers). Primary vector for the disease is Leptotrombidiumdeliense. Humen are infected when they come in contact with infected mite. The hot and humid climate and tropical weather are favorable conditions for mite activity and disease transmission. .Incubation period is 6-21 days².

In India initial cases were isolated during the sociopolitical turbulent times of 1930s³ and a large number of cases were initially described during World War II among the troops posted in the hilly regions of North East India⁴. The first outbreak of Scrub Typhus in India was reported in 2003-2004, when a cluster of paediatric patients presented with fever, hepatosplenomegaly and eschar. Scrub Typhus accounts for 25.3% of acute undifferentiated febrile illness in India with a community sero-prevalence of 34.2%⁵.

Scrub Typhus is an acute febrile illness with a wide range of clinical presentation starting from self-limiting illness with rash (50%), eschar (40%), lymphadenopathy, fever, headache, myalgia to life threatening complications including encephalitis⁶, interstitial pneumonia, ARDS, acute renal failure and acute hepatic failure. This possesses a great diagnostic challenge and dilemma while approaching an ill febrile patient. It is important to differentiate Scrub typhus from dengue, leptospirosis, malaria and enteric fever etc, so that appropriate treatment can be initiated.

³MD (Medicine), Assistant Professor

Received on : 03/08/2023 Accepted on : 28/11/2023 Editor's Comment :

The first clue to diagnose a case of scrub typhus is high index of suspicion in primary care physician particularly in this subcontinental scenario. Every case of Fever of Unknown Origin (FUO) should be searched for Scrub typhus due to its wide spectrum of presentation. Hence diagnosing and treating early can easily prevent potentially lethal complications.

The main pathogenesis is either focal or disseminated vasculitis or perivasculitis that may damage organ such as the kidney, lung, liver, brain and skin⁷. Late diagnosis of Scrub typhus and inadequate treatment may lead to complications with multiorgan involvement. Though the median mortality rate of untreated case is 6%, that of treated case is 1.4%, the mortality rate may increase up to 70% without adequate treatment^{8,9}.

Despite a few studies on this topic from this region, it is still being grossly under diagnosed due to its nonspecific clinical presentation, limited public awareness, low index of suspicion among the clinicians and lack of diagnostic facilities in some places.

CASE REPORT

A 25-years-old, male, occasional alcoholic for 2 years, was admitted in Midnapore Medical College & Hospital, West Bengal with yellowish discolouration of eyes and urine for 14 days. He also complained of high grade fever for 10 days, right sided progressive painful facial swelling with difficulty in chewing and swallowing for 7 days and altered sensorium for 2 days. On admission he was febrile, dyspnoeic, pulse - 100/min, MAP - 58 mm of Hg, dry tongue with CRT >3 sec, jaundice +, SPO2 92% in room air and RBS - 85 mg/dl. One black eschar (1x1cm) was seen over the forehead and right parotid gland was swollen (Fig 2). On palpation, RUQ of abdomen exhibited tenderness without hepato- spleenomegaly. On auscultation, S1 and S2 was audible, and vesicular breath sound in both the hemithorax was decreased (Right > Left). The patient was drowsy with a GCS of 10/15(E3V3M4), bilateral flexor

Department of General Medicine, Midnapore Medical College and Hospital, Midnapore, West Bengal 721101

¹DCH, MD, DM, Assistant Professor and Corresponding Author ²MBBS, Postgraduate Trainee

plantar response and bilaterally reactive pupil without neck rigidity.

On admission, relevant investigations showed: TLC-22,600/mm3 (N: 86, L:8, M:2, E:2), Haemoglobin - 9.5mg/dl, Platelet -1,73,000/mm³, ESR - 62 millimetres/ hour, CRP - 41.53 mg/ dL, microcytic hypochromic anaemia and toxic granules in neutrophils on peripheral blood smear. Thick and thin smear for MP/ MPDA was found to be negative. Biochemical parameters revealed: serum urea - 220 mg/dl, creatinine - 2.69 mg/dL. serum Na⁺ - 132 mmol/L, serum K⁺ - 4.22 mmol/L, total bilirubin 35.79 mg/dl with conjugated bilirubin 26.03 mg/dL. ALT/AST 482/316 U/L, ALP 382 IU/ L, serum albumin 2.6 g/dl, serum amylase/Lipase 132/78 IU/L, PT



Fig 1 — CECT of neck and face showing the features of Parotitis

22.32 sec and INR 2.4. Patient was found to be nonreactive for HIV I, II, HBsAg, Anti-HCV RNA, Hepatitis A IgM, Leptospira IgM, Dengue IgM and NS 1ag. After 48hour, 3 sites blood (both aerobic and aenorobic), and urine culture and sensitivity showed no growth. Sputum for AFB /CBNAAT and subsequently TB Quantiferron Gold was found to be negative. USG whole abdomen showed mild hepatomegaly, grade I fatty infiltration and increased echogenicity of Kidney (B/L). Chest X-Ray AP in supine position showed right sided minimal pleural effusion. Scrub IgM antibody with ELISA method was found to be positive, which was confirmed by Immunofluorescence method.

Patient was shifted to CCU, where he was conservatively managed with IV fluid, vasopressors, broadspectrum antibiotics, 6-units of FFP and subsequently 2 settings of bedside haemodialysis. After diagnosis of Scrub typhus was made, patient was put on injection IV Doxycycline 100 mg BD.

2D Echo with M mode was done to rule out vegetations. CSF study revealed lymphocytic predominance with 12 WBCS, sugar - 52 mg/dL and protein - 75mg /dl, with a negative Gram Stain and Z-N Stain. CT scan of the neck and face showed the features of Parotitis (Fig 1). FNAC was suggestive of necrotizing Parotitis.

After a 10 days course of IV doxycycline, the patient showed dramatic clinical improvements. He became afebrile, swelling decreased (Fig 2), mentation and appetite improved. The laboratory parameters before and after the treatment are enlisted below Table 1.

DISCUSSION

Scrub Typhus affects one million people around annually. Trombiculid mite (Chiggers) of the Leptotrombidium genus, which feeds on wild rats, infects human accidentally who are living in Sub Himalayan, dense forest area or the people whose livelihood depend



Fig 2 — Disappearance of Parotid Swelling after 10 days of Doxycycline therapy: right (before), I

mainly forest and agriculture². The ricketssial diseases which were once thought to have disappeared from India, are re-emerging as zoonosis in several parts of India such as Kashmir, Himachal Pradesh, Sikkim, Darjeeling (West

Table 1 — Comparison of the laboratory parameters before and after Doxycycline therapy				
Before	Parameters	After		
22,600	TLC (/mm ³)	4,300		
41.53	CRP (mg/dl)	3.56		
220	Urea (mg/dl)	16		
2.39	Creatinine (mg/dl)	0.62		
35.79	Bilirubin (mg/dl) (total)	4.25		
26.03	Bilirubin (mg/dl) (conjugate)	3.15		
316	AST (IU/L)	56		
482	ALT (IU/L)	78		

Bengal) and Western Ghat. Aetiological factors once thought eradicated have re-emerged due to migration of labour from other states, poor hygiene in public places and improper waste disposal.

The overwhelming clinical manifestations of Scrub Typhus are attributed to systemic vasculitis and endothelial dysfunction¹⁰. Signs and symptoms of Scrub Typhus are nonspecific and clinical features may mimic other febrile illness. Though most of the cases are self limiting in nature with spontaneous recovery, delay in diagnosis and inadequate treatment may lead to complications including Multi Organ Dysfunction (MOD). Though eschar is the most useful diagnostic clue of Scrub Typhus, many Indian studies revealed eschar in less number of patients¹¹.

Early use of antibiotics can prevent life-threatening scrub typhus. The effective antibiotics against Scrub Typhus are doxycycline, chloramphenicol, tetracycline and azithromycin. Azithromycin may be more tolerable than doxycycline with same effect. The clinicians should avoid rifampicin as a first-line agent due to the low-certainty evidence and the risk of resistance in undiagnosed tuberculosis¹².

Similar presentation of Scrub Typhus had been previously reported from Kasturba Medical College, Manipal in 2017, where a 38 year old male presented with fever, altered sensorium, parotid swelling and MODS¹. AKI was reported in 13-27% of patients in two different Indian studies. Parotitis is rare clinical manifestation of Scrub Typhus with MODS. Only a very few literature is available to support its rampant in southern districts of West Bengal¹³. This case report will help the clinicians to get insight about the diversity of this potentially fatal condition and increase the index of suspicion while dealing with an acute febrile patient with MOD.

CONCLUSION

Rickettsial diseases are prevalent but remain underdiagnosed in India. Scrub Typhus should be differentiated from other causes of febrile illnesses of short duration because of similar clinical and laboratory features. When patients present with fever and multi organ involvement, Scrub Typhus should always be considered in the differential diagnosis. High index of suspicion and prompt initiation of treatment will reduce the morbidity and mortality. **Declaration of patient consent :** The authors certify that they have obtained all appropriate patient consent. **Financial support and sponsorship :** Nil.

Conflict of interest : There are no conflicts of interest.

REFERENCES

- Mahajan SK Scrub typhus. J Assoc Physicians India 2005; 53: 954-8. PMID: 16515236.
- 2 Mohanty A, Kabi A, Gupta P, Jha M, Rekha US, Raj A Scrub typhus – A case series from the state of Sikkim, India. International Journal of Critical Illness and Injury Science 2019; 9(4): 194.
- 3 Roy BC Typhus fever; with special reference to its incidence in India. J Indian Med Assoc 1945; 15: 135-46. [PubMed] [Google Scholar]
- 4 Sayen JJ, Pond HS, Forrester JS, Wood FC Scrub typhus in Assam and Burma; a clinical study of 616 cases. *Medicine* (*Baltimore*) 1946; 25: 155-214. [PubMed] [Google Scholar]
- 5 Devasagayam E, Dayanand D, Kundu D, Kamath MS, Kirubakaran R, Varghese GM — The burden of scrub typhus in India: A systematic review. *PloS Negl Trop Dis* 2021; **15(7)**: e0009619.[CrossRef]
- 6 Dhanaraj M, Kar A, Dedeepiya D, Harikrishna K Acute encephalitis syndrome following scrub typhus infection. *Indian Journal of Critical Care Medicine [Internet]* 2014 [cited 2019 Nov 26]; **18(7):** 453-5. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4118511/1
- 7 Dogra S Recent advances in understanding pathophysiology of scrub typhus. JK Sci 2010; **12(2):** 19-23.
- 8 Taylor AJ, Paris DH, Newton PN A systematic review of mortality from untreated scrub typhus (Orientia tsutsugamushi). *PLoS Negl Trop Dis* 2015; **9(8):** e0003971.
- 9 Bonell A, Lubell Y, Newton PN, Crump JA, Paris DH Estimating the burden of scrub typhus: a systematic review. *PLoS Negl Trop Dis* 2017; **11(9):** e0005838
- 10 Loscalzo J, Fauci AS, Kasper DL, Hauser SL, Longo DL, Jameson JL— Harrison's principles of internal medicine. 21st ed.Vol1,page320-21 New York: McGraw Hill; 2022.
- 11 Mahajan SK, Rolain JM, Kashyap R Scrub typhus in Himalayas. *Emerg Infect Dis.* 2006;12:1590–1592. [PMC free article] [PubMed] [Google Scholar]
- 12 10.El Sayed I, Liu Q, Wee I, Hine P. Antibiotics for treating scrub typhus. *Cochrane Database Syst Rev* 2018; 9: Cd002150.
- 13 Sarma N, Chakraborty S Scrub Typhus in Southern Districts of West Bengal 2017; 62(5): 512.

Submit Article in JIMA — Online See website : https: // onlinejima.com

Any queries : (033) 2237-8092, +919477493027; +919477493033

Case Report

Neuropsychiatric Lupus with Apla Syndrome and Auto-immune Haemolytic Anaemia in a Patient of Hansen's Disease — A Diagnostic and Therapeutic Challenge

Sudipto Chakraborty¹, Parthajit Das², Kishalaya Karan³, Jayanta Sharma⁴

A 21-year-old gentleman of Hansen's disease on multi drug therapy presented with complaints of fever for 10 days and altered mental status for the last 5 days from the day of presentation. MRI brain which showed multiple infarcts in brain. He was evaluated with relevant investigations which were suggestive of Neuro Systemic Lupus Erythematosus with secondary Anti-phospholipid Syndrome and Auto-immune Haemolytic Anaemia. Immuno-suppressive therapy initiated with pulse dose of IV Corticosteroids but discontinued due to increasing TLC and new onset fever. Repeat Urine culture showed Candida tropicalis and he was started on IV Caspofungin. Then, Immuno-supressive therapy with IV Cyclophosphamide started. He had remarkable improvement in sensorium, his fever remitted and started to walk with help. He was discharged with oral steroids, warfarin and advice of IV Cyclophosphamide as per protocol.

[J Indian Med Assoc 2024; 122(12): 69-71]

Key words : APLA, Neuropsychiatric Lupus, Hansen, AIHA.

ansen's disease is caused by slow-growing acid-fast rod-shaped bacilli Mycobacterium leprae. It is a chronic infectious disease that affects the skin, peripheral nerves, upper respiratory tract, eyes and lining of the nasal mucosa and the diagnosis is done clinically¹.

Systemic Lupus Erythematosus (SLE) is a multisystem auto-immune disorder with multi-factorial causes which influence the loss of immunological tolerance against self-antigens leading to the formation of pathologic auto antibodies that cause tissue damage through various mechanisms and affects females more than males².

Anti-phospholipid Syndrome (APS) is a multi-system autoimmune disorder characterised by persistent presence of anti-phospholipid antibodies directed against phospholipid-binding proteins and may present as venous and arterial thrombosis and or pregnancy loss. The most common sites of venous and arterial thrombosis are lower limbs and cerebral arterial circulation, respectively. The anti-phospholipid antibodies are anti-cardiolipin antibody IgG and IgM, anti-beta2-gp1 antibody IgG and IgM and Lupus anticoagulant. APS can be primary, without any previous underlying disease or secondary, having underlying disease³.

CASE REPORT

A 21-year-old gentleman with previous history of Hansen's disease on multidrug therapy of Dapsone, Rifampicin and Clofazimine from rural India presented to the Emergency Department with complaints of fever for 10 days and altered mental status of drowsiness and

³MD, DM, Senior Consultant, Department of Neurology ⁴MD, Senior Consultant, Department of Medicine

Received on : 21/06/2023

Accepted on : 02/08/2023

Editor's Comment :

- APLA syndrome is common in SLE patients
- APLA syndrome can present as a very rare complication in patients with Hansen's disease
- Hansen's disease can mimic as Lupus. There are multiple reports where Hansen's disease were misdiagnosed as Lupus initially. Biopsy should be done when there is any diagnostic dilemma.
- Hansen's disease may also act as a trigger to cause SLE.
- Active infections should always be ruled out for patients planned for therapeutic immunosuppression. Presence of active Urinary Tract Infection delayed the commencement of immunosuppressive drugs in our patient.

restlessness for 5 days from the day of presentation. Fever was insidious in onset, gradually progressive, intermittent and low grade in nature. He also developed altered mental status of drowsiness and restlessness for the last 5 days following which family admitted him.

On further inquiry, family also said that he had headache, occasional nausea and vomiting for the same duration and one episode of loss of consciousness 2 days back. He did not experience any episode of neck pain, photophobia, sore throat, chest pain, cough, palpitations, abdominal pain, yellowish discoloration of eyes and urine, reduced urinary output or leg pains.

On examination, he was conscious and alert but drowsy at times and was only obeying to simple commands. Pulse rate was 110 per minute, regular in rhythm, normal in volume, normal in character, all the peripheral pulses were palpable and there was no radio-radial or radio-femoral delay. Respiratory rate 22 per minute regular, thoracoabdominal in nature and no accessory muscles of respirations were working. BP 90/60 mm Hg. Temperature 100°F. Pallor was present, there was no cyanosis, icterus, clubbing or oedema. No lymph node was palpable. Jugular vein pressure was not raised. Multiple hypopigmented patches of varying sizes along with multiple digital gangrene were present in both upper and lower limbs.

Apollo Gleneagles Hospitals, Kolkata, West Bengal 700054

¹MBBS, DNB Trainee, Department of Medicine and Corresponding Author

 $^{^2\}mbox{MD},$ MRCP, CCT, Senior Consultant, Department of Rheumatology

On neurological examination, GCS was E4V2M6. He was only obeying simple commands. There was neck stiffness; Kernig's sign and Brudzinski's sign were absent. Cranial nerve examination was within normal limits. There was reduced power in all the 4 limbs which was suggestive of quadriparesis. Deep tendon reflexes were all exaggerated with ankle clonus. On superficial reflex, Babinski's sign was positive. Sensory examination only revealed decreased sensation in the hypopigmented areas of the body along with multiple digital gangrene in both upper and lower limbs.

All the relevant investigations for fever work up were sent. He was initiated on IV fluids, empirical IV Meropenem, IV Vancomycin with IV antiviral Acyclovir and pulse dose of IV Methylprednisolone prophylactically. He did not improve with for the following treatment and there was persistent fever. MRI brain was done which detected multiple thrombo-embolic stroke. CSF studies detected increased protein level. Dermatological consult was taken in view of Hansen's disease and patient was continued on multi drug therapy of Dapsone, Rifampicin and Clofazimine. Rheumatology consult was taken. Additional blood reports were suggestive of Systemic Lupus Erythematosus with probable secondary antiphospholipid syndrome but could not be initiated on immuno-suppressive therapy in view of persisting fever, increasing trend of Total Leukocyte Count and Catheter related Urinary Tract Infection with Candida tropicalis. IV Caspofungin was initiated and IV Methylprednisolone was stopped. His blood reports showed a decreasing trend of haemoglobin levels and direct Coombs' test was positive. He was then initiated on with oral corticosteroids, oral Hydroxychloroquine and IV Cyclophosphamide on day 17 of admission. His fever subsided and his general condition improved dramatically. He was discharged in a hemodynamically stable afebrile condition with oral steroids and warfarin. On follow-up after 14 days of discharge, he had remarkable improvement, was walking and eating with assistance and is planned for further Cyclophoshamide doses.

OTHER REPORTS

DAY 1 : INR 1.21; COVID 19 RTPCR, MPDA, Dengue NS1 and IgM/IgG, Scrub Typhus IgM, Leptospira IgM, HBsAg, ANTI-HCV, HIV 1 AND 2, Blood CS and Urine CS Negative; Urine RE Normal.

EEG BRAIN : suggestive of encephalopathic pattern.

MRI BRAIN : Multiple scattered T1 hypotense, T2/ FLAIR hypertense intraparenchymal non-enhancing lesions of variable sizes showing intense diffusion restriction are seen in the bilateral basal ganglia, thalami, bilateral insular and fronto-parietal cortex, midbrain. Possible septic-embolic encephalitis or multiple cardioembolic infarct (Fig 1).

DAY 2 : Trans-esophageal Echo No Abnormality Detected; CSF Studies- Cell Count 3, Cell Type 100% Lymphocyte, Glucose 75 (Serum Glucose 142), Protein 110, TB Gene Xpert/ Gram Stain/AFB Stain/CNS Comprehensive Panel All Negative.

DAY 4 : ANA 3+ (1:180 titre in ELISA) (Fig 2)

DAY 7: ANTI ds-DNA Positive (>200); Antibody to nRNP 2+, SMITH 1+, SS-A 3+, Ro-52 3+, Rib P-Protin 2+, AMA-M2 2+; c-ANCA Negative; p-ANCA 3+ (Atypical Pattern); SERUM C3 59; SERUM C4 10; RA Factor Negative; Serum ACE 19; URINE ACR 21.

DAY 10 : Blood CS Negative; Urine CS Candida Tropicalis growth.

DAY 11 : Lupus Anticoagulant Negative; Anti Cardiolipin Antibody IgM Reactive; Anti Phospholipid Antibody Negative; BETA2 GP1 Antibody Negative.

DAY 16 : Direct Coombs' Test Positive; Stool For Occult Blood Negative (Table 1).

DISCUSSION

This patient of Hansen's disease on MDT and Systemic Lupus Erythematosus developed Secondary Antiphospholipid syndrome with autoimmune haemolytic anaemia which was improved after administration of IV Corticosteroids and IV Cyclophosphamide.

The main differentials that required to be ruled out for this patient initially were Lucio phenomenon due to Hansen's disease and Acute Systemic Lupus



Fig 1 — MRI Brain : Multiple scattered T1 hypotense, T2/FLAIR hypertense intraparenchymal non-enhancing lesio

Table 1 — Regular Blood Profile during Hospital stay							
Day of Admission	Day 1	Day 3	Day 7	Day 10	Day 14	Day 17	Day 21
Haemoglobin	9.5	9.3	9.8	8	7.6	8.5	9.1
TLC	19800 N81L7	12400 N84L12	21300 N84L5	17600 N71L11	15000 N70L15	7600 N62L3	311800 N84L11
Platelet	1.98	1.84	1.69	2.06	3.73	3.24	3.73
CRP / Procalcitonin		8.1/0.14			9.3/0.11		
Urea / Creatinine	57/0.9	72/0.8	45/0.6	32/0.7	30/0.6	39/0.5	36/0.6
Sodium / Potassium	136/4.4	135/4.2	126/4.1	133/3.3	127/4.2	130/4.1	126/4.2
Total / Unconjugated Bilirub	in 0.8/0.6	0.6/0.5	1/0.7	1.4/0.9	1/0.7		0.6/0.5
Albumin / Globulin	2.7/3.3	2.6/3.2	1.9/2.3	2.5/2.4	2.7/2.8		2.9/3.1
SGOT / SGPT / ALP	175/65/80	66/71/44	36/38/51	29/38/74	27/28/830		25/40/87

A settime to be a settime of the set of	Docted 4	14 antihody samely	: O ine Blot Assault
And a second sec	a second and		- familie - manual - ma

Antibody to n RNP/Sm	POSITIVE (2+)		
Antibody to Sm(Smith)	POSITIVE (1+)		
Antibody to SS-A(Re)	POSITIVE (3+)		
Antibody to SS-B(La)	NEGATIVE		
Argbody to Ro-52	POSITIVE (3+)		
Pody to Se170	NEGATIVE		
Antibody to PM-Scl	NEGATIVE		
Antibody to Jo-1	NEGATIVE		
Antibody to CENP B	NEGATIVE		
Antibody to PCNA	NEGATIVE		
Antibody to de DNA	NEGATIVE		
Antibody to Nucleosomes	NEGATIVE		
Antibody to Histones	NEGATIVE		
Antibody to Rib.P.Protein	POSITIVE (2+)		
Antibody to AMA-M2	POSITIVE (2+)		
Report Status Final			

Fig 2 — 14 antibody ANA panel suggestive of Neuropsychiatric Lupus

Erythematosus flare up, apart from the definitive diagnosis of Secondary APS which was contributed by Hansen's disease and or SLE.

Diagnosis of APS is based on a combination of clinical features and diagnostic findings as per revised Sapporo APS Classification Criteria. Clinical criteria include (if any one present): vascular thrombosis and pregnancy morbidity. Laboratory criteria include (if any one present): IgG and or IgM cardiolipin antibody, IgG and or IgM anti beta2 glycoprotein and Lupus anticoagulant⁴. This patient had clinical criteria of vascular thrombosis and laboratory criteria of positive IgM anti cardiolipin antibody.

Hansen's disease is associated with secondary Antiphospholipid syndrome- with previous documentations of case reports published previously as by Kaliyadan, *et al* in 2009 from India⁵. Previous studies of Hansen's disease associated APS has shown predominance of IgM subtype of antibodies of APS⁶.

One of the most common causes of secondary APS is Systemic Lupus Erythematosus which is around 35 per cent of all APS patients⁷. APS is one of the main causes of thrombosis risk in patients of SLE with the presence of anti-phospholipid antibodies as they attach to negatively charged phospholipid surface that may induce platelet activation, interfere with coagulation inhibitors and antifibrinolytics and initiate formation of a thrombus⁸.

Lucio phenomenon is an unusual presentation of Hansen's disease, a form of cutaneous vasculitis probably mediated by immune-complex deposition and present as large, sharply demarcated ulcerative lesions and thrombosis of bigger vessels and dermis and is treated by anti-leprotic medications, systemic glucocorticoids and other supportive medications⁹.

Systemic Lupus Erythematosus flare up is defined as a measurable increase in disease activity in one or more organ systems involving new or worse clinical signs and symptoms and/or laboratory measurements. It must be considered clinically significant by the assessor and usually there would be at least consideration of a change or an increase in treatment¹⁰.

Hansen's disease and SLE can mimic each other and having one can increase the risk of the occurrence of other disease¹¹. Also, there have been case reports that a case of Hansen's disease has been misdiagnosed as SLE¹².

Auto-immune haemolytic anaemia in primary APLA syndrome is very rare. Co-existence of auto-immune haemolytic anaemia and primary APLA may define as a subgroup of patients who may later develop SLE¹³.

Hence, we have been able to diagnose a patient of Systemic Lupus Erythematosus with Hansen's disease and secondary APLA syndrome and auto-immune haemolytic anaemia, which is very rare to come across.

CONCLUSION

Hence, we present a rare case of Neuropsychiatric Systemic Lupus Erythematosus with secondary Anti-Phospholipid Syndrome and auto-immune haemolytic anaemia in the background of Hansen's disease treated successfully with immuno-suppressive and anti-thrombotic therapy. It was a big challenge for coming to a diagnosis considering the presentation and even bigger challenge was the treatment course which was done judiciously.

REFERENCES

- National Institute of Allergy and Infectious diseases. Leprosy (Hansen's Disease). https://www.niaid.nih.gov/diseases-conditions/leprosy-hansens-disease.
- 2 Angel A Justiz Vaillant; Amandeep Goyal; Matthew Varacallo. Systemic Lupus Erythematosus. https://www.ncbi.nlm.nih.gov/books/NBK535405.
- 3 Jean G Bustamante; Amandeep Goyal; Mayank Singhal. Antiphospholipid Syndrome. https://www.ncbi.nlm.nih.gov/ books/NBK430980.
- 4 Diagnosis of antiphopholipid syndrome topic 4678 version 33.0. https://www.uptodate.com/contents/diagnosis-ofantiphospholipid-syndrome.
- 5 Kaliyadan F, Bhaskaran M, Dharmaratnam AD, Manoj J, Sreekanth G — Anti-phospholipid syndrome preceding a diagnosis of lepromatous leprosy. https://escholarship.org/uc/ item/2g75646r.
- 6 de Larrañaga GF, Forastiero RR, Martinuzzo ME, Carreras LO, Tsariktsian G, Sturno MM, Alonso BS. High prevalence of antiphospholipid antibodies in leprosy: evaluation of antigen reactivity. PMID: 11035434. https://pubmed.ncbi.nlm.nih.gov/ 11035434.
- 7 Pathogenesis of antiphospholipid syndrome topic 4676 version 23.0. https://www.uptodate.com/contents/pathogenesisof-antiphospholipid-syndrome.
- 8 Ibrahim A Al-Homood. Thrombosis in Systemic Lupus Erythematosus: A Review Article. https://www.hindawi.com/ journals/isrn/2012/428269.
- 9 Kasper DL, Braunwald E, Fauci AS Harrison's Principles of Internal Medicine. 16th edition. *McGraw-Hill* 2005; 1: 969.
- 10 Christina Adamichou and George Bertsias. Flares in systemic lupus erythematosus: diagnosis, risk factors and preventive strategies. PMCID: PMC7045928. PMID: 32185248. https:// www.ncbi.nlm.nih.gov/pmc/articles/PMC7045928/ #:~:text=Various%20SLE%20flare%20definitions%20 have,increase%20in%20proteinuria%20in%20the.
- 11 Horta-Baas G Multibacillary leprosy mimicking systemic lupus erythematosus: case report and literature review. PMID: 25761657. https://pubmed.ncbi.nlm.nih.gov/25761657.
- 12 Kumar, Rajiv Ranjan A Case of Leprosy Misdiagnosed as Lupus. https://journals.lww.com/jclinrheum/Fulltext/2020/12000/ A_Case_of_Leprosy_Misdiagnosed_ as_Lupus.20. aspx.
- 13 Bhagwat SN, Sharma JH, Patale DS A rare combination of autoimmune haemolytic anaemia and primary antiphospholipid syndrome evolving into systemic lupus erythematosus. Glob J Transfus Med 2016; 1: 88-90. https://www.gjtmonline.com/ article.asp?issn=2455-8893;year=2016;volume=1; issue=2;spage=88;epage=90;a ulast=Bhagwat#:~:text= Autoimmune%20hemolytic%20anemia%20(AIHA)% 20in,systemic%20lupus%20erythematosus %20(SLE).

Letter to the Editor

[The Editor is not responsible for the views expressed by the correspondents]

Does only Mupirocin enough for Nasal Decolonization in Methicillin-resistant *Staphylococcus Aureus* Carriers or Do we need more ?

SIR, — Staphylococcus aureus, particularly, its resistant variant Methicillin-resistant S Aureus (MRSA), is one of the most commonly prevalent nosocomial pathogens, creating a concern for the world population. For the treatment of severe MRSA infections, vancomycin is the drug of choice for physicians¹. However, due to the emergence of vancomycin-resistant MRSA, a topical glycopeptide antibiotic, Mupirocin is now frequently used for nasal decolonization of patients and Healthcare Workers (HCWs) to prevent the transmission of MRSA in a hospital setup². But few recent studies have documented Mupirocin-resistant MRSA, making the situation even more critical and raising the question, what is next? Or should we move to alternative preventive strategies?

During a sudden hospital outbreak of MRSA, the Hospital Infection Control Committee (HICC) has to initiate nasal swab surveillance from staff to find out the source of infection. If the source is identified, the treatment strategy will be as per HICC protocol. If no staff is found to have MRSA positive, a root cause analysis would need to be conducted, which could lead to the OPD or emergency department, which serves as the hospital's main entrance. If MRSA grows in any specimen, confirm it again and start Mupirocin treatment for 5-7 days in accordance with institutional policy.

For infection prevention, isolation of the source is also necessary in addition to treatment. Once the treatment is completed, fresh samples must be collected, and if MRSA does not grow, the treatment is successful. But if there is a growth of MRSA, treatment should be resumed and the sample should be collected again after completion of treatment. If again MRSA grows, the medical history of the source Health Care worker needs to be investigated further. Many times, medical history can reveal the reason behind MRSA growth even after mupirocin treatment. Mupirocin susceptibility of an isolated MRSA strain can be tested, and if require (colonization persists even after completion of two courses of mupirocin treatment or specimen detect mupirocin resistance), another drug such as chlorhexidine or Rifampicin, which is not often used to treat MRSA, can be added³. After the treatment is completed, collect the sample again to check for MRSA growth. Usually, after combination therapy, there will be a lesser chance of MRSA colonization still existing. In our center, a similar incident was experienced before, and we addressed it in accordance with institutional HICC protocol and as described above.

For the treatment of MRSA nasal carriage in healthy adult patients and HCWs, nasal formulation of mupirocin is frequently advised. It interferes with bacterial protein synthesis but continuous genetic evolution and the emergence of resistant strain causes treatment failure⁴. Nasal swab surveillance is necessary at regular intervals as well as during the outbreak of MRSA⁵. The purpose of surveillance is not only to collect samples and make treatment recommendations but also to retest samples after treatments to ensure that they were successfully completed⁶.

In conclusion, we would like to state that, although Mupirocin is the cornerstone of decolonization regimens, the existence of mupirocin resistance in MRSA strain is a cause for serious concern. It could be limited by the detection of Mupirocin resistance, regular surveillance, and effective infection control initiatives including standard precautions and environmental cleaning⁷. Nasal swab collection for MRSA carriers' detection during the preemployment health checkup could be a successful strategy to prevent healthcare-associated staphylococcal infections. Based on our real experience, we also strongly suggest that during MRSA nasal carrier surveillance, root cause analysis, which includes the entrance of the hospital setup, must be done, even if there is some positive evidence scattered here and there⁸.

REFERENCES

- Qureshi D, Gandham NR, Das NK, Vyawahare CR, Bhaumik S, Kannuri S, et al — Susceptibility of Contemporary Staphylococcus aureus Isolates to a Novel Antibiotic Levonadifloxacin: A Report from a Tertiary Care Center. Medical Journal of Dr. D.Y. Patil Vidyapeeth:, June 02, 2023. | DOI: 10.4103/ mjdrdypu.mjdrdypu_370_22
- 2 Agarwal L, Singh AK, Sengupta C, Agarwal A Nasal carriage of Methicillin- and Mupirocin-resistant S. aureus among health care workers in a tertiary care hospital. J Res Pharm Pract 2015; 4: 182-6.
- 3 Labercque S, Shah S, Fergus D, Parry FM Mupirocin susceptibility of staphylococci 2022: Is it time for a change in MRSA decolonization protocols? *Amerijan J Inf Cont* 2023; 51(7):725-8 https://doi.org/10.1016/j.ajic.2022.08.025.
- 4 Van rijen M, Bonten M, Wenzel R, Kluytmans J Mupirocin ointment for preventing Staphylococcus aureus infections in nasal carriers. *Cochrane Database Syst Rev* 2008; **2008(4)**: CD006216. doi: 10.1002/14651858.CD006216.pub2. PMID: 18843708; PMCID: PMC8988859.
- 5 Poovelikunnel T, Gethin G, Humphreys H Mupirocin resistance: clinical implications and potential alternatives for the eradication of MRSA, *Journal of Antimicrobial Chemotherapy* 2015; **70(10)**: 2681-92. https://doi.org/10.1093/jac/dkv169
- 6 Ouidri MA Screening of nasal carriage of methicillinresistant *Staphylococcus aureus* during admission of patients to Frantz Fanon Hospital, Blida, Algeria. *New Microbes New Infect* 2018; **23**: 52-60. doi: 10.1016/j.nmni.2018.02.006. PMID: 29692907; PMCID: PMC5913062.
- 7 Mukhida S, Palal D, Vyawahare C, Mali V, Das NK Is reuse of surgical and N95 masks dangerous? An opinion. J Patient Saf Infect Control 2022; 10: 57-8. DOI: 10.4103/jpsic.jpsic_34_22.
- 8 Liang SY, Riethman M, Fox J Infection Prevention for the Emergency Department: Out of Reach or Standard of Care? *Emerg Med Clin North Am* 2018; **36(4):** 873-87. doi: 10.1016/ j.emc.2018.06.013. Epub 2018 Sep 6. PMID: 30297010; PMCID: PMC6203442.

Department of Microbiology, Dr D Y Patil Medical College, Hospital & Research Centre, Maharashtra 411018 ¹MBBS, Postgraduate Resident ²PhD Scholar ³MD (Microbiology), Senior Resident ⁴MSc Student

⁵MBBS, Postgraduate Resident and Corresponding Author


(BR) Brief Report, (Commt) Commentary, (C) Correspondence, (CR) Case Report,
 (CS) Case Series, (DC) Drug Corner, (Ed) Editorial, (OA) Original Article,
 (PCME) Pictorial CME, (RA) Review Article, (SA) Special Article,
 (S Comm) Short Communication

SUBJECT INDEX

Α

- Abnormal Haemoglobinopathy in Tribal Population of Purulia A Clinico-epidemiological Study of Patients of, **Bera S, Biswas T, Majumdar A**, (OA), **122(9)**: 24-7.
- Acromegaly Atypical Presentation in a Lady with: A Case Report, Purkayastha A, Bhattacharjee P, Khakholary M, (CR), 122(11): 61-5.
- Active Phase of Labour Routine Oxytocin Infusion versus Discontinuation during: Does it Make a Difference in Outcome — A Prospective Longitudinal Study. **Chakraborty B, Basu M, Debangshi M, Mondal PK,** (OA), **122(5)**: 46-9.
- Acute Coronary Syndrome (ACS) in Only Armed Forces Tertiary Cardiac Care Centre in Eastern India during COVID-19 Pandemic A Study to Evaluate Time Delays and Management Strategy in Patients Presenting with, **Sharma M**, (OA), **122(9)**: 21-3.
- Acute Coronary Syndrome and their Impact in Prognostication High Sensitive CRP and Platelet Volume Indices in, **Trivedi AS**, **Chakrabarti D, Khasnabish S, Bhattacharyya AK**, (OA), **122(4)**: 38-42.
- Acute Iliofemoral Deep Venous Thrombosis A Research Protocol on Efficacious Evaluation of Thrombus Recanalization Techniques Employing Pigtail Catheter *versus* Spray Catheter for Vessel Patency in — A Randomized Controlled Trial, Arora SP, Mishra GV, Banode P, Dass A, Banode P, Naqvi WM, Rathi RK, (S Comm), 122(11): 66-9.
- Acute Ischemic Stroke Patients Neutrophil to Lymphocyte Ratio (NLR) as an Inflammatory Marker in Predicting Severity in, Sharma D, Nagar GL, Fageria N, Mangal AK, Kumar P, (OA), 122(12): 40-3.
- Acute Myocardial Infarction Comparative Analysis of Sensitivity and Specificity of Creatinine Kinase MB and Cardiac Troponin I for the Diagnosis of, **Purandare NR, Goyal A, Shinde S, Jain S**, (OA), **122(12)**: 47-51.
- Acute ST Elevation Myocardial Infarction Patients Association Between T p-e / QT Ratio In 12 Lead ECG and Major Adverse Cardiac Events during Hospital Stay among, **Patel AS, Bavaria DR**, (OA), **122(10)**: 33-8.
- Adenocarcinoma of Cervix Uncommon Presentation of, Soni NR, Gupta P. (CR), 122(1): 54-5.
- Adrenal Crisis Impending in Exogenous Cushing Following Cam Use, Mandal M, Chanda A, Roy N, (C), 122(7): 85.

- Aeroallergens in Patients of Bronchial Asthma and Allergic Rhinitis Reporting to a Tertiary Care Centre in North India To Determine the Prevalence Pattern of, Chaudhary K, Sen MK, Swati A, Sachdeva R, Kumar A, (OA), 122(8): 54-7.
- Alcoholic Liver Disease (ALD) Patients and to Correlate MPV to Platelet Count Ratio with Child Pugh Score to Predict the Severity To Study Platelet Indices in, **Sharma D, Mangal AK**, **Fageria N, Kumar P, Nagar GL**, (OA), **122(2)**: 34-8.
- Allergen Immunotherapy on Medication Score and Medicine Usage in Patients with Nasobronchial Allergy Impact of : A Single Centre, Prospective Observational Study in a Specialty Clinic in Eastern India, Samajdar SS, Mukherjee S, Moitra S, Tripathi SK, (OA), 122(11): 20-5.
- An Enigmatic Disease: Endometriosis, Banerjee S, Mitra S, (Ed), 122(4): 12-4.
- Anaemia Autoimmune Haemolytic Approach in a Tertiary Care Centre, Dikshit N, Basu A, Dikshit NA, Prasad GH, (OA), 122(9): 33-9.
- Ancient Vaidyas to Modern Doctors— Unsung Heroes, Saboo B, Parikh R, Gupta A, Vora A, Nayak A, Amin B, Nagar J, Pal J, Joshi S, Bhandari S, Shah V, (SA), 122(5): 71-2.
- Anti-retroviral Therapy Registered at a Nodal Centre A Cross Sectional Study on Factors Associated with Patients being MIS-3 on — Goa Medical College, Shah HK, Dessai BPS, Bandodkar K, Talaulikar DMS, Khandeparkar A, Cacodcar J, Sawant G, Umraskar L, Galyekar S, Bhagat S, (OA), 122(8): 63-9.
- Apla Syndrome and Auto-immune Haemolytic Anaemia in a Patient of Hansen's Disease Neuropsychiatric Lupus with — A Diagnostic and Therapeutic Challenge, **Chakraborty S, Das P, Karan K, Sharma J**, (CR), **122(12)**: 69-71.
- Aplasia Cervico-vaginal : A Rare Anomaly with Review of Literature, Potdukhe PS, Dhok AP, (CR), 122(2): 61-2.
- Are Deep Learning Algorithms Changing the Landscape of Al-Assisted LV-GLS Analysis in Cardiology, Offering Hope for Early Disease Detection? Gandhi MA, Jadhav AK, (C), 122(3): 81.
- Arterial Blood Gas Sampling Technique in Emergency Department Settings Investigating the Multifaceted Aspects that Affect Interns' Competence and Performance in, Vijay KSS, Shekhli S, Jose A, (OA), 122(2): 45-8.
- Artificial Intelligence in Diagnostic Microbiology Exploring the Potential of, **Pandya YG, Pandya A**, (C), **122(1)**: 61.

Artificial Intelligence in Shaping the Future of Medicine The Pivotal Role of. **Banerjee S, Samajdar SS**, (Ed), **122(3)**: 14-9.

Artificial Kidneys Evolution of, Banerjee S, (Ed), 122(9): 11-2.

- Association between Sleep Quality and Different Aspects of Memory along with Assessment of Post Exercise and Post Meditation Effects, **Deshmukh AD, Shinde SA, Phalak PJ**, (C), **122(3)**: 79.
- Atresias Arising in Watershed Areas Congenital Neonatal Colonic of the Colonic Blood Supply, Patel RV, Trambadia RA, Anthony FM, Chhaniara RA, Govani DR, Patel RR, Correia RC, (CS), 122(2): 55-60.
- Atypical Lymphocytes, Large Immature Cells, and Rising Hematocrit in Predicting Dengue Fever Severity The Clinical Utility of, Singh S, Srivastava MK, (C), 122(1): 60.
- Auto-rickshaw Drivers in Belagavi City, Karnataka Tobacco use among, Patil AP, Yogeshkumar S, (OA), 122(11): 26-31.

В

- Benign Lung Diseases at Tertiary Health Care Hospital in South Gujarat Diagnostic Yield of Bronchoscopy in, Gengadiya JK, Tamakuwala GP, Vadgama PK, Mangrola JV, Khatri GP, (OA), 122(12): 19-22.
- Biochemistry in Every Square Solve the Mystery of, Pratap A, Phalak PJ,(C), 122(8): 73-4.
- Blood Culture by an Automated System Assessment of Time To Positivity (TTP) and Loading Delay Influencing the Positivity of, **Patel N, Patel R, Patel R**, (OA), **122(7)**: 37-42.
- Breastfeeding in the Rural Field Practice Area in Tamil Nadu Knowledge, Attitude and Practice on, **Dhanaseakaran S**, Rajavelu S, Sree M. (OA), **122(4)**: 31-7.
- Bronchial Asthma and COPD Assessment of the Level of Knowledge and Skills Pertaining to Self-management among Adult Patients of, **Pal A, Biswas B, Shit P, Samajdar SS, Sen S, Tripathi SK**, (OA), **122(7)**:70-5.
- Broncho Alveolar Lavage Fluid (BAL) in Patients with Lower Respiratory Tract Infections (LRTIs) Bacteriological Profile and Antibiotic Susceptibility Patterns of Clinical Isolates of, Panigrahy S, Kumari PHP, Samatha P, (OA), 122(4): 56-60.
- Broncho Alveolar Lavage Fluid (BALF) in Patients with Lower Respiratory Tract Infections (LRTIs) Bacteriological Profile and Antibiotic Susceptibility Patterns of Clinical Isolates of, Panigrahy S, Kumari PHP, Samatha P, (OA), 122(4): 56-60.
- Bruxism in Children and the Psychosocial Status of their Parents The Relationship between, **Khataminia M**, **Meshki R**, **Araban M**, **Faratseh M**, (OA), **122(10)**: 21-5.

С

- Cancer Impact of Psychological Intervention on the Quality of Life of Primary Care Givers of Patients with : Preliminary Report in City based Cancer Hospital, India, Gangopadhyay S, Datta A, Nandy K, Dhar H, Jayaseelan P, Agarwal P, (OA), 122(8): 29-35.
- Cancer Patients Attending a Tertiary Cancer Hospital An Observational Study to Identify Socio-demographic Factors on Primary Caregivers' Quality of Life of, Gangapadhyay S, Datta
 A, Nandy K, Dhar H, Datta S, Agarwal P, Mondal A, Mondal PK, Das S, (OA), 122(3): 24-9.
- Carbapenem Resistant Gram-negative Bacteria with Phenotypic Detection of Metallo-beta-lactamase Prevalence and Antibiogram of, **Babu A, Patil V, Bhatawadekar SM, Jadhav P**, (OA), **122(9)**: 40-3.
- Carcinoma Base of Penis A Rare Case of: Evaluation on Ultrasound

Color Doppler, Computed Tomography and Magnetic Resonance Imaging, **Bahurupe SS, Phatak SV, Dhok A**, (CR), **122(3)**: 66-8.

- Cardiac Surgery Intravesical Knotting of an Infant Feeding Tube in a Child who Underwent, **Kurien JG**, (C), **122(10)**: 80.
- Cardiomyopathy Infiltrative Causing Heart Failure with Compromised Ejection Fraction (HFrEF) : An entity to look out for, **Owuor H**, **Gikonyo A, Ponoth PM, Gikonyo D**, (CR), **122(7)**: 81-4.
- Chronic Hepatitis B in Eastern India A Study on Response of Tenofovir in, Saha M, Sarkar D, Mondal SS, Ray A, (OA), 122(9): 17-20.
- Chronic Kidney Disease Cardiac Function Evaluation in Different Stages of, Dutta N, Sarkar D, Sinha PK, (OA), 122(9): 48-53.
- Chronic Kidney Disease Prevalence of Depression in: An Observational Study, **Shekhli S, Allichandi R, Herakall M**, (OA), **122(1)**: 29-31.
- Clavien-Dindo Classification of Surgical Complications to Emergency Laparotomies A Non-randomized Retrospective Study of Applicability of, **Singh A, Atal R, Choudhary R, Singh Parihar U, Palsania R**, (OA), **122(6)**: 36-42.
- Clinical Competence Assessment Evolution of, Bhat VG, Rangaswamy DR, Kamble N, (C), 122(4): 86.
- Compulsory Online Education Adolescent Health and Screen Time during a Period of : A Cross Sectional Study from Eastern India, Siddhanta S, Bandyopadhyay D, Bhowmick S, (OA), 122(3): 43-9.
- Congenital Hypothyroidism Screening for Umbilical Cord Blood TSH a Useful Tool : A Single Centre Eight Year Experience, Bhatia R, Bhatia G, Rajwaniya D, Mishra P, Kaur J, (OA), 122(7): 28-31.
- Conjunctivitis Current Epidemic in India A Tale of Two Bugs, Das NK, Chate SS, Kulkarni SS, Yadav GE, Watwe SA, Chavan S, (BR), 122(10): 79.
- COPD Patients Attending a Tertiary Care Hospital Association between BODE Index and Visceral Fat among — A Cross Sectional Study, **Swamy T, Anitha OR, Kiran N, Shivaprasad VR, Manoj R**, (OA), **122(7)**: 21-7.
- Coronary Artery Disease Assessing the Diagnostic Efficacy of Multidetector CT Angiography versus Digital Subtraction Angiography in : A Tertiary Care Centre Study, **Sekar A, Karthikeyan N, Amir AP, Vikram MA, Faizal A, Natarajan P**, (OA), **122(7)**: 58-63.
- COVID Associated Rhino-orbito- Cerebral Mucormycosis Epidemic Treated from a Tertiary Care Centre in Eastern India Experiences from First 100 Cases of: An Otorhinolaryngologist Prospective, Das SK, Chatterjee P, Chakraborty A, Sengupta A, Barman D, Mukhopadhyay D, Srivastava SR, Adhikari S, Biswas PS, Basu K, (OA), 122(5): 50-7.
- COVID-19 Pandemic A Comparative Study on Nebulised Levosalbutamol *versus* Adrenaline in Wheeze Associated Condition of Children Between 1 Month to 6 Months of Age Admitted in Paediatric Ward of a Tertiary Medical College during, **Das S, Sardar S, Basu R, Sinha Mahapatra TK**. (OA), **122(10)**: 55-9.
- COVID-19 Patients and its Association with Outcome Clinical Profile, Laboratory Parameters, Radiological Findings of — A Retrospective Observational Study, Meenakumari PB, Bhaskar A, Indu RM, Jose C, (RA), 122(11): 50-5.
- COVID-19 Patients in a Tertiary Care Hospital of India A Study of the Clinico-demographic Profile and Clinical Outcomes of, Nath R, Sutradhar D, Thakuria R, (OA), 122(12): 23-8.
- COVID-19 Patients in Relation to their Vaccination Status Mortality amongst, **Selvaraju B, Kumar MP, Lawrence P,** (OA), **122(5)**: 24-7.

Current Status & Future Prospects : Telemedicine, Khade (RA), 122(6): 55-8.

D

- Deep Vein Thrombosis Study of Prevalence of Vitamin B12 Deficiency and Hyperhomocysteinemia in Patients of, **Barot MV, Bagada JB, Makwana RA, Rathod GH**, (OA), **122(1)**: 13-6.
- Dengue Fever An Observational Study on Neurological Complications among Patients with, **Kumar P, Sambhaji KS**, (OA), **122(4)**: 51-5.
- Dengue Fever with Thrombocytopenia Ocular Manifestations of Patients with : Dengue Fever seen through the Eyes, Ali Khan N, Singh A, Yadav V, Kochar A, Kochar SK, Farooq S, (OA), 122(4): 61-4.
- Dengue In-patients in a Rural Based Tertiary Hospital in West Bengal A Cross-sectional Study - Are Dengue Patients Aware Enough? Thakur I, Majumdar S, Jatua SK, Makhal MK, Saha S. (OA), 122(2): 39-44.
- Dengue Severity in Children in a Tertiary Care Hospital A Prospective, Observational Study of Serum Triglyceride and Cholesterol Level as Markers of, **Bose SS, Panda S, Mahapatra MK, Das AK**, (OA), **122(5)**: 19-23.
- Depression and Anxiety among Elderly Patients Visiting Tertiary Care Hospital Prevalence of, Kour S, Singh M, Gupta P, Arora R, Gupta U, (OA), 122(12): 52-6.
- Dermatophytosis in Children in a Tertiary Health Care in Karnataka A Clinico-epidemiological Study of, **Rathod MP, Rajegowda HM, Madegowda SB, Kalegowda D**. (OA), **122(3)**: 20-3.
- Diabetes Care Revolutionizing with a Patient-Friendly GLP-1 Receptor Agonist : Oral Semaglutide. Banerjee S, Samajdar SS, (Ed), 122(10): 11-20.
- Diabetes Mellitus among Tuberculosis Patients Attending District Tuberculosis Centre, Vijayapur Levels and Risk Factors of, Yadavannavar MC, Pattankar TP, Kouttalmatt DP, (OA), 122(12): 29-34.
- Diabetic Foot Ulcer Management Effect of Papain Urea *versus* EUSOL in, Sharath R, Philip AJ, Rajkumar S, Rahul RCL, Chand A, Umaakanth SCS, (OA), 122(10): 64-9.
- Diabetic Old Man The Utility of MatriDerm (Skin Substitute) with Autologous Thin Split Skin Graft to Cover Deperiosted Exposed Tibia 6x4 cm Wound in, **Singh S**, (RA), **122(7)**: 76-8.
- Disability and Behavioral Disturbances of the Children with Intellectual Disability and the Coping Mechanisms of their Parents Extent of: An Observational Study from a Tertiary Care Hospital of Eastern India. **Patra P, Dan A, Sarkar S, Mukhopadhyay A, Sanyal D**, (OA), **122(12)**: 35-9.
- Does only Mupirocin enough for Nasal Decolonization in Methicillinresistant Staphylococcus Aureus Carriers or do we need more? Mukhida S, Datta A, Edavaloth P, Pawar V, Mukherjee S, (C), 122(12): 72.

E

- Electrothermal Coupling Bowel Injury Indirect A Rare Complication of Laparoscopy, **Gawankar DS, Sarang B, Ruke M, Bulusu S**, (CR), **122(6)**: 63-4.
- Endocarditis with Pulmonary Septic Emboli Presenting as PUO in a Young Adult Female Managed Successfully with Surgical Intervention Isolated Native Tricuspid Valve, Jadhav AK, Khan AK, Nalawade DD, (CR), 122(10): 76-8.
- Endometrial Stromal Sarcomas A Case Series of : A Relook at a rare kind of Uterine Malignancy, Bandyopadhyay P, Ray

Chaudhuri G, Bandyopadhyay A, Ghosh Sengupta S, (CS), 122(11): 56-60.

- Endometrial Thickness versus Endometrial Pattern on Implantation in North Indian Females To Study the Correlation of : A Retrospective Observational Study, **Singh P, Singh RK, Rastogi R, Gandhi K**, (OA), **122(3)**: 30-2.
- Endoscopic Variceal Band Ligation and Assessing Utility of Dedicated Score (BICAP Score) to Identify High Risk Groups Risk Factors for Early Re-bleed following, **Unnikrishnan RK**, **Shafique A, Rejoice P, Kumar EK, Durai G, Kishore KR**, (OA), **122(8)**: 36-9.
- ENT Infections Exploring the Efficacy and Safety of Lincomycin 1000 mg SR Tablets in : A Clinical Investigation, **Desai A, Das P, Nair S, Sakpal A**, (DC), **122(11)**: 76-80.
- Epilepsy A Cross Sectional Study of Socio-cultural and Clinical Determinants of Health-related Quality of Life in Indian Patients with, Asole D, Bharote H, Thakre M. (OA), 122(8): 43-6.

F

- Factors Affecting Patient Attitude towards Private Clinical Establishments in India An Exploratory Study on, Sengupta M, Baishya K, Roy A, Chakrabarti S, Mukhopadhyay I, (OA), 122(4): 43-50.
- Faculty Development Programme for the 'Tutors & Residents' in a Medical College of Kolkata Introduction of a 'Capsule-course' in, Das Sarkar M, Damke S, Ray K, Roy H, (OA), 122(3): 33-8.
- Fetal Kidney Length in Estimation of Gestational Age Role of: Second Trimester versus Third Trimester, Luther N, (OA), 122(6): 17-20.
- Flagellate Pigmentation Following Chemotherapy An Unusual Case of, Nandy P, Rudra O, Chatterjee G, Mandal A, (C), 122(11): 82.
- Fungal Sinus Diseases with Maxillary Bone Erosion Invasive : Is Aggressive Maxillectomy Justified in all Cases? Sethuraman S, Adithya S. (OA), 122(4): 65-9.

G

- GIT Diseases An Update on Oral & Systemic Manifestations of: A Narrative Review, Vohra P, (RA), 122(4): 70-2.
- Growing Threat to Public Health : Carbapenem-Resistant Acinetobacter baumannii — A Review of Current Understanding and Future Directions, **Pate BS, Meshram S, Mudey GA, Shaw D**, (RA), **122(4)**: 73-7.

Н

- Handson Workshop in AETCOM Modules for Faculty in a Teaching Medical College Evaluation of, Gandham P, Goka S, Arcot R, (OA), 122(6): 15-6.
- Headache Disorders by Indian Medical Association Headache Algorithm : A Guide to Differential Diagnosis and Management of, Ravishankar K, Banerjee TK, Srinivasan AV, Patel DK, Jayakrishnan AV, Lele J, Shah B, Shah R, (Commt), 122(11): 71-5.
- Health in Changing Scenario Mental : Present Status and Future Direction, Banerjee S, Bhattacharyya R, (Ed), 122(8): 19-23.
- Healthcare Professionals Working in a COVID Hospital in West Bengal during COVID-19 Pandemic Assessment of Perceived Stress in, Halder S, Seikh P, Sarkar A, Chatterjee S, Mallick AK. (OA), 122(10): 46-8.
- Healthcare Reforms in India Challenges in, **Gupta VK**, (C), **122(4)**: 85.
- Heinous Offences with Demographic, Socio-economic Factors and

Personality Traits among the Children in Conflict with Law Staying in Juvenile Justice Homes A Study on Association of, **Chakrovarty A, Nath S, Guha Ray T, Batabyal S**, (OA), **122(4)**: 26-30.

- Hematological Parameters Including Novel Research Parameters in Population of Eastern India Biological Reference Intervals for, **Sengupta A, Pal S**, (OA), **122(5)**: 35-9.
- Hematoma in a Newborn Cutaneous Venous Malformation Masquerading as, **Khare C, Kumar A**, (CR), **122(4)**: 84.
- Hemodialysis Study of Assessment of Change in Lipid Profile Pattern in Patients on, **Deba Z, Jambale TA, Kalasker PS, Jaweed SA**, (OA), **122(7)**: 55-7.
- Hepatitis and Community : A Roadmap for 2030, Verma P, Johnson S, Palal D, Mukhida S. (C), 122(9): 77.
- HIV Associated Ophthalmic Diseases among the Patients Enrolled for Antiretroviral Therapy and its Correlation with CD4-T Cell Count Current Spectrum of — A Cross Sectional Study, **Biswas U**, **Ray Chaudhuri P, Ghosh S, Dey AK, Ray S, Chakraborty M**, (OA), **122(8)**: 47-53.
- Human Health in Northern Karnataka Evaluating Farmers' knowledge, Attitude and Practice Regarding Pesticide Use and Its Impact on - A Cross Sectional study, Nagendra A. Patil SS, (OA), 122(12): 57-60.
- Humerus Shaft Fracture with Combination Screw Prospective Study for Anterior Bridge Plate (ABP) for: Innovative SPV Technique, Patel SD, Desai AKA, Patel AA, (OA), 122(7): 64-9.
- Hypertension Uncontrolled and Resistant : A major challenges for Clinicians, Trailokya AA, Shirsat A, (C), 122(2): 67-8.
- Hypothermia in Newborn in Post Neonatal Ward and Factors Contributing It A Study of Occurrence of, **Dodiya AJ, Kakkad KM, Prajapati VN, Parmar H**, (OA), **122(2)**: 52-4.

- Infections in Paediatrics A Cornerstone Antibiotic for Treatment of: Amoxicillin / Clavulanate, Neha N, Kotak B, Ghosh KU, Shankar AVR, Veligandla KC, Rathod R, (RA), 122(9): 54-61.
- Infections in Post Renal Transplant Recipients in a Tertiary Care Hospital in Chennai A Prospective Observational Follow-up Study of, Varman M, Shanmugam B, Sharmili S, (OA), 122(1): 35-41.
- Infertile Females with Special Reference to Radiological Observations, Laparoscopic Findings and Endometrial Histology A Clinico-Anatomical Study on Primary and Secondary, Goswami N, Chakraborty N, Kundu D, Chakraborty S, Bhowmick S, (OA), 122(1): 32-4.
- Insulin A 100 Years Journey of, Chaurasia RC, (RA), 122(3): 58-62.
- Insulin Therapy Weekly: Where Do We Stand? Banerjee S, Raychaudhuri M. (Ed), 122(6): 12-4.
- Invasive Breast Carcinoma as an Independent Prognostic Factor in Comparison with Already Established Ones Microvascular Density of: A Study from Northeast India, **Devi J, Suhana A**, (OA), **122(12)**: 44-6.
- Is the Herbal Medicine in the Market Really Herbal? An opinion on Malpractices, Bhavsar HM, Kumar L, Mahashabde M, Bhimani Y, Mukhida S, (C), 122(10): 81.
- IVF Cycles Directly Correlates with the Dynamically Changing Endometrial Thickness Endometrial Perfusion on day of hCG Trigger in : A Pilot Study Examining Potential Implications for Endometrial Receptivity and the Development of an Integrated Model to Assess Endometrial Function, Ghosh Dastidar B, Ghosh Dastidar S, Majumdar J, Chakraborty C, Ghosh

Dastidar K, (OA), 122(2): 30-3.

IVF Pregnancies Postpartum Hemorrhage in — A Case Series, Basarakod SS, Halemani C, Rajashree SP, Chaitra N, (CS), 122(3): 63-5.

J

Jugular Tubercle in Dry Human Skull Osseomorphometric Study of, Biswas A, Pal A, Bhar A, (OA), 122(9): 44-7.

Κ

- Klebsiella Sepsis among Neonates Admitted in NICU at a Tertiary Care Centre of Western India Outcome of MDR — A Retrospective Study, **Das SK, Prajapati N**. (OA), **122(9)**: 28-32.
- Koch's Abdomen Diagnostic Peritoneal Biopsy in: A Case Report and Review of Literature, Kattamreddy AR, Kalyankar PP, Singh Bhalla R, Gupta N, (CR), 122(8): 70-2.

L

- Lesions of Liver among Patients Presenting in a Tertiary Care Center in India Etiologies of Space Occupying, Biswas T, Naskar A, Halder P, Mallik S, Mondal A, Saha B, (OA), 122(1): 42-7.
- Lifestyle Factors on the Health Among Students from Different Professional Backgrounds Comparitive Study of the Impact of, Akhil R, Jayaram V, Srivastava K, Johnson S, Mangala B, Vajjala SM ,(OA), 122(11): 38-44.
- Liver Abscess Cases from A Tertiary Care Hospital, Kolkata A Study on Risk Factor Assessment, Clinical Profile and Outcome Analysis of, **Modak D, Chatterjee R**, (OA), **122(6)**: 47-51.
- Lung Cancer Patients Psychological Distress and Health-related Quality of Life in, Kumar A, Verma AK, Bajpai J, Pandey AK, Kant S, Kar SK, Verma SK, Kushwaha RAS, Kumar S, Garg R, Srivastava A, Bajaj DK, Kumar A, Tripathi P, (OA), 122(7): 47-51.

Μ

- Malaria Parasite in Vivax Malaria Patients after Completion of Antimalaria Therapy in a Tertiary Care Hospital of Kolkata Presence of : A Real World Observational Study, Mukherjee K, Ghosh UC, Talukdar A, Ghosh A, Samajdar SS, (OA), 122(1): 26-8.
- Mastalgia Effectiveness of Evening Primrose Oil (EPO) versus Tamoxifen in the Treatment of, **Chrungoo I, Bhardwaj R**, (OA), **122(3)**: 39-42.
- Medical Educators Regarding the Integration of Standardized Teaching-learning Modules for Training Communication Skills in Medical Undergraduate Students Perceptions of, Ganganahalli PS, Udgiri R (OA), 122(4): 15-8.
- Medical Negligence, Banerjee S, (Ed), 122(2): 11-4.
- Moderately Severe COVID-19 Cases Use of Remdesivir as an Investigational Therapy in, **Talukdar AJ, Das RP, Hazarika B, Saikia P, Dutta S, Baishya AC**, (OA), **122(2)**: 15-20.
- Medical Sciences Artificial Intelligence In, Banerjee S, Banerjee C, (Ed), 122(11): 11-4.
- Medicine A Year of Transformative Strides in : 2024, Banerjee S, Samajdar SS, (Ed), 122(12): 11-8.
- Metabolic Dysfunction-associated Steatotic Liver Disease in India and the way forward Burden of, **Bandyopadhyay S**, (RA), **122(11)**: 45-9.
- Methicillin-resistant Staphylococcus aureus (MRSA) among Health Care Workers Working in Intensive and Critical Care Units at a Teaching Hospital in North India Nasal Carriage of, **Kilikdar M, Saha N, Mohan KR, Rehman Z**, (OA), **122(1)**: 21-5.

- Microfilaria at Unusual Sites with Varied Clinical Presentation Incidental Cytological Findings of : A Retrospective Study, **Sahu BK**, **Nayak S**, **Panigrahi C**. (OA), **122(2)**: 26-9.
- MIS-C Temporally Associated with COVID-19 Clinicoepidemiological Profile of - A Hospital Based Retrospective, Cross Sectional, Observational Study. What is new? Chatterjee A, Kundu TK, Bera M, Chowdhury A, Khan I, (OA), 122(2): 21-5.

N

- Non-motor Manifestations in Young Onset Parkinsons Disease (YOPD) in Eastern Indian Population A Study on, Yadav PK, Samsudheen KK, (OA), 122(7): 43-6.
- Non-traumatic Brown-Sequard Syndrome with Rare and Diverse Etiologies, A Case Series of, **Asole D, Bharote H, Thakre M**, (CS), **122(5)**: 58-61.
- Nuclear Scan A Reappraisal with Special Reference to : Elephantiasis, **Paul R**, (PCME), **122(11)**: 70.

0

- Obstetrics Hysterectomy Emergency : A Retrospective Study in Tertiary Care Hospital in Western India, Panchal PP, Maheta SD, Raval BM, Mehta AV, Shah SR, Shroff DJ, Shah RM, (OA), 122(7): 17-20.
- Operated by Mini Open Rotator Cuff Repair in Tertiary Care Hospital of Ahmedabad City Functional Outcome of Patients, **Mistry J**, **Patel TR, Patel PM**, (OA), **122(7)**: 52-4.
- Oral Lincomycin in the Treatment of Upper Respiratory Tract Infection Open-label, Prospective, Interventional Clinical Study to Evaluate the Efficacy and Safety of, **Desai A, Das P, Nair S, Parab S**, (DC), **122(9)**: 73-6.
- Osteopetrosis Mandibular Osteomyelitis Secondary to : A Management Conundrum, Uddin I, Dey S, Sikarwar S, Ganguly S, Anwar AA, (CR), 122(4): 80-3.
- Ovarian Cysts in Infants Auto-amputated : Current Status, Sinha CK, Anbarasan R, Patel R, Joshi A, Cherian A, (CR), 122(5): 62-4.
- Overweight Normoglycemic Offspring of T2DM Parents and Nondiabetic Parents Comparison of Effect of Exercise on Lipid Profiles of, **Taiwo EO**, (OA), **122(11)**: 15-9.
- Ovulation Monitoring of By Adopting "Dutta's New Scoring" Technique & Pregnancy Outcome, Dutta DK, Dutta I, Banerjee Dutta R, (OA), 122(10): 29-32.

Ρ

- Pan facial Fracture Interesting Case of, Ramesh XRW, Emperor CE, Dharsini OTS, (CR), 122(9): 66-70.
- Parenteral Human Placental Extract and Oral Azithromycin Versus Oral Azithromycin Alone in Chronic and Recurrent Pelvic Inflammatory Disease Efficacy and Safety of: An Open-Label Randomized Controlled Trial in Indian Patients, Mahapatra PC, Pandey A, Mayadeo NM, Dwivedi JS, Vani I, Venkatesh V, Mitra M, (OA), 122(8): 58-62.
- Parkinson's Disease Defective Systemic Iron Metabolism in, Mahindrakar YS, Jadhav BH, Salunkhe SN, (OA), 122(6): 52-4.
- Parvovirus B19 in an Immunocompetent Patient Refractory Pure Red Cell Aplasia Secondary to : A Challenging Case Report, Bohare SM, Lohar R, Chaudhari AS, Shriwastav UK, Rijal P, Kant R, (CR), 122(1): 51-3.
- Peak Expiratory Flow Rate (PEFR) in Children Residing in Urban

Areas of Dhule District Evaluation of, Nikam KD, Hatkar NK, Ghate S, (OA), 122(10): 26-8.

- Peripartum Cardiomyopathy Multidisciplinary Management of Maternal Near Miss (MNM) due to — A Case Report, Chakrabarti Bhattacharyya N, Roy A, Ghosh Haldar R, Chatterjee S, Mandal D, Haque S, Bhattacharyya S Chakrabarti Bhattacharyya N, Roy A, Ghosh Haldar R, Chatterjee S, Mandal D, Haque S, Bhattacharyya S, (CR), 122(5): 65-7.
- PET Scan of Brain and Dementia, Paul R, (PCME), 122(5): 73.
- Pneumococcal Conjugate Vaccine among Medical and Nursing Students studying at Civil Hospital Campus, Ahmedabad A Comparative Study to Estimate Knowledge about, Parmar RK, Patel MK, Bhatt RJ, (OA), 122(5): 33-4.
- Post-COVID-19 Pulmonary Interstitial Pneumonitis with Methylene Blue Nebulization Therapy Resolution of — A Case Report, Golwalkar D, Pandya J, (CR), 122(9): 71-2.
- Preterm Birth in Indian Scenario Clinical Outcome of Arabin Cervical Pessary in Women at Risk of, Yadav P, Singh R, Agrawal R, Verma A, Malhotra N, Kumar N, (OA), 122(5): 40-5.
- Primary Headache Disorders in India Understanding the Changing Landscape of, **Banerjee S** (Ed), **122(5)**: 12-3.
- Prophylactic Retention Sutures on Closure of Laparotomy Wound in High-risk Patients in a Tertiary Care Hospital Effects of, **Dagra R**, **Mukherjee R**, **Das S**, **Roy A**, **Saha S**, (OA), **122(8)**: 40-2.
- Proton Pump Inhibitors at a Tertiary Care Hospital in South India Drug Utilization Pattern of, Shanmugapriya S, Saravanan T, Suriya Dharsini J, Shuruthi S, Saravanan A, (OA), 122(6): 21-7.
- Publication Spree in Medical Colleges. Where does this lead us in long term? Das NK, Chate SS, Watwe SA, Chavan S, Mukhida S, (C), 122(1): 60-1.

Q

Quality over Quantity in Health Care "Life isn't about quantity, it's about quality" – Malorie Blackman, Landge JA, (C), 122(6): 65-6

R

- Reinforced Training of BLS on the Ability to Retain the Imparted Knowledge and Skill amongst OT Personnel Including Nursing Staff and Technicians Evaluation of the Impact of : A Prospective Interventional Study. Acharya GR, Lagoo JY, Chate SU, (OA), 122(10): 39-45.
- Renal Agenesis Unilateral Müllerian Duct Fusion Anomaly Associated with, Vishwaprem RDR, Naveen D, Mallikarjunappa B, Gupta A. (CR), 122(7): 79-80.

REVIEW :

HIV-AIDS in India & Developing Countries, **Krishna YM**, **122(11)**: 81.

- Rheumatoid Arthritis with Waitlist Control Group A Comparative Study of Effect of Mindfulness-based Stress Reduction on Psychological Stress & Quality of Life in Patients of, **Sen A, Bandyopadhyay T, Bhattacharyya R, Saha PK, Das D** (OA), **122(5)**: 28-32.
- Rhino-Orbito-Cerebral Mucormycosis Post COVID-19 : Retrospective Clinical Observational Study & Analysis of the Patients Presenting in Kheda District, Gujarat, India, Prabhu S, Mehta R, Steward E, Doria G, Bavishi M, Masariya B, (OA), 122(5): 14-8.
- Rifampicin induced Thrombocytopenia : a rare complication, Seval M, Jain P, Meena V, Verma A, (C), 122(2): 69.

My Health, My Right. Banerjee S, (Ed), 122(7): 15-6.

S

- SARS-CoV-2 Infection A Cross Sectional Clinical Study to Evaluate the Pattern of Acute Hair Loss in Patients after, Barbhaya M, Modasiya A, Patel A, Bapat N, Patel K, (OA), 122(3): 50-3.
- SARS-CoV-2 Rapid Antigen Test as Compared to RT-PCR Test Sensitivity of, Mundhada SG, Deogade N, Tanna J, Deshmukh M, Mundhada AS, (OA), 122(7): 32-6.
- Seizures Disorder by Magnetic Resonance Imaging in a Tertiary Care Hospital Cross-sectional Observational Study to Determine Morphology of the Hippocampus in, Vishwaprem RDR, Khan Mohd SMS, Naveen D, Srinivasa BCR, Mallikarjunappa B, (OA), 122(6): 28-31.
- Self-directed Learning Compared to the Traditional Method of Learning for Undergraduate Medical Students Study of Effectiveness of, Mohapatra D, Sahoo PK, Meher A, Behera M, Mishra T. (OA), 122(10): 60-3.
- Should JN 1 Trigger an Alarm? Kamble N, Kavatagi K, Rangaswamy DR, (C), 122(6): 65.
- Simplifying Day-to-Day Practice with Al, Rangaswamy DR, Kamble N, (C), 122(2): 68-9.
- Skin and Soft Tissue Infections and Surgical Site Infections Assessing the Therapeutic Effectiveness and Tolerability of Lincomycin Injectables in: A Comprehensive Real-World Evidence Study, Ruke M, Desai A, Anand S, Nair S, (DC), 122(1): 56-9.
- Skin Sore Unraveling the Mystery of Chigger Fever or Cigarette Burn, Rajagopalan R, (C), 122(5): 74.
- Smartphone Exposure among Children <5 Years of Age Attending Out-patient Department of a Tertiary Care Hospital Pattern of: A Cross-sectional Study, Das A, Biswas N, Agarwala B, Mandal PK, (OA), 122(2): 49-51.
- Snake Bite Envenomation at a Tertiary Care Centre in Eastern Rajasthan Clinico-epidemiological Profile of : A Longitudinal Prospective Study, Jain PK, Goswami H, Bhat PK, Seval M, (OA), 122(3): 54-7.
- Spermatic Cord Hydrocele in 2-year-old Child A Rare Case of Encysted Type of, Gund SV, Dhok AP, Phatak SV, Onkar PM, Mitra K, Trimukhe DM, (CR), 122(4): 78-9.
- Spinal Cord Disc Diseases in Bankura Sammilani Medical College and Hospital A Comparative Study on the Efficacy of Pregabalin Over Gabapentin in Controlling Neuropathic Pain due to, Choudhury U, Banerjee C, Sural R, (OA), 122(1): 17-20.
- Steatosis Cardiac An Emerging Entity, Bagchi AS, Bagchi A, (RA), 122(12): 61-5.
- Successful Use of Total Parenteral Nutrition in Patients with Paraquat Poisoning, **Biswas U**, **Mukhopadhyay A**, **Datta S**,: (C), **122(3)**: 79-80.

Surgery: Robots in, Banerjee S, (Ed), 122(1): 11-2.

- Surgical Site Infection (SSI) and Skin & Soft Tissue Infection (SSTI) A Real-World Evidence Study on Effectiveness and Tolerability of Topical Lincomycin in the Treatment of, Ruke M, Desai A, Anand S, Nair S, (DC), 122(2): 63-6.
- Symptomatic Direct Inguinal Hernia in an Apparently Healthy Boy Laparoscopic Repair of, **Patel RV, Correia RC**, (C), **122(10)**: 82.

Т

TB Households in Northern Karnataka A Cross-sectional Study Assessing Indoor Air Pollution Sources among, Ali SS, Patil SS, (OA), 122(9): 13-6.

- Telemedicine Services among the Health Care Providers and Beneficiaries in Purba Bardhaman District, West Bengal Perception and Constraints of : A Qualitative Study, **Sannigrahi S, Bandyopadhyay S, Thakur RP, Taraphdar P, Jha A, Mallick A**, (OA), **122(11)**: 32-7.
- Temporary Contraceptive Methods in Ahmedabad City, Gujarat, India An Assessment of Knowledge and Attitude of Beneficiaries Attending Tertiary Care Hospital regarding Temporary Family Planning Methods and to Determine their Experienced Side Effects of. Saini VK, Bhabhor MB, Panchal DM, Patel PP, Chaudhari VK, Dagra JJ. (OA), 122(8): 24-8.
- Testicular Perfusion after Laparoscopic Hernioplasty Evaluation of Alteration in : A Prospective Study Conducted at one of the Tertiary Care Centres of Western India, Shah SV, Chavada RR, Thakar PM, (OA), 122(10): 70-5.
- Therapeutics Reimagined Hypertension : Nebivolol and Telmisartan — A Contemporary Review, Kumbla DK, (DC), 122(3): 69-74.
- Thrombotic Events in Indian Patients Statin Intake and A Cross Sectional Study, Sharma A, Donepudi A, Padmaja M, (OA), 122(1): 48-50.
- Transient Ventricular Asystole following High Spinal Cord Injury Silent Beats : Spontaneous Recovery of, Malani SK, Nalawade DD, Jadhav A, Wadhokar P, (CR), 122(5): 68-70.
- Tubercular Lesions Spinal Intramedullary A Case Series of Clinicoradiological Manifestations, Khan I, Srivastava T, Meena S, (CS), 122(6): 59-62.
- Tuberculosis among adult Residents of Urban Field Practice Area of Shri B M Patil Medical College, Vijayapura Knowledge, Attitude and Practices Regarding, **Patil SD, Ganganahalli P, Yankanchi S.** (OA), **122(6)**: 43-6.
- Tuberculosis in Women Presenting with Infertility Peritoneal Fluid from Pouch of Douglas is not a Suitable Specimen for Molecular Testing in the Diagnosis of Female Genital, Mishra D, Turuk J, Palo S, Ray PK, Mishra B, Pati S, (OA), 122(10): 49-54.
- Type-2 Diabetes Mellitus Prevalence of Diabetic Kidney Disease and Its Associated Risk Factors in — A Tertiary Care Experience, Shah SJ, Shah JH, Prajapati P, (OA), 122(4): 19-25.
- Typhus Scrub A Case Series from Tertiary Care Hospital of Rural Bengal, Mahapatra U, Bhunia PK, Murmu BK, Kundu S, (CS), 122(9): 62-5.

U

- Umbilical Cord Blood Banking Truth Unveiled : ICMR Guidelines Debunks Common Misconceptions about, Kattamreddy AR, (C), 122(4): 85-6.
- Unilateral Parotitis and Encephalopathy Scrub Typhus with: A rare case report and literature review, Mahapatra U, Das Bakshi S, Mukhopadhyay S, Gayen BK,(CR), 122(12): 66-8.
- Urgent Need for Dedicated Clinical Pharmacology Departments and to Address Faculty Shortage in West Bengal, Samajdar SS, Tripath SK, (C), 122(8): 73.
- Urinary Bladder in Human Fetuses Histogenesis of, Devi YE, Singh NS, Singh CR, Cindy L, (OA), 122(6): 32-5.

V

Vitamin D3 Formulations Advancements in : A Review of UNS D3 Ultra Nano 60 Thousand, Reddy KJ, Reddy J, Reddy S, Desai A, Garg MR, (DC), 122(3): 75-8.

AUTHOR INDEX

Α

Acharya GR, Lagoo JY, Chate SU: Evaluation of the Impact of Reinforced Training of BLS on the Ability to Retain the Imparted Knowledge and Skill amongst OT Personnel Including Nursing Staff and Technicians : A Prospective Interventional Study. (OA), 122(10): 39-45.

Adhikari S: see Das SK.

Adithya S: see Sethuraman S.

Agarwal P: see Gangapadhyay S.

Agarwal P: see Gangopadhyay S.

Agarwala B: see Das A.

Agrawal R: see Yadav P.

- Akhil R, Jayaram V, Srivastava K, Johnson S, Mangala B, Vajjala
 SM: Lifestyle Factors on the Health Among Students from Different Professional Backgrounds, (OA), 122(11): 38-44.
- Ali Khan N, Singh A, Yadav V, Kochar A, Kochar SK, Farooq S: Dengue Fever seen through the Eyes : Ocular Manifestations of Patients with Dengue Fever with Thrombocytopenia, (OA), 122(4): 61-4.
- Ali SS, Patil SS: A Cross-sectional Study Assessing Indoor Air Pollution Sources among TB Households in Northern Karnataka, (OA), 122(9): 13-6.

Allichandi R: see Shekhli S.

Amin B: see Saboo B.

Amir AP: see Sekar A.

Anand S: see Ruke M.

Anand S: see Ruke M.

Anbarasan R: see Sinha CK.

Anitha OR: see Swamy T.

Anthony FM: see Patel RV.

Anwar AA: see Uddin I.

Araban M: see Khataminia M.

Arcot R: see Gandham P.

Arora R: see Kour S

- Arora SP, Mishra GV, Banode P, Dass A, Banode P, Naqvi WM, Rathi RK: A Research Protocol on Efficacious Evaluation of Thrombus Recanalization Techniques Employing Pigtail Catheter versus Spray Catheter for Vessel Patency in Acute Iliofemoral Deep Venous Thrombosis — A Randomized Controlled Trial, (S Comm), 122(11): 66-9.
- Asole D, Bharote H, Thakre M: A Case Series of Non-traumatic Brown-Sequard Syndrome with Rare and Diverse Etiologies, (CS), **122(5)**: 58-61.
- Asole D, Bharote H, Thakre M: A Cross Sectional Study of Sociocultural and Clinical Determinants of Health-related Quality of Life in Indian Patients with Epilepsy, (OA), 122(8): 43-6.
 Atal R: see Singh A.

В

Babu A, Patil V, Bhatawadekar SM, Jadhav P: Prevalence and Anti-biogram of Carbapenem Resistant Gram-negative Bacteria with Phenotypic Detection of Metallo-beta-lactamase, (OA), 122(9): 40-3.

Bagada JB: see Barot MV.

Bagchi A: see Bagchi AS.

- Bagchi AS, Bagchi A: Cardiac Steatosis An Emerging Entity, (RA), 122(12): 61-5.
- Bahurupe SS, Phatak SV, Dhok A: Carcinoma Base of Penis : Evaluation on Ultrasound Color Doppler, Computed Tomography and Magnetic Resonance Imaging, (CR), **122(3)**: 66-8.

Baishya AC: see Talukdar AJ.

Baishya K: see Sengupta M.

Bajaj DK: see Kumar A.

Bajpai J: see Kumar A.

- Bandodkar K: see Shah HK.
- Bandyopadhyay A: see Bandyopadhyay P.
- Bandyopadhyay D: see Siddhanta S.
- Bandyopadhyay P, Ray Chaudhuri G, Bandyopadhyay A, Ghosh Sengupta S: A Case Series of Endometrial Stromal Sarcomas: A Relook at a rare kind of Uterine Malignancy, (CS), **122(11)**: 56-60.
- Bandyopadhyay S: Burden of Metabolic Dysfunction-associated Steatotic Liver Disease in India and the way forward, (RA), 122(11): 45-9.

Bandyopadhyay S: see Sannigrahi S.

Bandyopadhyay T: see Sen A.

Banerjee C: see Banerjee S.

- Banerjee C: see Choudhury U.
- Banerjee Dutta R: see Dutta DK.
- Banerjee S, Banerjee C: Artificial Intelligence In Medical Sciences, (Ed), 122(11): 11-4.
- Banerjee S, Bhattacharyya R: Mental Health in Changing Scenario: Present Status and Future Direction, (Ed), **122(8)**: 19-23.
- Banerjee S, Mitra S: Endometriosis : An Enigmatic Disease, (Ed), 122(4): 12-4.
- Banerjee S, Raychaudhuri M: Weekly Insulin Therapy : Where Do We Stand? (Ed), 122(6): 12-4.
- Banerjee S, Samajdar SS : Oral Semaglutide : Revolutionizing Diabetes Care with a Patient-Friendly GLP-1 Receptor Agonist. (Ed), **122(10)**: 11-20.
- Banerjee S, Samajdar SS: 2024 : A Year of Transformative Strides in Medicine, Surgery and Public Health. (Ed), 122(12): 11-8.
- Banerjee S, Samajdar SS: The Pivotal Role of Artificial Intelligence in Shaping the Future of Medicine. (Ed), **122(3)**: 14-9.
- Banerjee S: Evolution of Artificial Kidneys, (Ed), 122(9): 11-2.
- Banerjee S: Medical Negligence. (Ed), 122(2): 11-4.

Banerjee S: My Health, My Right. (Ed), 122(7): 15-6.

- Banerjee S: Robots in Surgery. (Ed), 122(1): 11-2.
- Banerjee S: Understanding the Changing Landscape of Primary Headache Disorders in India, (Ed), **122(5)**: 12-3.
- Baneriee TK: see Ravishankar K.

Banode P: see Arora SP.

Banode P: see Arora SP.

Bapat N: see Barbhaya M.

- Barbhaya M, Modasiya A, Patel A, Bapat N, Patel K: A Cross Sectional Clinical Study to Evaluate the Pattern of Acute Hair Loss in Patients after SARS-CoV-2 Infection, (OA), 122(3): 50-3.
 Barman D: see Das SK
- Barot MV, Bagada JB, Makwana RA, Rathod GH: Study of Prevalence of Vitamin B12 Deficiency and Hyperhomocysteinemia in Patients of Deep Vein Thrombosis, (OA), 122(1): 13-6.
- Basarakod SS, Halemani C, Rajashree SP, Chaitra N: Postpartum Hemorrhage in IVF Pregnancies — A Case Series, (CS), **122(3)**: 63-5.
- Basu A: see Dikshit N.
- Basu K: see Das SK.

Basu M: see Chakraborty B.

Basu R: see Das S.

Batabyal S: see Chakrovarty A.

80

Bavaria DR: see Patel AS. Bavishi M: see Prabhu S. Behera M: see Mohapatra D. Bera M: see Chatteriee A. Bera S, Biswas T, Majumdar A: A Clinico-epidemiological Study of Patients of Abnormal Haemoglobinopathy in Tribal Population of Purulia, (OA), 122(9): 24-7. Bhabhor MB: see Saini VK. Bhagat S: see Shah HK. Bhandari S: see Saboo B Bhar A: see Biswas A. Bhardwaj R: see Chrungoo I. Bharote H: see Asole D. Bharote H: see Asole D. Bhaskar A: see Meenakumari PB. Bhat PK: see Jain PK. Bhat VG, Rangaswamy DR, Kamble N: Evolution of Clinical Competence Assessment, (C), 122(4): 86. Bhatawadekar SM: see Babu A. Bhatia G: see Bhatia R. Bhatia R, Bhatia G, Rajwaniya D, Mishra P, Kaur J: Screening for Congenital Hypothyroidism — Umbilical Cord Blood TSH a Useful Tool : A Single Centre Eight Year Experience, (OA),

122(7): 28-31.

Bhatt RJ: see Parmar RK.

- Bhattacharjee P: see Purkayastha A.
- Bhattacharyya AK: see Trivedi AS.
- Bhattacharyya R: see Banerjee S.
- Bhattacharyya R: see Sen A.,
- Bhattacharyya S: see Chakrabarti Bhattacharyya N.
- Bhavsar HM, Kumar L, Mahashabde M, Bhimani Y, Mukhida S: Is the Herbal Medicine in the Market Really Herbal? An opinion on Malpractices, (C), 122(10): 81.

Bhimani Y: see Bhavsar HM.

Bhowmick S: see Goswami N.

- Bhowmick S: see Siddhanta S.
- Bhunia PK: see Mahapatra U.
- Biswas A, Pal A, Bhar A: Osseomorphometric Study of Jugular Tubercle in Dry Human Skull, (OA), 122(9): 44-7.

Biswas B: see Pal A.

Biswas N: see Das A.

Biswas PS: see Das SK.

Biswas T, Naskar A, Halder P, Mallik S, Mondal A, Saha B: Etiologies of Space Occupying Lesions of Liver among Patients Presenting in a Tertiary Care Center in India, (OA), 122(1): 42-7

Biswas T: see Bera S.

- Biswas U, Mukhopadhyay A, Datta S: Successful Use of Total Parenteral Nutrition in Patients with Paraguat Poisoning, (C), 122(3): 79-80.
- Biswas U, Ray Chaudhuri P, Ghosh S, Dey AK, Ray S, Chakraborty M: Current Spectrum of HIV Associated Ophthalmic Diseases among the Patients Enrolled for Antiretroviral Therapy and its Correlation with CD4-T Cell Count A Cross Sectional Study, (OA), 122(8): 47-53.
- Bohare SM, Lohar R, Chaudhari AS, Shriwastav UK, Rijal P, Kant R: Refractory Pure Red Cell Aplasia Secondary to Parvovirus B19 in an Immunocompetent Patient : A Challenging Case Report, (CR), 122(1): 51-3
- Bose SS, Panda S, Mahapatra MK, Das AK: A Prospective, Observational Study of Serum Triglyceride and Cholesterol Level as Markers of Dengue Severity in Children in a Tertiary Care Hospital, (OA), 122(5): 19-23.

Bulusu S: see Gawankar DS.

С

Cacodcar J: see Shah HK.

Chaitra N: see Basarakod SS.

Chakrabarti Bhattacharyya N, Roy A, Ghosh Haldar R, Chatterjee S, Mandal D, Haque S, Bhattacharyya S: Multidisciplinary Management of Maternal Near Miss (MNM) due to Peripartum Cardiomyopathy - A Case Report, (CR), 122(5): 65-7.

Chakrabarti D: see Trivedi AS.

Chakrabarti S: see Sengupta M. Chakraborty A: see Das SK.

- Chakraborty B, Basu M, Debangshi M, Mondal PK: Routine Oxytocin Infusion versus Discontinuation during Active Phase of Labour : Does it Make a Difference in Outcome - A Prospective Longitudinal Study. (OA), 122(5): 46-9.
- Chakraborty C: see Ghosh Dastidar B.
- Chakraborty M: see Biswas U.
- Chakraborty N: see Goswami N.
- Chakraborty S, Das P, Karan K, Sharma J: Neuropsychiatric Lupus with Apla Syndrome and Auto-immune Haemolytic Anaemia in a Patient of Hansen's Disease - A Diagnostic and Therapeutic Challenge, (CR), 122(12): 69-71.
- Chakraborty S: see Goswami N.
- Chakrovarty A, Nath S, Guha Ray T, Batabyal S: A Study on Association of Heinous Offences with Demographic, Socioeconomic Factors and Personality Traits among the Children in Conflict with Law Staying in Juvenile Justice Homes, (OA), 122(4): 26-30.
- Chand A: see Sharath R.

Chanda A: see Mandal M.

Chate SS: see Das NK.

Chate SS: see Das NK.

- Chate SU: see Acharya GR.
- Chatterjee A, Kundu TK, Bera M, Chowdhury A, Khan I: Clinicoepidemiological Profile of MIS-C Temporally Associated with COVID-19 - A Hospital Based Retrospective, Cross Sectional, Observational Study. What is new? (OA), 122(2): 21-5.

Chatterjee G: see Nandy P.

Chatterjee P: see Das SK.

- Chatterjee R: see Modak D.
- Chatterjee S: see Chakrabarti Bhattacharyya N.

Chatterjee S: see Halder S.

Chaudhari AS: see Bohare SM.

Chaudhari VK: see Saini VK.

Chaudhary K, Sen MK, Swati A, Sachdeva R, Kumar A: To Determine the Prevalence Pattern of Aeroallergens in Patients of Bronchial Asthma and Allergic Rhinitis Reporting to a Tertiary Care Centre in North India, (OA), 122(8): 54-7.

Chaurasia RC: A 100 Years Journey of Insulin, (RA), 122(3): 58-62.

Chavada RR: see Shah SV.

Chavan S: see Das NK.

Chavan S: see Das NK.

Cherian A: see Sinha CK

Chhaniara RA: see Patel RV.

Choudhary R: see Singh A.

Choudhury U, Banerjee C, Sural R: A Comparative Study on the Efficacy of Pregabalin Over Gabapentin in Controlling Neuropathic Pain due to Spinal Cord Disc Diseases in Bankura Sammilani Medical College and Hospital, (OA), 122(1): 17-20. Chowdhury A: see Chatterjee A.,

Chrungoo I, Bhardwaj R: Effectiveness of Evening Primrose Oil (EPO) versus Tamoxifen in the Treatment of Mastalgia, (OA), 122(3): 39-42.

Cindy L: see Devi YE.

Correia RC: see Patel RV. Correia RC: see Patel RV.

D

Dagra JJ: see Saini VK.

Dagra R, Mukherjee R, Das S, Roy A, Saha S: Effects of Prophylactic Retention Sutures on Closure of Laparotomy Wound in High-risk Patients in a Tertiary Care Hospital, (OA), 122(8): 40-2.

Damke S: see Das Sarkar M.

Dan A: see Patra P.

Das A, Biswas N, Agarwala B, Mandal PK: Pattern of Smartphone Exposure among Children <5 Years of Age Attending Out-patient Department of a Tertiary Care Hospital : A Cross-sectional Study, (OA), 122(2): 49-51.

Das AK: see, Bose SS.

Das Bakshi S: see Mahapatra U.

Das D: see Sen A.

- Das NK, Chate SS, Kulkarni SS, Yadav GE, Watwe SA, Chavan S: Current Conjunctivitis Epidemic in India — A Tale of Two Bugs, (BR), 122(10): 79.
- Das NK, Chate SS, Watwe SA, Chavan S, Mukhida S: Publication Spree in Medical Colleges. Where does this lead us in long term? (C), 122(1): 60-1.

Das P: see Chakraborty S.

Das P: see Desai A.

Das P: see Desai A.

- Das RP: see Talukdar AJ.
- Das S, Sardar S, Basu R, Sinha Mahapatra TK: A Comparative Study on Nebulised Levosalbutamol versus Adrenaline in Wheeze Associated Condition of Children Between 1 Month to 6 Months of Age Admitted in Paediatric Ward of a Tertiary Medical College during COVID-19 Pandemic, (OA), 122(10): 55-9.

Das S: see Dagra R.

- Das S: see Gangapadhyay S.
- Das Sarkar M, Damke S, Ray K, Roy H: Introduction of a 'Capsulecourse' in Faculty Development Programme for the 'Tutors & Residents' in a Medical College of Kolkata, (OA), 122(3): 33-8.
- Das SK, Chatterjee P, Chakraborty A, Sengupta A, Barman D, Mukhopadhyay D, Srivastava SR, Adhikari S, Biswas PS, Basu K: Experiences from First 100 Cases of COVID Associated Rhino-orbito- Cerebral Mucormycosis Epidemic Treated from a Tertiary Care Centre in Eastern India : An Otorhinolaryngologist Prospective, (OA), 122(5): 50-7.
- Das SK, Prajapati N: Outcome of MDR Klebsiella Sepsis among Neonates Admitted in NICU at a Tertiary Care Centre of Western India — A Retrospective Study, (OA), 122(9): 28-32.

Dass A: see Arora SP.

- Datta A: see Gangapadhyay S.
- Datta A: see Gangopadhyay S.

Datta A: see Mukhida S.

- Datta S: see Gangapadhyay S.
- Datta S:: see Biswas U
- Deba Z, Jambale TA, Kalasker PS, Jaweed SA: Study of Assessment of Change in Lipid Profile Pattern in Patients on Hemodialysis, (OA), 122(7): 55-7.

Debangshi M: see Chakraborty B.

Deogade N: see Mundhada SG.

Desai A, Das P, Nair S, Parab S: Open-label, Prospective, Interventional Clinical Study to Evaluate the Efficacy and Safety of Oral Lincomycin in the Treatment of Upper Respiratory Tract Infection, (DC), 122(9): 73-6.

Desai A, Das P, Nair S, Sakpal A: Exploring the Efficacy and Safety

of Lincomycin 1000 mg SR Tablets in ENT Infections : A Clinical Investigation, (DC), **122(11)**: 76-80.

Desai A: see Reddy KJ.

Desai A: see Ruke M

Desai A: see Ruke M.

Desai AKA: see Patel SD.

Deshmukh AD, Shinde SA,, Phalak PJ: Association between Sleep Quality and Different Aspects of Memory along with Assessment of Post Exercise and Post Meditation Effects, (C), 122(3): 79. Deshmukh M: see Mundhada SG.

Dessai BPS: see Shah HK.

- Devi J, Suhana A: Microvascular Density of Invasive Breast Carcinoma as an Independent Prognostic Factor in Comparison with Already Established Ones : A Study from Northeast India, (OA), 122(12): 44-6.
- Devi YE, Singh NS, Singh CR, Cindy L: Histogenesis of Urinary Bladder in Human Fetuses, (OA), **122(6)**: 32-5.
- Dey AK: see Biswas U.

Dey S: see Uddin I.

- Dhanaseakaran S, Rajavelu S, Sree M: Knowledge, Attitude and Practice on Breastfeeding in the Rural Field Practice Area in Tamil Nadu, (OA), **122(4)**: 31-7.
- Dhar H: see Gangapadhyay S.
- Dhar H: see Gangopadhyay S.

Dharsini OTS: see Ramesh XRW.

Dhok A: see Bahurupe SS.

Dhok AP: see Gund SV.

- Dhok AP: see Potdukhe PS.
- Dikshit N, Basu A, Dikshit NA, Prasad GH: Autoimmune Haemolytic Anaemia — Approach in a Tertiary Care Centre, (OA), 122(9): 33-9.
- Dikshit NA: see Dikshit N.
- Dodiya AJ, Kakkad KM, Prajapati VN, Parmar H: A Study of Occurrence of Hypothermia in Newborn in Post Neonatal Ward and Factors Contributing It, (OA), **122(2)**: 52-4.

Donepudi A: see Sharma A.

- Doria G: see Prabhu S.
- Durai G: see Unnikrishnan RK.
- Dutta DK, Dutta I, Banerjee Dutta R: Monitoring of Ovulation By Adopting "Dutta's New Scoring" Technique & Pregnancy Outcome, (OA), 122(10): 29-32.
- Dutta I: see Dutta DK.
- Dutta N, Sarkar D, Sinha PK: Cardiac Function Evaluation in Different Stages of Chronic Kidney Disease, (OA), 122(9): 48-53.

Dutta S: see Talukdar AJ.

Dwivedi JS: see Mahapatra PC.

E

Edavaloth P: see Mukhida S. Emperor CE: see Ramesh XRW.

F.

Fageria N: see Sharma D. Fageria N: see Sharma D. Faizal A: see Sekar A. Faratseh M: see Khataminia M. Faroog S: see Ali Khan N.

G

Galyekar S: see Shah HK.

Gandham P, Goka S, Arcot R: Evaluation of Handson Workshop in AETCOM Modules for Faculty in a Teaching Medical College, (OA), **122(6)**: 15-6.

Gandhi K: see Singh P.

- Gandhi MA, Jadhav AK: Are Deep Learning Algorithms Changing the Landscape of AI-Assisted LV-GLS Analysis in Cardiology, Offering Hope for Early Disease Detection? (C), 122(3): 81.
 Ganganahalli P: see Patil SD.
- Ganganahalli PS, Udgiri R: Perceptions of Medical Educators Regarding the Integration of Standardized Teaching-learning Modules for Training Communication Skills in Medical Undergraduate Students, (OA), **122(4)**: 15-8.
- Gangapadhyay S, Datta A, Nandy K, Dhar H, Datta S, Agarwal P, Mondal A, Mondal PK, Das S: An Observational Study to Identify Socio-demographic Factors on Primary Caregivers' Quality of Life of Cancer Patients Attending a Tertiary Cancer Hospital, (OA), 122(3): 24-9.
- Gangopadhyay S, Datta A, Nandy K, Dhar H, Jayaseelan P, Agarwal P: Impact of Psychological Intervention on the Quality of Life of Primary Care Givers of Patients with Cancer : Preliminary Report in City based Cancer Hospital, India, (OA), 122(8): 29-35.
- Ganguly S: see Uddin I.

Garg MR: see Reddy KJ.

- Garg R: see Kumar A.
- Gawankar DS, Sarang B, Ruke M, Bulusu S: Indirect Electrothermal Coupling Bowel Injury — A Rare Complication of Laparoscopy, (CR), **122(6)**: 63-4.
- Gayen BK: see Mahapatra U.
- Gengadiya JK, Tamakuwala GP, Vadgama PK, Mangrola JV, Khatri GP: Diagnostic Yield of Bronchoscopy in Benign Lung Diseases at Tertiary Health Care Hospital in South Gujarat, (OA), 122(12): 19-22.
- Ghate S: see Nikam KD.
- Ghosh A: see Mukherjee K.
- Ghosh Dastidar B, Ghosh Dastidar S, Majumdar J, Chakraborty C, Ghosh Dastidar K: Endometrial Perfusion on day of hCG Trigger in IVF Cycles Directly Correlates with the Dynamically Changing Endometrial Thickness : A Pilot Study Examining Potential Implications for Endometrial Receptivity and the Development of an Integrated Model to Assess Endometrial Function, (OA), 122(2): 30-3.
- Ghosh Dastidar K: see Ghosh Dastidar B.
- Ghosh Dastidar S: see Ghosh Dastidar B.
- Ghosh Haldar R: see Chakrabarti Bhattacharyya N.
- Ghosh KU: see Neha N.
- Ghosh S: see Biswas U.
- Ghosh Sengupta S: see Bandyopadhyay P.
- Ghosh UC: see Mukherjee K.
- Gikonyo A: see Owuor H.
- Gikonyo D: see Owuor H.
- Goka S: see Gandham P.
- Golwalkar D, Pandya J: Resolution of Post-COVID-19 Pulmonary Interstitial Pneumonitis with Methylene Blue Nebulization Therapy — A Case Report, (CR), **122(9)**: 71-2.
- Goswami H: see Jain PK.
- Goswami N, Chakraborty N, Kundu D, Chakraborty S, Bhowmick S: A Clinico-Anatomical Study on Primary and Secondary Infertile Females with Special Reference to Radiological Observations, Laparoscopic Findings and Endometrial Histology, (OA), **122(1)**: 32-4.

Govani DR: see Patel RV.

- Goyal A: see Purandare NR.
- Guha Ray T: see Chakrovarty A.
- Gund SV, Dhok AP, Phatak SV, Onkar PM, Mitra K, Trimukhe DM: A Rare Case of Encysted Type of Spermatic Cord Hydrocele in 2-year-old Child, (CR), **122(4)**: 78-9.

Gupta A: see Saboo B.

Gupta A: see Vishwaprem RDR.

Gupta N: see Kattamreddy AR.

Gupta P: see Kour S.

Gupta P: see Soni NR.

Gupta U: see Kour S.

Gupta VK: Challenges in Healthcare Reforms in India, (C), 122(4): 85.

Н

Halder P: see Biswas T.

Halder S, Seikh P, Sarkar A, Chatterjee S, Mallick AK: Assessment of Perceived Stress in Healthcare Professionals Working in a COVID Hospital in West Bengal during COVID-19 Pandemic, (OA), 122(10): 46-8.

Halemani C: see Basarakod SS.

- Haque S: see Chakrabarti Bhattacharyya N.
- Hatkar NK: see Nikam KD.

Hazarika B: see Talukdar AJ.

Herakall M: see .Shekhli S.

Indu RM: see Meenakumari PB.

J

- Jadhav A: see Malani SK
- Jadhav AK, Khan AK, Nalawade DD: Isolated Native Tricuspid Valve Endocarditis with Pulmonary Septic Emboli Presenting as PUO in a Young Adult Female Managed Successfully with Surgical Intervention, (CR), **122(10)**: 76-8.
- Jadhav AK: see Gandhi MA.
- Jadhav BH: see Mahindrakar YS.

Jadhav P: see Babu A.

- Jain P: see Seval M.
- Jain PK, Goswami H, Bhat PK, Seval M: Clinico-epidemiological Profile of Snake Bite Envenomation at a Tertiary Care Centre in Eastern Rajasthan : A Longitudinal Prospective Study, (OA), 122(3): 54-7.
- Jain S: see Purandare NR.

Jambale TA: see Deba Z.

Jatua SK: see Thakur I.

Jaweed SA: see Deba Z.

Jayakrishnan AV: see Ravishankar K.

Jayaram V: see Akhil R.

- Jayaseelan P: see Gangopadhyay S.
- Jha A: see Sannigrahi S.
- Johnson S: see Akhil R.

Johnson S: see Verma P.

Jose A: see Vijay KSS.

Joshi S: see Saboo B.

- Jose C: see Meenakumari PB.
- Joshi A: see Sinha CK.

Κ

Kakkad KM: see Dodiya AJ.

- Kalasker PS: see Deba Z.
- Kalegowda D: see Rathod MP.
- Kalyankar PP: see Kattamreddy AR.
- Kamble N, Kavatagi K, Rangaswamy DR: Should JN 1 Trigger an
- Alarm? (C), **122(6)**: 65.
- Kamble N: see Bhat VG.
- Kamble N: see Rangaswamy DR.

Kant R: see Bohare SM.

Kant S: see Kumar A.

- Kar SK: see Kumar A.
- Karan K: see Chakraborty S.

Karthikeyan N: see Sekar A.

- Kattamreddy AR, Kalyankar PP, Singh Bhalla R, Gupta N: Diagnostic Peritoneal Biopsy in Koch's Abdomen : A Case Report and Review of Literature, (CR), **122(8)**: 70-2.
- Kattamreddy AR: Truth Unveiled : ICMR Guidelines Debunks Common Misconceptions about Umbilical Cord Blood Banking, (C), 122(4): 85-6.
- Kaur J: see Bhatia R.

Kavatagi K: see Kamble N.

- Khade AM: Telemedicine : Current Status & Future Prospects, (RA), 122(6): 55-8.
- Khakholary M: see Purkayastha A.

Khan AK: see Jadhav AK.

- Khan I, Srivastava T, Meena S: Spinal Intramedullary Tubercular Lesions — A Case Series of Clinicoradiological Manifestations, (CS), 122(6): 59-62.
- Khan I: see Chatterjee A.
- Khan Mohd SMS: see Vishwaprem RDR.
- Khandeparkar A: see Shah HK.
- Khare C, Kumar A: Cutaneous Venous Malformation Masquerading as Hematoma in a Newborn, (CR), **122(4)**: 84.
- Khasnabish S: see Trivedi AS.
- Khataminia M, Meshki R, Araban M, Faratseh M: The Relationship between Bruxism in Children and the Psychosocial Status of their Parents, (OA), 122(10): 21-5.
- Khatri GP: see Gengadiya JK.
- Kilikdar M, Saha N, Mohan KR, Rehman Z: Nasal Carriage of Methicillin-resistant Staphylococcus aureus (MRSA) among Health Care Workers Working in Intensive and Critical Care Units at a Teaching Hospital in North India, (OA), 122(1): 21-5.
 Kiran N: see Swamy T.

Kishore KR: see Unnikrishnan RK.

- Kochar A: see Ali Khan N.
- Kochar SK: see Ali Khan N.
- Kotak B: see Neha N.
- Kour S, Singh M, Gupta P, Arora R, Gupta U: Prevalence of Depression and Anxiety among Elderly Patients Visiting Tertiary Care Hospital, (OA), 122(12): 52-6.
- Kouttalmatt DP: see Yadavannavar MC.

Kulkarni SS: see Das NK.

- Kumar A, Verma AK, Bajpai J, Pandey AK, Kant S, Kar SK, Verma SK, Kushwaha RAS, Kumar S, Garg R, Srivastava A, Bajaj DK, Kumar A, Tripathi P: Psychological Distress and Healthrelated Quality of Life in Lung Cancer Patients, (OA), 122(7): 47-51.
- Kumar A: see Chaudhary K.
- Kumar A: see Khare C.
- Kumar A: see Kumar A.
- Kumar EK: see Unnikrishnan RK.
- Kumar L: see Bhavsar HM.
- Kumar MP: see Selvaraju B.
- Kumar N: see Yadav P
- Kumar P, Sambhaji KS: An Observational Study on Neurological Complications among Patients with Dengue Fever, (OA), 122(4): 51-5.
- Kumar P: see Sharma D.
- Kumar P: see Sharma D.
- Kumar S: see Kumar A.
- Kumari PHP: see Panigrahy S.
- Kumbla DK: Hypertension Therapeutics Reimagined : Nebivolol and Telmisartan — A Contemporary Review, (DC), 122(3): 69-74.

Kundu D: see Goswami N.

Kundu S: see Mahapatra U.

Kundu TK: see Chatterjee A.

 Kurien JG: Intravesical Knotting of an Infant Feeding Tube in a Child who Underwent Cardiac Surgery, (C), 122(10): 80.
 Kushwaha RAS: see Kumar A.

L

Lagoo JY: see Acharya GR.

Landge JA: Quality over Quantity in Health Care "Life isn't about quantity, it's about quality" – Malorie Blackman, (C), **122(6)**: 65-6.

Lawrence P: see Selvaraju B.

Lele J: see Ravishankar K.

Lohar R: see Bohare SM.

Luther N: Role of Fetal Kidney Length in Estimation of Gestational Age : Second Trimester *versus* Third Trimester, (OA), **122(6)**: 17-20.

Μ

Madegowda SB: see Rathod MP.

Mahapatra MK: see, Bose SS.

- Mahapatra PC, Pandey A, Mayadeo NM, Dwivedi JS, Vani I, Venkatesh V, Mitra M: Efficacy and Safety of Parenteral Human Placental Extract and Oral Azithromycin Versus Oral Azithromycin Alone in Chronic and Recurrent Pelvic Inflammatory Disease : An Open-Label Randomized Controlled Trial in Indian Patients, (OA), **122(8)**: 58-62.
- Mahapatra U, Bhunia PK, Murmu BK, Kundu S: Scrub Typhus — A Case Series from Tertiary Care Hospital of Rural Bengal, (CS), 122(9): 62-5.
- Mahapatra U, Das Bakshi S, Mukhopadhyay S, Gayen BK: Scrub Typhus with Unilateral Parotitis and Encephalopathy: A Rare Case Report and Literature Review, (CR), **122(12)**: 66-8.

Mahashabde M: see Bhavsar HM.

- Maheta SD: see Panchal PP.
- Mahindrakar YS, Jadhav BH, Salunkhe SN: Defective Systemic Iron Metabolism in Parkinson's Disease, (OA), **122(6)**: 52-4.
- Majumdar A: see Bera S.

Majumdar J: see Ghosh Dastidar B.

Majumdar S: see Thakur I.

Makhal MK: see Thakur I.

- Makwana RA: see Barot MV.
- Malani SK, Nalawade DD, Jadhav A, Wadhokar P: Silent Beats : Spontaneous Recovery of Transient Ventricular Asystole following High Spinal Cord Injury, (CR), **122(5)**: 68-70.

Malhotra N: see Yadav P.

Mallick A: see Sannigrahi S.

Mallick AK: see Halder S.

Mallik S: see Biswas T.

- Mallikarjunappa B: see Vishwaprem RDR.
- Mallikarjunappa B: see Vishwaprem RDR.
- Mandal PK: see Das A.
- Mandal A: see Nandy P.
- Mandal D: see Chakrabarti Bhattacharyya N.
- Mandal M, Chanda A, Roy N: Impending Adrenal Crisis in Exogenous Cushing Following Cam Use, (C), **122(7)**: 85.
- Mangal AK: see Sharma D.
- Mangal AK: see Sharma D.
- Mangala B: see Akhil R.
- Mangrola JV: see Gengadiya JK.
- Manoj R: see Swamy T.
- Masariya B: see Prabhu S.

Mayadeo NM: see Mahapatra PC.

Meena S: see Khan I.

Meena V: see Seval M.

Meenakumari PB, Bhaskar A, Indu RM, Jose C: Clinical Profile, Laboratory Parameters, Radiological Findings of COVID-19 Patients and its Association with Outcome — A Retrospective Observational Study, (RA), 122(11): 50-5.

Meher A: see Mohapatra D.

Mehta AV: see Panchal PP.

Mehta R: see Prabhu S.

Meshki R: see Khataminia M.

Meshram S: see Pate BS.

Mishra B: see Mishra D.

- Mishra D, Turuk J, Palo S, Ray PK, Mishra B, Pati S: Peritoneal Fluid from Pouch of Douglas is not a Suitable Specimen for Molecular Testing in the Diagnosis of Female Genital Tuberculosis in Women Presenting with Infertility, (OA), 122(10): 49-54.
- Mishra GV: see Arora SP.
- Mishra P: see Bhatia R.

Mishra T: see Mohapatra D.

- Mistry J, Patel TR, Patel PM: Functional Outcome of Patients Operated by Mini Open Rotator Cuff Repair in Tertiary Care Hospital of Ahmedabad City, (OA), 122(7): 52-4.
- Mitra K: see Gund SV.

Mitra M: see Mahapatra PC.

Mitra S: see Banerjee S.

Modak D, Chatterjee R: A Study on Risk Factor Assessment, Clinical Profile and Outcome Analysis of Liver Abscess Cases from A Tertiary Care Hospital, Kolkata, (OA), **122(6)**: 47-51.

Modasiya A: see Barbhaya M.

- Mohan KR: see Kilikdar M.
- Mohapatra D, Sahoo PK, Meher A, Behera M, Mishra T: Selfdirected Learning Compared to the Traditional Method of Learning for Undergraduate Medical Students, (OA), 122(10): 60-3.

Moitra S: see Samajdar SS.

Mondal A: see Biswas T. Mondal A: see Gangapadhyay S.

Mondal PK: see Chakraborty B.

- Mondal PK: see Gangapadhyay S.
- Mondal SS: see Saha M.
- Mudey GA: see Pate BS
- Mukherjee K, Ghosh UC, Talukdar A, Ghosh A, Samajdar SS: Presence of Malaria Parasite in Vivax Malaria Patients after Completion of Anti-malaria Therapy in a Tertiary Care Hospital of Kolkata : A Real World Observational Study, (OA), 122(1): 26-8.

Mukherjee R: see Dagra R.

Mukherjee S: see Mukhida S.

Mukherjee S: see Samajdar SS.

Mukhida S, Datta A, Edavaloth P, Pawar V, Mukherjee S: Does only Mupirocin enough for Nasal Decolonization in Methicillinresistant Staphylococcus Aureus Carriers or do we need more? (C), 122(12): 72.

Mukhida S: see Bhavsar HM.

Mukhida S: see Das NK.

Mukhida S: see Verma P

- Mukhopadhyay A: see Patra P.
- Mukhopadhyay A: see Biswas U
- Mukhopadhyay D: see Das SK.
- Mukhopadhyay I: see Sengupta M.
- Mukhopadhyay S: see Mahapatra U.

Mundhada AS: see Mundhada SG.

Mundhada SG, Deogade N, Tanna J, Deshmukh M, Mundhada AS: Sensitivity of SARS-CoV-2 Rapid Antigen Test as Compared

to RT-PCR Test, (OA), **122(7)**: 32-6.

Murmu BK: see Mahapatra U.

Ν

- Nagar GL: see Sharma D.
- Nagar GL: see Sharma D.

Nagar J: see Saboo B.

- Nagendra A. Patil SS: Evaluating Farmers' knowledge, Attitude and Practice Regarding Pesticide Use and Its Impact on Human Health in Northern Karnataka - A Cross Sectional study, (OA), 122(12): 57-60.
- Nair S: see Desai A.

Nair S: see Desai A.

Nair S: see Ruke M.

Nair S: see Ruke M.

Nalawade DD: see Jadhav AK.

- Nalawade DD: see Malani SK.
- Nandy K: see Gangapadhyay S.
- Nandy K: see Gangopadhyay S.
- Nandy P, Rudra O, Chatterjee G, Mandal A: An Unusual Case of Flagellate Pigmentation Following Chemotherapy, (C), 122(11): 82.

Nagvi WM: see Arora SP.

Naskar A: see Biswas T.

Nataraian P: see Sekar A.

- Nath R, Sutradhar D, Thakuria R: A Study of the Clinicodemographic Profile and Clinical Outcomes of COVID-19 Patients in a Tertiary Care Hospital of India, (OA), 122(12): 23-8.
- Nath S: see Chakrovarty A.
- Naveen D: see Vishwaprem RDR.
- Naveen D: see Vishwaprem RDR.
- Nayak A: see Saboo B.

Navak S: see Sahu BK.

- Neha N, Kotak B, Ghosh KU, Shankar AVR, Veligandla KC, Rathod R: Amoxicillin / Clavulanate : A Cornerstone Antibiotic for Treatment of Infections in Paediatrics, (RA), 122(9): 54-61.
- Nikam KD, Hatkar NK, Ghate S: Evaluation of Peak Expiratory Flow Rate (PEFR) in Children Residing in Urban Areas of Dhule District, (OA), 122(10): 26-8.

0

Onkar PM: see Gund SV.

Owuor H, Gikonyo A, Ponoth PM, Gikonyo D: Infiltrative Cardiomyopathy Causing Heart Failure with Compromised Ejection Fraction (HFrEF): An entity to look out for, (CR), **122(7)**: 81-4.

Ρ

Padmaja M: see Sharma A.

Pal A, Biswas B, Shit P, Samajdar SS, Sen S, Tripathi SK: Assessment of the Level of Knowledge and Skills Pertaining to Self-management among Adult Patients of Bronchial Asthma and COPD, (OA), **122(7)**: 70-5.

Pal A: see Biswas A.

Pal J: see Saboo B.

Pal S: see Sengupta A.

Palal D: see Verma P.

Palo S: see Mishra D.

Palsania R: see Singh A.

Panchal DM: see Saini VK.

Panchal PP, Maheta SD, Raval BM, Mehta AV, Shah SR, Shroff DJ, Shah RM: Emergency Obstetrics Hysterectomy : A Retrospective Study in Tertiary Care Hospital in Western India, (OA), 122(7): 17-20.

Panda S: see, Bose SS.

Pandey A: see Mahapatra PC.

Pandey AK: see Kumar A.

Pandya A: see Pandya YG.

Pandya J: see Golwalkar D.

Pandya YG, Pandya A: Exploring the Potential of Artificial Intelligence in Diagnostic Microbiology, (C), 122(1): 61.

Panigrahi C: see Sahu BK.

Panigrahy S, Kumari PHP, Samatha P: Bacteriological Profile and Antibiotic Susceptibility Patterns of Clinical Isolates of Broncho Alveolar Lavage Fluid (BAL) in Patients with Lower Respiratory Tract Infections (LRTIs), (OA), 122(4): 56-60.

Parab S: see Desai A.

Parikh R: see Saboo B.

Parmar H: see Dodiya AJ.

- Parmar RK, Patel MK, Bhatt RJ: A Comparative Study to Estimate Knowledge about Pneumococcal Conjugate Vaccine among Medical and Nursing Students studying at Civil Hospital Campus, Ahmedabad, (OA), 122(5): 33-4.
- Pate BS, Meshram S, Mudey GA, Shaw D: Carbapenem-Resistant Acinetobacter baumannii : Growing Threat to Public Health — A Review of Current Understanding and Future Directions, (RA), 122(4): 73-7.

Patel A: see Barbhaya M.

Patel AA: see Patel SD

Patel AS, Bavaria DR: Association Between T p-e / QT Ratio In 12 Lead ECG and Major Adverse Cardiac Events during Hospital Stay among Acute ST Elevation Myocardial Infarction Patients, (OA), 122(10): 33-8.

Patel DK: see Ravishankar K.

Patel K: see Barbhaya M.

Patel MK: see Parmar RK.

Patel N, Patel R, Patel R: Assessment of Time To Positivity (TTP) and Loading Delay Influencing the Positivity of Blood Culture by an Automated System, (OA), **122(7)**: 37-42.

Patel PM: see Mistry J.

Patel PP: see Saini VK.

Patel R: see Patel N.

Patel R: see Patel N.

Patel R: see Sinha CK.

Patel RR: see Patel RV.

- Patel RV, Correia RC: Laparoscopic Repair of Symptomatic Direct Inguinal Hernia in an Apparently Healthy Boy, (C), 122(10): 82.
- Patel RV, Trambadia RA, Anthony FM, Chhaniara RA, Govani DR, Patel RR, Correia RC: Congenital Neonatal Colonic Atresias Arising in Watershed Areas of the Colonic Blood Supply, (CS), 122(2): 55-60.
- Patel SD, Desai AKA, Patel AA: Prospective Study for Anterior Bridge Plate (ABP) for Humerus Shaft Fracture with Combination Screw : Innovative SPV Technique, (OA), **122(7)**: 64-9.

Patel TR: see Mistry J.

Pati S: see Mishra D.

Patil AP, Yogeshkumar S: Tobacco use among Auto-rickshaw Drivers in Belagavi City, Karnataka, (OA), **122(11)**: 26-31.

Patil SD, Ganganahalli P, Yankanchi S: Knowledge, Attitude and Practices Regarding Tuberculosis among adult Residents of Urban Field Practice Area of Shri B M Patil Medical College, Vijayapura, (OA), 122(6): 43-6.

Patil SS: see Ali SS.

Patil SS: see Nagendra A. Patil V: see Babu A.

Falli V. See Babu A.

Patra P, Dan A, Sarkar S, Mukhopadhyay A, Sanyal D: Extent of Disability and Behavioral Disturbances of the Children with Intellectual Disability and the Coping Mechanisms of their Parents : An Observational Study from a Tertiary Care Hospital of Eastern India. (OA), **122(12)**:35-9.

Pattankar TP: see Yadavannavar MC.

Paul R: Dementia and PET Scan of Brain, (PCME), 122(5): 73.

Paul R: Elephantiasis : A Reappraisal with Special Reference to Nuclear Scan, (PCME), 122(11): 70.

Pawar V: see Mukhida S.

Phalak PJ: see Deshmukh AD.

Phalak PJ: see Pratap A.

Phatak SV: see Bahurupe SS.

Phatak SV: see Gund SV.

Philip AJ: see Sharath R.

- Ponoth PM: see Owuor H.
- Potdukhe PS, Dhok AP: Cervico-vaginal Aplasia : A Rare Anomaly with Review of Literature, (CR), **122(2)**: 61-2.
- Prabhu S, Mehta R, Steward E, Doria G, Bavishi M, Masariya B: Post COVID-19 Rhino-Orbito-Cerebral Mucormycosis : Retrospective Clinical Observational Study & Analysis of the Patients Presenting in Kheda District, Gujarat, India, (OA), 122(5): 14-8.

Prajapati N: see Das SK.

Prajapati P: see Shah SJ.

Prajapati VN: see Dodiya AJ.

- Prasad GH: see Dikshit N.
- Pratap A, Phalak PJ: Solve the Mystery of Biochemistry in Every Square, (C), 122(8): 73-4.
- Purandare NR, Goyal A, Shinde S, Jain S: Comparative Analysis of Sensitivity and Specificity of Creatinine Kinase MB and Cardiac Troponin I for the Diagnosis of Acute Myocardial Infarction, (OA), 122(12): 47-51.
- Purkayastha A, Bhattacharjee P, Khakholary M: Atypical Presentation in a Lady with Acromegaly : A Case Report, (CR), 122(11): 61-5.

R

Rahul RCL: see Sharath R.

Rajagopalan R: Chigger Fever or Cigarette Burn — Unraveling the Mystery of Skin Sore, (C), 122(5): 74.

Rajashree SP: see Basarakod SS.

Rajavelu S: see Dhanaseakaran S.

Rajegowda HM: see Rathod MP.

Rajkumar S: see Sharath R.

Rajwaniya D: see Bhatia R.

- Ramesh XRW, Emperor CE, Dharsini OTS: Interesting Case of Pan facial Fracture, (CR), 122(9): 66-70.
- Rangaswamy DR, Kamble N: Simplifying Day-to-Day Practice with AI, (C), 122(2): 68-9.
- Rangaswamy DR: see Bhat VG.

Rangaswamy DR: see Kamble N.

Rastogi R: see Singh P.

Rathi RK: see Arora SP.

Rathod GH: see Barot MV.

Rathod MP, Rajegowda HM, Madegowda SB, Kalegowda D: A Clinico-epidemiological Study of Dermatophytosis in Children in a Tertiary Health Care in Karnataka, (OA), **122(3)**: 20-3.

Rathod R: see Neha N.

Raval BM: see Panchal PP.

Ravishankar K, Banerjee TK, Srinivasan AV, Patel DK,

Jayakrishnan AV, Lele J, Shah B, Shah R: Headache Algorithm: A Guide to Differential Diagnosis and Management of Headache Disorders by Indian Medical Association, (Commt), 122(11): 71-5.

Ray A: see Saha M.

Ray Chaudhuri G: see Bandyopadhyay P.

Ray Chaudhuri P: see Biswas U.

Ray K: see Das Sarkar M.

Ray PK: see Mishra D.

Ray S: see Biswas U.

Raychaudhuri M: see Banerjee S,

Reddy J: see Reddy KJ.

Reddy KJ, Reddy J, Reddy S, Desai A, Garg MR: Advancements in Vitamin D3 Formulations : A Review of UNS D3 Ultra Nano 60 Thousand, (DC), **122(3)**: 75-8.

Reddy S: see Reddy KJ.

Rehman Z: see Kilikdar M.

Rejoice P: see Unnikrishnan RK.

Rijal P: see Bohare SM

Roy A: see Chakrabarti Bhattacharyya N.

Roy A: see Dagra R.

Roy A: see Sengupta M.

Roy H: see Das Sarkar M.

Roy N: see Mandal M.

Rudra O: see Nandy P.

- Ruke M, Desai A, Anand S, Nair S: A Real-World Evidence Study on Effectiveness and Tolerability of Topical Lincomycin in the Treatment of Surgical Site Infection (SSI) and Skin & Soft Tissue Infection (SSTI), (DC), **122(2)**: 63-6.
- Ruke M, Desai A, Anand S, Nair S: Assessing the Therapeutic Effectiveness and Tolerability of Lincomycin Injectables in Skin and Soft Tissue Infections and Surgical Site Infections : A Comprehensive Real-World Evidence Study, (DC), 122(1): 56-9.
 Ruke M: see Gawankar DS.

S

Saboo B, Parikh R, Gupta A, Vora A, Nayak A, Amin B, Nagar J, Pal J, Joshi S, Bhandari S, Shah V: Unsung Heroes — Ancient Vaidyas to Modern Doctors, (SA), **122(5)**: 71-2.

Sachdeva R: see Chaudhary K.

Saha B: see Biswas T.

Saha M, Sarkar D, Mondal SS, Ray A: A Study on Response of Tenofovir in Chronic Hepatitis B in Eastern India, (OA), 122(9): 17-20

Saha N: see Kilikdar M.

Saha PK: see Sen A.

Saha S: see Dagra R.

Saha S: see Thakur I.

Sahoo PK: see Mohapatra D.

Sahu BK, Nayak S, Panigrahi C: Incidental Cytological Findings of Microfilaria at Unusual Sites with Varied Clinical Presentation : A Retrospective Study, (OA), 122(2): 26-9.

Saikia P: see Talukdar AJ.

Saini VK, Bhabhor MB, Panchal DM, Patel PP, Chaudhari VK, Dagra JJ: An Assessment of Knowledge and Attitude of Beneficiaries Attending Tertiary Care Hospital regarding Temporary Family Planning Methods and to Determine their Experienced Side Effects of Temporary Contraceptive Methods in Ahmedabad City, Gujarat, India. (OA), 122(8): 24-8.

Sakpal A: see Desai A.

Salunkhe SN: see Mahindrakar YS.

Samajdar SS, Mukherjee S, Moitra S, Tripathi SK: Impact of Allergen Immunotherapy on Medication Score and Medicine Usage in Patients with Nasobronchial Allergy : A Single Centre, Prospective Observational Study in a Specialty Clinic in Eastern India, (OA), **122(11)**: 20-5.

- Samajdar SS, Tripath SK: Urgent Need for Dedicated Clinical Pharmacology Departments and to Address Faculty Shortage in West Bengal, (C), 122(8): 73.
- Samajdar SS: see Banerjee S.

Samajdar SS: see Banerjee S.

Samajdar SS: see Banerjee S.

Samajdar SS: see Mukherjee K.

- Samajdar SS: see Pal A.
- Samatha P: see Panigrahy S.
- Sambhaji KS: see Kumar P.

Samsudheen KK: see Yadav PK.

Sannigrahi S, Bandyopadhyay S, Thakur RP, Taraphdar P, Jha A, Mallick A: Perception and Constraints of Telemedicine Services among the Health Care Providers and Beneficiaries in Purba Bardhaman District, West Bengal : A Qualitative Study, (OA), 122(11): 32-7.

Sanyal D: see Patra P.

Sarang B: see Gawankar DS.

Saravanan A: see Shanmugapriya S.

Saravanan T: see Shanmugapriya S.

Sardar S: see Das S.

Sarkar A: see Halder S.

Sarkar D: see Dutta N.

Sarkar D: see Saha M.

Sarkar S: see Patra P.

Sawant G: see Shah HK.

Seikh P: see Halder S.

- Sekar A, Karthikeyan N, Amir AP, Vikram MA, Faizal A, Natarajan P: Assessing the Diagnostic Efficacy of Multi-detector CT Angiography versus Digital Subtraction Angiography in Coronary Artery Disease : A Tertiary Care Centre Study, (OA), 122(7): 58-63.
- Selvaraju B, Kumar MP, Lawrence P: Mortality amongst COVID-19 Patients in Relation to their Vaccination Status, (OA), **122(5)**: 24-7.
- Sen A, Bandyopadhyay T, Bhattacharyya R, Saha PK, Das D: A Comparative Study of Effect of Mindfulness-based Stress Reduction on Psychological Stress & Quality of Life in Patients of Rheumatoid Arthritis with Waitlist Control Group, (OA), **122(5)**: 28-32.
- Sen MK: see Chaudhary K.

Sen S: see Pal A.

Sengupta A, Pal S: Biological Reference Intervals for Hematological Parameters Including Novel Research Parameters in Population of Eastern India, (OA), 122(5): 35-9.

Sengupta A: see Das SK.

- Sengupta M, Baishya K, Roy A, Chakrabarti S, Mukhopadhyay I: An Exploratory Study on Factors Affecting Patient Attitude towards Private Clinical Establishments in India, (OA), 122(4): 43-50.
- Sethuraman S, Adithya S: Invasive Fungal Sinus Diseases with Maxillary Bone Erosion : Is Aggressive Maxillectomy Justified in all Cases ? (OA), **122(4)**: 65-9.
- Seval M, Jain P, Meena V, Verma A: Rifampicin induced Thrombocytopenia : a rare complication, (C), 122(2): 69.

Seval M: see Jain PK.

Shafique A: see Unnikrishnan RK.

Shah B: see Ravishankar K.

Shah HK, Dessai BPS, Bandodkar K, Talaulikar DMS, Khandeparkar A, Cacodcar J, Sawant G, Umraskar L, Galyekar S, Bhagat S: A Cross Sectional Study on Factors Associated with Patients being MIS-3 on Anti-retroviral Therapy Registered at a Nodal Centre — Goa Medical College, (OA), 122(8): 63-9.

Shah JH: see Shah SJ.

Shah R: see Ravishankar K.

Shah RM: see Panchal PP.

Shah SJ, Shah JH, Prajapati P: Prevalence of Diabetic Kidney Disease and Its Associated Risk Factors in Type-2 Diabetes Mellitus — A Tertiary Care Experience, (OA), **122(4)**: 19-25.

Shah SR: see Panchal PP.

Shah SV, Chavada RR, Thakar PM: Evaluation of Alteration in Testicular Perfusion after Laparoscopic Hernioplasty : A Prospective Study Conducted at one of the Tertiary Care Centres of Western India, (OA), 122(10): 70-5.

Shah V: see Saboo B.

Shankar AVR: see Neha N.

Shanmugam B: see Varman M.

- Shanmugapriya S, Saravanan T, Suriya Dharsini J, Shuruthi S, Saravanan A: Drug Utilization Pattern of Proton Pump Inhibitors at a Tertiary Care Hospital in South India, (OA), **122(6)**: 21-7.
- Sharath R, Philip AJ, Rajkumar S, Rahul RCL, Chand A, Umaakanth SCS: Effect of Papain Urea *versus* EUSOL in Diabetic Foot Ulcer Management, (OA), **122(10)**: 64-9.
- Sharma A, Donepudi A, Padmaja M: Statin Intake and Thrombotic Events in Indian Patients — A Cross Sectional Study, (OA), 122(1): 48-50.
- Sharma D, Mangal AK, Fageria N, Kumar P, Nagar GL: To Study Platelet Indices in Alcoholic Liver Disease (ALD) Patients and to Correlate MPV to Platelet Count Ratio with Child Pugh Score to Predict the Severity, (OA), 122(2): 34-8.
- Sharma D, Nagar GL, Fageria N, Mangal AK, Kumar P: Neutrophil to Lymphocyte Ratio (NLR) as an Inflammatory Marker in Predicting Severity in Acute Ischemic Stroke Patients, (OA), 122(12): 40-3.

Sharma J: see Chakraborty S.

 Sharma M: A Study to Evaluate Time Delays and Management Strategy in Patients Presenting with Acute Coronary Syndrome (ACS) in Only Armed Forces Tertiary Cardiac Care Centre in Eastern India during COVID-19 Pandemic, (OA), 122(9): 21-3.
 Sharmili S: see Varman M.

Shaw D: see Pate BS.

Shekhli S, Allichandi R, Herakall M: Prevalence of Depression in Chronic Kidney Disease : An Observational Study, (OA), 122(1): 29-31.

Shekhli S: see Vijay KSS.

- Shinde S: see Purandare NR.
- Shinde SA: see Deshmukh AD.
- Shirsat A: see Trailokya AA.

Shit P: see Pal A.

Shivaprasad VR: see Swamy T.

Shriwastav UK: see Bohare SM.

Shroff DJ: see Panchal PP.

Shuruthi S: see Shanmugapriya S.

Siddhanta S, Bandyopadhyay D, Bhowmick S: Adolescent Health and Screen Time during a Period of Compulsory Online Education : A Cross Sectional Study from Eastern India, (OA), 122(3): 43-9.

Sikarwar S: see Uddin I.

Singh A, Atal R, Choudhary R, Singh Parihar U, Palsania R: A Non-randomized Retrospective Study of Applicability of Clavien-Dindo Classification of Surgical Complications to Emergency Laparotomies, (OA), 122(6): 36-42. Singh A: see Ali Khan N.

Singh Bhalla R: see Kattamreddy AR.

Singh CR: see Devi YE.

Singh M: see Kour S.

Singh NS: see Devi YE.

Singh P, Singh RK, Rastogi R, Gandhi K: A Retrospective Observational Study : To Study the Correlation of Endometrial Thickness versus Endometrial Pattern on Implantation in North Indian Females, (OA), **122(3)**: 30-2.

Singh Parihar U: see Singh A.

Singh R: see Yadav P.

Singh RK: see Singh P.

- Singh S, Srivastava MK: The Clinical Utility of Atypical Lymphocytes, Large Immature Cells, and Rising Hematocrit in Predicting Dengue Fever Severity, (C), **122(1)**: 60.
- Singh S: The Utility of MatriDerm (Skin Substitute) with Autologous Thin Split Skin Graft to Cover Deperiosted Exposed Tibia 6x4 cm Wound in Diabetic Old Man, (RA), **122(7)**: 76-8.
- Sinha CK, Anbarasan R, Patel R, Joshi A, Cherian A: Autoamputated Ovarian Cysts in Infants : Current Status, (CR), 122(5): 62-4.

Sinha Mahapatra TK: see Das S.

Sinha PK: see Dutta N.

- Soni NR, Gupta P : Uncommon Presentation of Adenocarcinoma of Cervix, (CR), 122(1): 54-5.
- Sree M: see Dhanaseakaran S.

Srinivasa BCR: see Vishwaprem RDR.

Srinivasan AV: see Ravishankar K.

Srivastava A: see Kumar A.

Srivastava K: see Akhil R.

- Srivastava MK: see Singh S.
- Srivastava SR: see Das SK.

Srivastava T: see Khan I.

Steward E: see Prabhu S.

Suhana A: see Devi J.

Sural R: see Choudhury U.

- Suriya Dharsini J: see Shanmugapriya S
- Sutradhar D: see Nath R.
- Swamy T, Anitha OR, Kiran N, Shivaprasad VR, Manoj R: Association between BODE Index and Visceral Fat among COPD Patients Attending a Tertiary Care Hospital — A Cross Sectional Study, (OA), 122(7): 21-7.

Swati A: see Chaudhary K.

Т

Taiwo EO: Comparison of Effect of Exercise on Lipid Profiles of Overweight Normoglycemic Offspring of T2DM Parents and Non-diabetic Parents, (OA), 122(11): 15-9.

Talaulikar DMS: see Shah HK.

Talukdar A: see Mukherjee K.

Talukdar AJ, Das RP, Hazarika B, Saikia P, Dutta S, Baishya AC: Use of Remdesivir as an Investigational Therapy in Moderately Severe COVID-19 Cases, (OA), **122(2)**: 15-20.

Tamakuwala GP: see Gengadiya JK.

Tanna J: see Mundhada SG.

Taraphdar P: see Sannigrahi S.

Thakar PM: see Shah SV.

Thakre M: see Asole D.

Thakre M: see Asole D.

Thakur I, Majumdar S, Jatua SK, Makhal MK, Saha S: Are Dengue Patients Aware Enough ? - A Cross-sectional Study among Dengue In-patients in a Rural Based Tertiary Hospital in West Bengal, (OA), 122(2): 39-44.

- Thakur RP: see Sannigrahi S.
- Thakuria R: see Nath R.

Trailokya AA, Shirsat A: Uncontrolled and Resistant Hypertension : A major challenges for Clinicians, (C), 122(2): 67-8.

Trambadia RA: see Patel RV.

Trimukhe DM: see Gund SV.

Tripath SK: see Samajdar SS.

Tripathi P: see Kumar A.

Tripathi SK: see Pal A.

- Tripathi SK: see Samajdar SS.
- Trivedi AS, Chakrabarti D, Khasnabish S, Bhattacharyya AK: High Sensitive CRP and Platelet Volume Indices in Acute Coronary Syndrome and their Impact in Prognostication, (OA), 122(4): 38-42.

Turuk J: see Mishra D.

Ш

Uddin I, Dey S, Sikarwar S, Ganguly S, Anwar AA: Mandibular Osteomyelitis Secondary to Osteopetrosis : A Management Conundrum, (CR), 122(4): 80-3.

Udgiri R: see Ganganahalli PS.

Umaakanth SCS: see Sharath R.

Umraskar L: see Shah HK.

Unnikrishnan RK, Shafigue A, Rejoice P, Kumar EK, Durai G, Kishore KR: Risk Factors for Early Re-bleed following Endoscopic Variceal Band Ligation and Assessing Utility of Dedicated Score (BICAP Score) to Identify High Risk Groups, (OA), 122(8): 36-9.

V

Vadgama PK: see Gengadiya JK. Vajjala SM: see Akhil R.

Vani I: see Mahapatra PC.

Varman M, Shanmugam B, Sharmili S: A Prospective Observational Follow-up Study of Infections in Post Renal Transplant Recipients in a Tertiary Care Hospital in Chennai, (OA), 122(1): 35-41.

Veligandla KC: see Neha N.

Venkatesh V: see Mahapatra PC.

Verma A: see Seval M. Verma A: see Yadav P.



Verma P, Johnson S, Palal D, Mukhida S: Hepatitis and Community: A Roadmap for 2030, (C), 122(9): 77.

Verma SK: see Kumar A.

Vijay KSS, Shekhli S, Jose A: Investigating the Multifaceted Aspects that Affect Interns' Competence and Performance in Arterial Blood Gas Sampling Technique in Emergency Department Settings, (OA), 122(2): 45-8.

Vikram MA: see Sekar A.

- Vishwaprem RDR, Khan Mohd SMS, Naveen D, Srinivasa BCR, Mallikarjunappa B: Cross-sectional Observational Study to Determine Morphology of the Hippocampus in Seizures Disorder by Magnetic Resonance Imaging in a Tertiary Care Hospital, (OA), 122(6): 28-31.
- Vishwaprem RDR, Naveen D, Mallikarjunappa B, Gupta A: Müllerian Duct Fusion Anomaly Associated with Unilateral Renal Agenesis, (CR), 122(7): 79-80.
- Vohra P: An Update on Oral & Systemic Manifestations of GIT Diseases : A Narrative Review, (RA), 122(4): 70-2. Vora A: see Saboo B.

W

Wadhokar P: see Malani SK. Watwe SA: see Das NK. Watwe SA: see Das NK.

Υ

Yadav GE: see Das NK.

- Yadav P, Singh R, Agrawal R, Verma A, Malhotra N, Kumar N: Clinical Outcome of Arabin Cervical Pessary in Women at Risk of Preterm Birth in Indian Scenario, (OA), 122(5): 40-5.
- Yadav PK, Samsudheen KK: A Study on Non-motor Manifestations in Young Onset Parkinsons Disease (YOPD) in Eastern Indian Population, (OA), 122(7): 43-6.

Yadav V: see Ali Khan N.

Yadavannavar MC, Pattankar TP, Kouttalmatt DP: Levels and Risk Factors of Diabetes Mellitus among Tuberculosis Patients Attending District Tuberculosis Centre, Vijayapur. (OA), 122(12): 29-34.

Yankanchi S: see Patil SD. Yogeshkumar S: see Patil AP.

We thank all the Hob'ble Referees who contributed a lot during the year 2024



Dr Mahuya Chattopadhyay Ophthalmology, WB



Medicine, WB





Dr Sujoy Dasgupta Obstetrics and Gynaecology, WB





ology, WB



Dr Pranab Kumar Biswas Obstetrics and Gynaecology, WB



Dr Nilendu Sarma Dermatology & Venereology, WB

W/R Obst



Dr Avijit Roy eral Surgery, , WB



ine & Toxicology, WB Forensic Me



We thank all the Hob'ble Referees who contributed a lot during the year 2024



Dr Abhimanyu Basu Surgery, WB



Dr Anurag Srivastava Surgery, Delhi



Dr Bidisha Ghosh Naskar Radiotherapy, WB



Dr Dilip Karmakar Urology, WB



Dr Hironmoy Roy Anatomy, WB



Dr Kaushik Mitra Community Medicine, WB



Dr Mrinal Kanti Roy Neurology, Bengal



Dr Nilanjan Ghosh Paediatrics, WB



Dr Alakendu Ghosh Rheumatology, WB



Dr Amab Gupta Surgical Oncology, WB



Dr Bitan K Chattopadhyay Surgery, WB



Dr Anirban Dalui Community Medicine, WB,

Dr Dilip Dutta Gynaecologist, WB



Dr Indranil Chakraborty Biochemistry, WB



Dr Kanai Lal Karmakar Nephrology, Bengal



Dr Mukul Ch Biswas Ophthalmology, WB



Dr Manas Banerjee Internal Medicine, Bengal

Dr Niladri Sarkar Medicine, WB



Dr Anita Babasaheb Tandale Dentistry&Endodontics, MS



Dr Arup Das Biswas Cardiology, WB



Dr Biswanath Mukhopadhyay Paediatric Surgery, WB



Dr Dolanchampa Modak Tropical Medicine, WB



Dr Joydeep Deb Chest Medicine, WB



Dr Kaushik Samajdar Physiology, WB



Dr Maitreyee Bhattacharya Haematology, Bengal



Dr Nemai C Nath Surgery, Bengal



Dr Atanu Chandra General Medicine, WB



Dr Ashish Kr Basu Endocrinology, WB



Dr Chandan Chatterjee Pharmacology, WB



Dr D P Singh Chest Medicine, Bihar



Dr Jayashree Paul Senior Consultant, WB



Dr K Mugundhan Neurology, Chenna



Dr Manab Nandy Pharmacology, Bengal



Dr Nilanjan Sinha Haematology, WB



Dr Abhijit Chaudhuri Surgery, WB



Dr Ashish Saha Medicine, WB



Dr Chanchal Kr Jana Medicine, WB



Dr. Girish Mathur Gen.Physician, Rajasthan



Dr KK Mukherjee Orthopaedics, WB



Dr Malay Mundle Community Medicine, WB



Dr Meenakshi Girish Pediatrics, Maharashtra



Dr O P Singh Psychiatry, Bengal



Dr Anup Kr Sadhu Radiology, WB



Dr Asoke Ghoshal Dermatology, Bengal



Dr Debasish Mukherjee ENT Surgeon, WB

Dr. Golokbihari Maji Orthopaedics, WB



Dr Kingshuk K Dhar Gastroenterology, WB



Dr Mausumi De Banerjee Gynaecologist, WB



Dr Mriganka Ghosh General Surgery, Bengal



Dr Praveen Kr Yadav Neurologist, WB

89

We thank all the Hob'ble Referees who contributed a lot during the year 2024



Dr Pradeep Kumar Nemani Surgeon, WB



Dr Rupnarayan Bhattacharya Plastic Surgery, WB



Dr S K Jindal Pulmonology, Chandigarh



Dr Samarendra K Basu Obs. & Gynaecology, WB



Dr Sattik Siddanta Endocronology, WB



Dr Somnath Das FMT, Bengal



Dr Subir Ganguly Radiotherapy, WB









Dr R.N.Sarkar Medicine, Bengal

31

Dr S B Ganguly Medicine, WB



Dr Shamita Chatterjee Surgery, WB





Dr Soma Saha Medicine, Agartala



Dr Soumitra Roy Cardiology, WB



Dr Sujay Maitra Paediatric Surgery, WB



Neurology, Kerala

Dr Udas Ch.Ghosh Medicine, WB



Dr Rana Bhattacharya Endocr.& Metabolism, WB Dr Rajashree Ray Psychiatry, WB



Dr Santanu Kr Tripathi Pharmacology, WB



Dr Shiva K. Misra Surgery, UP



Dr. Soma Dutta, Gynaecologist, WB



Dr Subhankar Home Ophthalmology, WB



Dr Sudip Kumar Das ENT, WB



Dr Gautamananda Roy Stroke Medicine, UK



Dr Pijush Kanti Biswas Medicine, WB



Dr Rudrajit Paul Medicine, WB

Dr Rita Pal Anaesthesiology, WB

Dr Ramdip Ray Liver Transplant, WB

Dr Sandip Ghosh Medicine, WB

Dr Sharmistha Bhattacherjee Community Medicine, WB

Dr Subhash C Biswas Obs. & Gynae, WB

Dr Sudhir Mehta Medicine, Rajasthan

Dr Shohael Md Arafat

Medicine, Dhaka



Dr Reena Ray (Ghosh) Microbiology, WB,



Dr Sanjoy Banerjee Gastroenterology, WB



Dr Shobha Sehgal Immunopathology,Chandigarh



Dr Subhasish K Guha Tropical Medicine, WB



Dr Surya Kant Respiratory Medicine, UP



Dr Partha Ray Neurology, UK



90

Dr Rajdeep Singh Surgery, Delhi



Dr Saikat Datta Medicine, WB



Dr Sanjay Kalra Endocrinology, Haryana



Dr Sarbajit Roy Medicine, Bengal



Dr Santanu Banerjee Orthopaedics, WB



Dr Sukanta Chatterji Paediatrics, WB



Dr Tapan K Naskar Obs.& Gynae, Bengal



Dr Nur Hossain B Sahin Surgery, Bangladesh





The smallest size tablet to combAT BPH

In BPH for long term relief



The first tablet formulation of Tamsulosin in India

In Urinary Tract Infections



In Urinary Tract Infection,



SPREADING PORTRAIT OF SMILLES



INTAS PHARMACEUTICALS LTD. Corporate Office, Near Sola Bridge, S.G.Highway, Thaltej, Ahmedabad-380054, Gujarat, INDIA | www.intaspharma.com



JOURNAL OF THE INDIAN MEDICAL ASSOCIATION :

Sir Nilratan Sircar IMA House, 53, Sir Nilratan Sarkar Sarani (Creek Row), Kolkata - 700 014 Phone : (033) 2237- 8092, Mobile : +919477493027; E-mail : jima1930@rediffmail.com Website : https://onlinejima.com ; www.ima-india.org/ejima Head office : Indian Medical Association, IMA House, Indraprastha Marg, New Delhi - 110 002 Telephones : +91-11-2337 0009, 2337 8680, Email : hsg@ima-india.org : Website : www.ima-india.org

Date of Publication : 20th December, 2024

Registration No. KOL RMS / 476 / 2023 - 2025

RNI Regd. No. 2557/1957 Vol. 68, No. 12, December 2024, Kolkata



If not delivered please return to Journal of the IMA (JIMA) 53, Sir Nilratan Sarkar Sarani, (Creek Row), Kolkata - 700014 Printed and Published by **Dr Sibabrata Banerjee** on behalf of Indian Medical Association and printed at Prabaha, 45, Raja Rammohan Sarani, Kolkata - 700009 and Published from Sir Nilratan Sircar IMA House, 53, Sir Nilratan Sarkar Sarani (Creek Row), Kolkata - 700014, Editor : **Dr Sanjoy Banerjee**